

S0040-4039(96)00031-7

Enantioselective Synthesis of Curacin A. 2. Total Synthesis of Curacin A by Condensation of C1-C7, C8-C17, and C18-C22 Segments

Hisanaka Ito^a, Nobuyuki Imai^a, Ken-ichi Takao^a, and Susumu Kobayashi^{b*}

^a Sagami Chemical Research Center, Nishi-Ohnuma, Sagamihara, Kanagawa 229, Japan

^b Faculty of Pharmaceutical Sciences, Science University of Tokyo, Ichigaya-Funagawara-machi, Shinjuku-ku, Tokyo 162, Japan

Abstract: Total synthesis of curacin A, a novel antimitotic antiproliferative antibiotic, was achieved by the connection of C1-C7, C8-C17, and C18-C22 segments by Julia coupling and iminocther condensation.

Curacin A (1), isolated by Gerwick *et al.* from the marine cyanobacterium Lyngbya majuscula is considered to be one of the interesting antitumor agents.¹ Structural feature and biological activity of curacin A have brought a considerable interest toward the total synthesis and structure-activity relationships.^{2,3} In the preceding paper, we have described an efficient synthetic route to the segments C8-C17 (2), C1-C7 (3), and C18-C22 (4).⁴ In this paper, we report the total synthesis of curacin A (1) by the condensation of three segments.



The coupling of three segments of curacin A was accomplished in a straightforward manner (Scheme 1). Thus, Julia coupling of the sulfone 2 and the aldehyde 3 was first carried out to afford 5 in 50 % overall yield (E : Z = 5 : 1).⁵ Although the stereoselectivity is moderate in the present Julia coupling, the minor isomer could be removed by medium pressure liquid chromatography after conversion to 6. Conversion of thiazolidine 5 into free aminothiol 7 was next examined. Cleavage of the thiazolidine ring under acidic condition⁶ (6N HCl in MeOH reflux) resulted in the formation of complex mixture of products, and was accomplished in a stepwise manner. Initially, Boc group was removed by 6N HCl in MeOH at 55 °C to give 6,⁷ and the resulting thiazolidine 6 was reacted with AgNO3 in CH₃CN-H₂O. Then the silver mercaptide was treated with cysteamine to obtain aminothiol 7. The crude aminothiol 7 thus obtained was finally reacted with iminoether 8, prepared *in situ* from the amide 4 and TfOMe, to complete the total synthesis of curacin A. ¹H, ¹³C NMR spectra of the synthetic curacin A {[α]_D 75° (c 0.44, CHCl₃), lit.¹ [α]_D 86° (c 0.64, CHCl₃)} were in good accordance with those reported by Gerwick *et al.*¹

Scheme 1



Reagents: a) 2 and nBuLi, THF, -78 °C, 20 min, then 3, -78 °C, 30 min, then BzCl, Et₃N, DMAP, r.t., 10 h; b) Na(Hg), MeOH-THF, -12 °C, 2 h, two steps 50 %; c) 6N HCl, MeOH, 55 °C, 2 h; d) AgNO₃, CH₃CN-H₂O, r.t., 1 h, then cysteamine, r.t., 2 h; e) MeOTf, CH₂Cl₂, r.t., 14 h; f) MeOH, 55 °C, 2 days, 32 % from 5.

The present approach can provide a facile route to variously substituted thiazoline derivatives. Structureactivity relationships of the related compounds are now in progress and will be reported in due course.

References and Notes

- 1. Gerwick, W. H.; Proteau, P. J.; Nagle, D. G.; Hamel, E.; Blokhin, A.; Slate, D. L. J. Org. Chem. 1994, 59, 1243-1245.
- 2. Total synthesis of curacin A: White, J. D.; Kim, T.-S.; Nambu, M. J. Am. Chem. Soc. 1995, 117, 5612-5613.
- 3. Synthesis of partial structure of curacin A: Onoda, T.; Shirai, R.; Koiso, Y.; Iwasaki, S. Tetrahedron Lett. 1995, 36, 5765-5768.
- 4. Ito, H.; Imai, N.; Tanikawa, S.; Kobayashi, S. Tetrahedron Lett. the preceding paper.
- 5. Kocienski, P. J.; Lythgoe, B.; Waterhouse, I. J. Chem. Soc. Perkin Trans. 1 1980, 1045-1050.
- 6. Sheehan, J. C.; Yang, D.-D. H. J. Am. Chem. Soc. 1958, 80, 1158-1164.
- 7. 6: $[\alpha]_D^{20}$ -41.2° (c 2.35, CHCl₃); IR (neat) 3401, 2971, 1640, 1439, 1096, and 965 cm⁻¹; ¹H NMR (CDCl₃) δ 6.26 (1H, dd, J = 15.0, 10.8 Hz), 5.75-5.86 (1H, m), 5.81 (1H, d, J = 10.8 Hz), 5.53-5.62 (2H, m), 5.37 (1H, dd, J = 10.8, 8.5 Hz), 5.04-5.12 (2H, m), 4.21 (1H, dddd, J = 9.4, 8.5, 6.0, 0.9 Hz), 3.34 (3H, s), 3.16-3.24 (2H, m), 2.68 (1H, dd, J = 10.3, 9.4 Hz), 2.01-2.33 (8H, m), 1.73 (3H, s), 1.68 (3H, s), 1.56-1.63 (2H, m), 1.57 (3H, s); ¹³C NMR (CDCl₃) δ 136.9, 134.8, 132.8, 131.0, 128.5, 127.4, 124.6, 117.0, 79.9, 75.5, 60.9, 56.6, 43.2, 37.7, 35.4, 33.0, 32.7, 31.6, 31.4, 28.1, 16.6. HRMS calcd for C21H35NOS 349.2437, found 349.2425.

(Received in Japan 11 December 1995; accepted 27 December 1995)