

## SYNTHESIS OF DEUTERATED PENTACENES

B.H. Kwant

Department of Physical Chemistry, University of Groningen,  
Groningen, The Netherlands.

Received February 16, 1977

Revised March 28, 1977

## SUMMARY

Pentacene- $d_{14}$ , pentacene- $d_{12}$  and pentacene- $d_4$  were obtained from 6,13-pentacenequinone- $d_{12}$  and 6,13-pentacenequinone- $d_4$  by refluxing with aluminumcyclohexoxide- $d_{33}$  or aluminumcyclohexoxide. The former quinones were prepared from 1,2-benzenedicarboxaldehyde- $d_6$  or 1,2-benzenedicarboxaldehyde- $d_2$  and 1,4-cyclohexanedione- $d_8$ . The aldehydes were prepared from naphthalene- $d_8$  through ozonization or by reduction of 1,2-benzenedicarboxylic acid dimethylester with  $LiAlD_4$ . All synthetic methods were described in detail.

Key Words: Pentacene, Deuterium, Pentacenequinone, Aluminumcyclohexoxide, Benzenedicarboxaldehyde, Cyclohexanedione.

## INTRODUCTION

In spectroscopic research there is a growing interest in perdeuterated compounds for which there now exist a variety of synthetic methods. This article describes the synthesis of the titled compounds from already perdeuterated chemicals.

Initially the needed perdeuteropentacene was tried to prepare by acid-catalysed exchange of the H analog with  $D_2O$ <sup>(1)</sup> and by the use of an homogeneous catalyst and benzene- $d_6$ <sup>(2)</sup>. Both methods however were without succes. The third synthetic method followed Ried and Anthofer<sup>(3)</sup>, using 1,2-benzenedicarboxaldehyde- $d_6$ <sup>(4)</sup> (I) and 1,4-cyclohexanedione- $d_8$ <sup>(5)</sup> (II), to yield 6,13-pentacenequinone- $d_{12}$  (III) 6,13-Pentacenequinone 5,7,11,14- $d_4$  (VII) was obtained from II and 1,2 benzenedicarboxaldehyde- $d_2$  (VI). The next step followed the method of Bruckner et al<sup>(6,7)</sup> by treatment of III with aluminumcyclohexoxide- $d_{33}$  to yield the desired penta-



mass anal.,  $m/e$  140,  $C_8D_6O_2$ ;  $m/e$  110,  $C_7D_5O$ ;  $m/e$  82,  $C_6D_5$ ;  $m/e$  54,  $C_4D_3$ .

1,4-Cyclohexanedione- $d_8$  (II):

1,4-Cyclohexanedione (5 g, 45 mmol.) was dissolved in an already reacted mixture of  $D_2O$  (30 ml., 1.7 mol.) and  $PCl_5$  (0.5 g, 2.4 mmol.). The solution was stirred 24 hrs at  $50^\circ$ . After cooling, the solid was filtered off and was washed with a small amount  $D_2O$  and the same exchange procedure was repeated twice. The degree of exchange was followed by NMR measurements. After the third exchange, the solid was dried and sublimed two times, yielding 4.0 g white solid, m.p.  $79.6^\circ$ , lit.  $78-80^\circ$ . Ir,  $2125\text{ cm}^{-1}$ ,  $CD_2$ ;  $2940\text{ cm}^{-1}$ ,  $CH_2$  was not found. Anal. calcd. C, 29.97; D, 13.40; found C, 60.01; D, 13.24, mass anal.  $m/e$  120,  $C_6D_8O_2$ ;  $m/e$  60,  $C_4D_6$ .

6,13-Pentacenequinone- $d_{12}$  (III):

I (0.9 g, 6.4 mmol.) and II (0.36 g, 3 mmol.) were dissolved in abs. ethanol (100 ml.) and heated to  $50^\circ$ , thereafter 10% KOH in  $D_2O$  (1.5 ml.) was added to the solution. The mixture was stirred during  $\frac{1}{2}$  hr at  $50^\circ$ , cooled, whereafter the solid was filtered off and washed with a small amount of ethanol and recrystallized from N,N dimethylformamide (DMF), yielding 0.71 g yellow needles. Ir,  $2290\text{ cm}^{-1}$ ,  $CD$ ;  $1675\text{ cm}^{-1}$ ,  $CO$ ;  $2940$  and  $1680\text{ cm}^{-1}$  for CH and CO in the H analog, both were not found. Anal. calcd. C, 82.49; D, 7.52; found C, 82.16; D, 7.68. Mass anal.  $m/e$  320,  $C_{22}D_{12}O_2$ ;  $m/e$  292; 264;  $m/e$  160,  $C_{11}D_6O$ ;  $m/e$  134; 130. Uv, absorbtion max. in benzene, 303 and 402 nm; lit. H analog 300 and 403 nm.

Pentacene- $d_{14}$  (IV):

Aluminum foil (0.05 g, 1.7 mmol.), cyclohexanol- $d_{12}$  (5 ml., 43 mmol.), carbon-tetrachloride (2.5 mg) and  $HgCl_2$  (2.5 mg) were heated under reflux and dry  $N_2$  until the aluminum foil was dissolved. After cooling the mixture, III (0.05 g, 0.16 mmol.) was added under stirring, another 4 hrs refluxed, cooled and thereafter  $D_2O$  (5 ml.) and 37% DCl in  $D_2O$  (3 ml.) were added. The pentacene was filtered off, was washed with  $D_2O$  and dried. Under high vacuum was sublimed, yielding 10 mg deep blue powder. Mass anal.  $m/e$  292,  $C_{22}D_{14}$ ;  $m/e$  146,  $C_{11}D_7$ . Uv, absorbtion max. in pyridine, 495, 534 and 580 nm; lit. 495, 534 and 580 nm for the H analog.

Pentacene 1,2,3,4,5,7,8,9,10,11,12,14-d<sub>12</sub> (V):

Aluminum foil (0.116 g, 4.3 mmol.), cyclohexanol (5 ml., 48 mmol.), carbontetrachloride (0.05 g) and HgCl<sub>2</sub> (3 mg) were heated under reflux and dry N<sub>2</sub> until the aluminum foil was dissolved. After cooling the mixture, III (0.116 g, 0.36 mmol.) was added and then for 16 hrs heated at 200°. The mixture was then cooled water (5 ml.) and 37% HCl (3 ml.) were added, further diluted with water (95 ml.) and the blue mixture filtered off, washed with water and ethanol and dried. The powder was sublimed under high vacuum, yielding 40 mg deep blue product. Ir, 3060 cm<sup>-1</sup>, CH; 2280 cm<sup>-1</sup>, CD. Mass anal. m/e 290, C<sub>22</sub>D<sub>12</sub>H<sub>2</sub>; m/e 145, C<sub>11</sub>D<sub>6</sub>H. Uv, absorption max. in pyridine, gave the same spectrum as IV.

1,2-Benzenedicarboxaldehyde-d<sub>2</sub> (VI):

IX (4.0 g, 27.8 mmol.) and SeO<sub>2</sub> (3.2 g, 28.8 mmol.) were slowly heated to 150° and the water was distilled off. The dry residue was heated in a bulb-to-bulb distillation apparatus above 200°, the distilled product was taken up in diethyl ether and was dried. The solvent was removed and the residue was sublimed, yielding 1.4 g light-yellow crystals. M.p. 55.0°; lit. 53.2°; 56-57° for the H analog. Ir, 2130 cm<sup>-1</sup>, CD; 1675 cm<sup>-1</sup>, CD. Mass anal. m/e 136, C<sub>8</sub>H<sub>4</sub>D<sub>2</sub>O<sub>2</sub>; m/e 106, C<sub>7</sub>H<sub>4</sub>DO; m/e 78, C<sub>6</sub>H<sub>4</sub>D; m/e 51, C<sub>4</sub>H<sub>3</sub>; a small amount of m/e 148, C<sub>8</sub>H<sub>4</sub>O<sub>3</sub> was also found.

6,13-Pentacenequinone 5,7,12,14-d<sub>4</sub> (VII):

VI (1 g, 7.4 mmol.) and II (0.4 g, 3.3 mmol.) were dissolved in abs. ethanol (125 ml.) and heated to 50-60° and thereafter 10% KOH in water (1.5 ml.) was added to the solution. The mixture was stirred 1 hr at 60°, cooled and the solid was filtered off, washed with ethanol and recrystallized from hot DMF. The fine yellow needles so obtained were washed with diethyl ether and dried, yielding 0.83 g. Ir, 3070 cm<sup>-1</sup>, CH; the CD peak was not visible; 1675 cm<sup>-1</sup>, CD. Uv, absorption max. in benzene, 302.5 and 402 nm, see also the uv spectra of compound III. Mass anal. m/e 312, C<sub>22</sub>D<sub>4</sub>H<sub>8</sub>O<sub>2</sub>; m/e 284; 256; m/e 156, C<sub>11</sub>D<sub>2</sub>H<sub>4</sub>O; m/e 128 and m/e 127.

Pentacene 5,7,12,14-d<sub>4</sub> (VIII):

Aluminum foil (0.09 g, 3.3 mmol.), cyclohexanol (1.8 ml., 17.3 mmol.), bromine (0.01 g) were heated at 200° under dry N<sub>2</sub>. The reaction was very slow and after

24 hrs there was still aluminum foil left. Then there was added  $\text{HgCl}_2$  (0.01 g), whereafter the reaction started. After all the aluminum foil had dissolved, the mixture was cooled and VII (0.09 g, 0.3 mmol.) was added. The blue mixture was heated during 48 hrs at  $200^\circ$ , cooled, the product was filtered off, washed with water and benzene and dried. The blue product was sublimed under high vacuum, yielding 25 mg deep blue powder. Ir,  $3060\text{ cm}^{-1}$ , CH;  $2280\text{ cm}^{-1}$ , CD;  $1675\text{ cm}^{-1}$ , CD was missing. Uv, absorbtion max. in pyridine, as product IV. Mass anal. m/e  $282, \text{C}_{22}\text{D}_4\text{H}_{10}$ ; m/e  $281, \text{C}_{22}\text{D}_3\text{H}_{11}$ ; m/e  $280, \text{C}_{22}\text{D}_2\text{H}_{12}$ ; m/e  $279, \text{C}_{22}\text{D}\text{H}_{13}$ ; m/e  $278, \text{C}_{22}\text{H}_{14}$ ; all five derivatives in roughly 20% concentration; m/e  $141, \text{C}_{11}\text{D}_2\text{H}_5$ ; m/e  $140, \text{C}_{11}\text{D}\text{H}_6$ ; m/e  $139, \text{C}_{11}\text{H}_7$ , all three derivatives in roughly 33% concentration.

1,2-Bis(hydroxymethyl-d<sub>2</sub>) benzene (IX):

1,2-Benzenedicarboxylic acid dimethylester (10 g, 52 mmol.) in diethyl ether (250 ml.) and tetrahydrofuran (50 ml.) (all three chemicals dried before usage) was reduced with  $\text{LiAlD}_4$  (2.65 g, 63 mmol.), refluxed 1 hr and left overnight at roomtemperature. The mixture was cooled at  $0^\circ$ , whereafter  $\text{D}_2\text{O}$  (2.6 ml.), 15% NaOH in  $\text{D}_2\text{O}$  (2.6 ml.) and  $\text{D}_2\text{O}$  (7.5 ml.) were added in this order. (By this method<sup>(8)</sup> a dry granular precipitate was obtained) The precipitate was filtered off, washed with diethyl ether, the upper layer was separated, dried, filtered and the solvent removed. After distillation of the residue, b.p.  $120^\circ/0.01\text{ mm}$ , 4.3 g white solid was obtained, m. p.  $63.2^\circ$ , lit.  $64.2-64.8^\circ$  for the H analog. Ir,  $1735\text{ cm}^{-1}$ , CD was missing. Without further purification this was used for the preparation of 1,2-benzenedicarboxaldehyde-d<sub>2</sub> (VI).

ACKNOWLEDGEMENTS

The author wishes to thank Mr. A. Kievit and Mr. A. de Lange for measuring the mass spectra and the staff of the analytical department for the C H measurements.

REFERENCES

1. Brown W.G. and Garnett J.L. -J. Amer. Chem. Soc. 80: 5272 (1958)
2. Garnett J.L., Long M.A., Vining R.F.W. and Mole T. -J. Amer. Chem. Soc. 94: 5913 (1972)
3. Ried W. und Anthöfer F. -Angew. Chem. 65: 601 (1953)
4. B.H. Kwant, previous article in this journal.
5. Wood G., Woo E.P. and Miskow M.H. -Can. J. of Chem. 47: 429 (1969)
6. Bruckner V. und Tomasz J. -Acta Chim. Hung. Tomus 28: 405 (1961)
7. Bruckner V., Karczag (Wilhelms) A., Körmeny K., Meszaros M. und Tomasz J. -Tetrahedron Letters 1: 5 (1960)
8. Mićović V.M. and Mihailović M. LJ. -J. Org. Chem. 18: 1190 (1953)