

Hydrazine-Selective Chromogenic and Fluorogenic Probe Based on Levulinated Coumarin


Myung Gil Choi, Jiyoung Hwang, Jung Ok Moon, Jaeyoung Sung, and Suk-Kyu Chang*

Department of Chemistry, Chung-Ang University, Seoul 156-756, Korea

skchang@cau.ac.kr

Received August 6, 2011

ABSTRACT



Chemosignaling of hydrazine by selective deprotection of levulinated coumarin was investigated. In the presence of hydrazine, levulinated coumarin was selectively deprotected, resulting in chromogenic and fluorescent turn-on type signaling. The selective naked-eye detectable signaling of hydrazine was possible in the presence of representative metal ions and common anions in an aqueous environment.

Hydrazine is widely used as a fuel in rocket and missile propulsion systems¹ and as a reactant in fuel cells.² It is a highly reactive base and a strong reducing agent and is used as an important reactant in the preparation of pharmaceuticals, pesticides, photography chemicals, emulsifiers, and dyes in various chemical industries.³ In industry it is often applied as a chemical blowing agent and corrosion inhibitor for heating systems.⁴ Hydrazine, however, is highly toxic and readily absorbed by oral, dermal, or

inhalation routes of exposure. Long-term studies with laboratory animals indicate that hydrazine is mutagenic and carcinogenic.⁵

Due to its widespread applications and human toxicity, developing reliable and sensitive analytical methods for the selective detection of hydrazine is highly desirable. Hydrazine can be routinely analyzed by a wide variety of chromatographic techniques, such as gas chromatography, high performance liquid chromatography, and ion chromatography.⁶ Electrochemical detection using a variety of chemically modified electrodes has also frequently been used.⁷ However, there are few reports of optical analysis systems, despite their convenience. Oxidation of

(1) (a) Hydrazine and Its Derivatives. In *Kirk-Othmer Encyclopedia of Chemical Technology*, 5th ed.; Kroschwitz, J. I., Seidel, A., Eds.; Wiley: New York, 2005; Vol. 13, pp 562–607. (b) Mo, J.-W.; Ogorevc, B.; Zhang, X.; Pihlar, B. *Electroanalysis* **2000**, *12*, 48–54. (c) Zelnick, S. D.; Mattie, D. R.; Stepaniak, P. C. *Aviat. Space Environ. Med.* **2003**, *74*, 1285–1291.

(2) Serov, A.; Kwak, C. *Appl. Catal., B* **2010**, *98*, 1–9.

(3) (a) Ragnarsson, U. *Chem. Soc. Rev.* **2001**, *30*, 205–213. (b) Garrod, S.; Bollard, M. E.; Nicholls, A. W.; Connor, S. C.; Connelly, J.; Nicholson, J. K.; Holmes, E. *Chem. Res. Toxicol.* **2005**, *18*, 115–122. (c) Narayanan, S. S.; Scholz, F. *Electroanalysis* **1999**, *11*, 465–469.

(4) Vieira, I. C.; Lupetti, K. O.; Fatibello-Filho, O. *Anal. Lett.* **2002**, *35*, 2221–2231.

(5) International Agency for Research on Cancer: Re-evaluation of some organic chemicals, hydrazine, and hydrogen peroxide. IARC monographs on the evaluation of carcinogenic risk of chemicals to humans. Lyon, IARC, 1999, Vol. 71, pp 991–1013. <http://monographs.iarc.fr/ENG/Monographs/vol71/mono71-43.pdf>.

(6) (a) Elder, D. P.; Snodin, D.; Teasdale, A. *J. Pharm. Biomed. Anal.* **2011**, *54*, 900–910. (b) Haghani, A.; Eaton, A.; Wan, J.; Cha, Y. Y. *Proceedings-Water Quality Technology Conference and Exposition 2008*, hagh1/1–hagh1/17.

(7) (a) Batchelor-McAuley, C.; Banks, C. E.; Simm, A. O.; Jones, T. G. J.; Compton, R. G. *Analyst* **2006**, *131*, 106–110. (b) Umar, A.; Rahman, M. M.; Kim, S. H.; Hahn, Y.-B. *Chem. Commun.* **2008**, 166–168.

(8) Ensafi, A. A.; Rezaei, B. *Talanta* **1998**, *47*, 645–649.

(9) (a) Collins, G. E.; Rose-Pehrsson, S. L. *Anal. Chim. Acta* **1993**, *284*, 207–215. (b) Collins, G. E.; Rose-Pehrsson, S. L. *Analyst* **1994**, *119*, 1907–1913.


hydrazine by Ti(III) to produce fluorescent Ti(I)⁸ and reaction with arenedicarboxaldehydes to produce fluorescent derivatives have been reported.⁹ Fluorescence signaling with disruption of the internal hydrogen bonding by hydrazine in a carbazolopyridinophane was also reported.¹⁰ Swager et al. reported interesting trace hydrazine detection results using a turn-on type fluorescent conjugated polymer film.¹¹

Among the many sophisticated optical signaling systems, reactive chemical probes have received much interest due to their specificity and cumulative signaling effects.¹² There are many smart reactive probe systems for the analysis of metal ions, anions, and neutral molecules utilizing unique chemical transformations.¹³ Particularly, deprotection of specific protecting groups has been used as a versatile signaling tool, for example the silyl ether for fluoride,¹⁴ the boronate for hydrogen peroxide,¹⁵ the benzenesulfonyl group for superoxide,¹⁶ the hydrazone for Cu²⁺,¹⁷ and the thioacetals for Hg²⁺.¹⁸

Levulinoyl ester, a versatile protecting group often applied in synthetic organic chemistry,¹⁹ can be removed selectively by treatment with hydrazine.²⁰ Interestingly, levulinated triphenol has been used as a hydrazine cleavable chromogenic protecting group for hydroxyl groups.²¹ Recently, we reported that levulinated resorufin could be used as a selective chromogenic probe for the determination of sulfites.²² With this background for levulinates, we devised a new hydrazine-selective signaling system. A levulinate of 3-cyano-7-hydroxycoumarin was successfully deprotected under mild conditions and acted as a selective chromogenic and fluorogenic probe for hydrazine.²³

Levulinate **1** was prepared by the reaction of 3-cyano-7-hydroxycoumarin **2** with levulinyl chloride (65%, CH₂Cl₂) (Scheme 1).

Scheme 1. Preparation of Levulinate Probe **1**



The chromogenic signaling behavior of 3-cyano-7-hydroxycoumarin levulinate **1** was investigated in a 30% aqueous DMSO solution at pH 4.5 (acetate buffer, 10 mM). Levulinate **1** revealed moderate UV–vis absorption at 307 and 336 nm. Upon the interaction of **1** with hydrazine, a prominent absorption band centered at 426 nm developed (Figure 1). Concomitantly, a greenish yellow color, which is a characteristic of **2**, developed that allowed a colorimetric detection of hydrazine by the naked eye. The changes in absorption bands by the hydrazine-induced deprotection process were remarkable and provided ratio-metric analysis for the transformation of probe **1** to **2**. With 100 equiv of hydrazine, the absorbance ratio A_{426}/A_{336} at the two characteristic wavelengths of 426 and 336 nm increased over 500-fold (from 0.014 to 7.14). Other common cations and anions were relatively nonresponsive, with A_{426}/A_{336} values varying in a limited range between 0.019 (for Pb²⁺) and 0.078 (for Fe³⁺) for metal ions, 0.014 (for Br⁻) and 0.16 (for N₃⁻) for anions, respectively (Figures S1 and S2, Supporting Information).

(10) Brown, A. B.; Gibson, T. L.; Baum, J. C.; Ren, T.; Smith, T. M. *Sens. Actuators, B* **2005**, *110*, 8–12.

(11) Thomas, S. W., III; Swager, T. M. *Adv. Mater.* **2006**, *18*, 1047–1050.

(12) Jun, M. E.; Roy, B.; Ahn, K. H. *Chem. Commun.* **2011**, *47*, 7583–7601.

(13) (a) Mohr, G. J. *Anal. Bioanal. Chem.* **2006**, *386*, 1201–1214. (b) Cho, D.-G.; Sessler, J. L. *Chem. Soc. Rev.* **2009**, *38*, 1647–1662.

(14) (a) Kim, S. Y.; Hong, J.-I. *Org. Lett.* **2007**, *9*, 3109–3112. (b) Bhalla, V.; Singh, H.; Kumar, M. *Org. Lett.* **2010**, *12*, 628–631.

(15) (a) Chang, M. C. Y.; Pralle, A.; Isacoff, E. Y.; Chang, C. J. *J. Am. Chem. Soc.* **2004**, *126*, 15392–15393. (b) Dickinson, B. C.; Chang, C. J. *J. Am. Chem. Soc.* **2008**, *130*, 9638–9639.

(16) Maeda, H.; Yamamoto, K.; Kohno, I.; Hafsi, L.; Itoh, N.; Nakagawa, S.; Kanagawa, N.; Suzuki, K.; Uno, T. *Chem.—Eur. J.* **2007**, *13*, 1946–1954.

(17) Kim, M. H.; Jang, H. H.; Yi, S.; Chang, S.-K.; Han, M. S. *Chem. Commun.* **2009**, 4838–4840.

(18) Cheng, X.; Li, Q.; Qin, J.; Li, Z. *ACS Appl. Mater. Interfaces* **2010**, *2*, 1066–1072.

(19) Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 3rd ed.; John Wiley & Sons: New York, 1999; pp 168 and 278.

(20) Geurink, P. P.; Florea, B. I.; Li, N.; Witte, M. D.; Verasdonck, J.; Kuo, C.-L.; van der Marel, G. A.; Overkleeft, H. S. *Angew. Chem., Int. Ed.* **2010**, *49*, 6802–6805.

(21) Leikauf, E.; Köster, H. *Tetrahedron* **1995**, *51*, 5557–5562.

(22) Choi, M. G.; Hwang, J.; Eor, S.; Chang, S.-K. *Org. Lett.* **2010**, *12*, 5624–5627.

(23) We tried to apply this approach to another widely used phenolic dye resorufin. The resorufin derivative also exhibited satisfactory signaling behavior toward hydrazine; however, the interference from azide ions was significant (Figure S11, Supporting Information). On the other hand, the 7-hydroxycoumarin derivative revealed only turn-on type fluorescence signaling without any chromogenic behavior, which is inferior to the present system. Zhou, Z.; Li, N.; Tong, A. *Anal. Chim. Acta* **2011**, *702*, 82–86.




Figure 1. UV–vis spectra of probe **1** in the presence of hydrazine, representative metal ions, and anions. [**1**] = 1.0×10^{-5} M, [hydrazine] = [M^{n+}] = [A^{n-}] = 1.0×10^{-3} M in a mixture of acetate buffer (pH 4.5, 10 mM) and DMSO (3:7, v/v). Measured after 15 min of mixing.

The fluorogenic signaling behavior of **1** toward hydrazine was measured. Probe **1** demonstrated very weak fluorescence emission at 458 nm (Figure 2). Upon interaction with hydrazine, the fluorescence intensity at 458 nm increased 250-fold and the solution color, under illumination

with a UV lamp, changed from colorless to bright blue. The other metal ions and anions exhibited almost no changes in emission behavior; the emission intensity ratio in the presence and absence of various ions at 458 nm, I/I_0 , varied within a narrow range between 0.81 (Zn^{2+}) and 1.05 (K^+) for metal ions, 1.07 (HCO_3^-) and 2.37 (N_3^-) for anions, respectively (Figures S3 and S4, Supporting Information).





Figure 2. Fluorescence spectra of **1** in the presence of hydrazine, representative metal ions, and anions. $[1] = 5.0 \times 10^{-6}$ M, $[hydrazine] = [M^{n+}] = [A^{n-}] = 5.0 \times 10^{-4}$ M in a mixture of acetate buffer (pH 4.5, 10 mM) and DMSO (3:7, v/v). Measured after 15 min of mixing. $\lambda_{ex} = 360$ nm.

Signaling occurs due to the selective deprotection of the levulinate group of probe **1** by hydrazine (Scheme 2). Cleavage of the levulinate group of **1** by reaction with hydrazine is known to proceed first to the carbonyl group at the 4-position of the levulinate group and then to subsequent amide ring formation leading to cleavage of the ester function.²⁴ Thus generated 3-cyano-7-hydroxycoumarin exhibited its characteristic chromogenic and fluorogenic signaling behaviors.

Scheme 2. Signaling of Hydrazine by Levulinated Hydroxycoumarin **1**



The proposed hydrazine induced deprotection was evidenced by 1H NMR and UV-vis measurements. By 1H NMR spectroscopy, resonances for the coumarin moiety

(24) Ho, T. L.; Wong, C. M. *Synthesis* **1974**, *45*, 45–45.

of probe **1** were observed at 8.68, 7.79, 7.23, and 7.15 ppm. However, the 1H NMR spectrum of probe **1** upon treatment with 2 equiv of hydrazine was almost identical to that of 3-cyano-7-hydroxycoumarin (Figure 3). In addition, the resonances of the reaction product 4,5-dihydro-6-methylpyridazin-3(2*H*)-one was also observed (Figure S5, Supporting Information). The UV-vis spectrum of probe **1** in the presence of hydrazine was also almost identical to that of the proposed deprotection product of 3-cyano-7-hydroxycoumarin itself (Figure S6, Supporting Information).




Figure 3. Partial 1H NMR spectra of **1** only, **1** + hydrazine, and **2** + hydrazine. $[1] = [2] = 5.0 \times 10^{-3}$ M, $[hydrazine] = 1.0 \times 10^{-2}$ M in $D_2O/DMSO-d_6$ (3:7, v/v).

Competition experiments on the signaling of the **1**–hydrazine system revealed that hydrazine-induced UV-vis changes of **1** were not significantly altered in the presence of 100 equiv of coexisting common anions and metal ions except for Cu^{2+} and Hg^{2+} (Figures S7 and S8, Supporting Information). The interference from Cu^{2+} and Hg^{2+} ions was successfully circumvented by using a chelating resin Chelex-100 (Figure S9, Supporting Information). Signaling of hydrazine was finished in less than 15 min, while probe **1** showed no responses at all under the measurement conditions (Figure S10, Supporting Information).




Figure 4. UV-vis titration of **1** with hydrazine. $[1] = 1.0 \times 10^{-5}$ M in a mixture of acetate buffer (pH 4.5, 10 mM) and DMSO (3:7, v/v). Measured after 15 min of each mixing.

Finally, the quantitative analytical behavior of probe **1** for the analysis of hydrazine was investigated by UV–vis absorption titration. In spectra taken 15 min after the addition of hydrazine to **1**, the absorbance ratio (A_{426}/A_{336}) increased steadily in response to the increases in hydrazine to about 14 equiv of titration (Figure 4). From the concentration-dependent UV–vis absorption changes, the detection limit of probe **1** for the determination of hydrazine was estimated to be $2.46 \mu\text{M}$ (0.08 ppm) in a 30% aqueous DMSO solution.²⁵

In summary, we have developed a new chemosignaling system for the selective sensing of hydrazine by deprotection of the levulinate group. The probe revealed a selective

(25) Shortreed, M.; Kopelman, R.; Kuhn, M.; Hoyland, B. *Anal. Chem.* **1996**, *68*, 1414–1418.

chromogenic and fluorogenic signaling behavior in response to hydrazine in a 30% aqueous DMSO solution. Selective signaling is based on the efficient deprotection of the levulinate group by hydrazine. The developed system could be used as a convenient signaling tool for the optical determination of hydrazine in an aqueous environment.

Acknowledgment. This work was supported by a fund from the Korea Research Foundation of the Korean government (2011-0004870).

Supporting Information Available. Experimental details, NMR spectra, and additional chemosignaling behavior of **1** are reported. This material is available free of charge via the Internet at <http://pubs.acs.org>.