

## SELECTIVE OXIDATION METHODS FOR PREPARATION OF N-ALKYLPHENOTHIAZINE SULFOXIDES AND SULFONES

Monica Toşa<sup>1</sup>, Csaba Paizs<sup>1</sup>, Cornelia Majdik<sup>1</sup>, László Poppe<sup>2</sup>, Pál Kolonits<sup>2</sup>, Ioan A. Silberg<sup>3</sup>,  
Lajos Novák<sup>2</sup>, Florin-Dan Irimie<sup>\*1</sup>

<sup>1</sup> Department of Biochemistry and Biochemical Engineering, Babes-Bolyai University, RO-3400, Cluj-Napoca, Arany János 11, Romania

<sup>2</sup> Institute for Organic Chemistry and Research Group for Alkaloid Chemistry of the Hungarian Academy of Sciences, Budapest University of Technology and Economics, H-1111 Budapest, Gellért tér 4., Hungary

<sup>3</sup> Department of Organic Chemistry, Babes-Bolyai University, RO-3400, Cluj-Napoca, Arany János 11, Romania

**Abstract:** Efficient and selective oxidation methods for preparation of *N*-alkylphenothiazine sulfoxides **2a-h** and sulfones **3a-h** starting from *N*-alkylphenothiazines **1a-h** are described.


In the metabolism of the phenothiazine-based drugs often occurring metabolites are the corresponding sulfoxides, 7-hydroxylated derivatives or even sulfones (1-4).

Phenothiazines have been oxidized by various oxidizing agents. Most frequently H<sub>2</sub>O<sub>2</sub> in various solvents has been used for *S*-oxidation (1). In ethanol (5) or in ethanol-acetone mixture (6), H<sub>2</sub>O<sub>2</sub> produced phenothiazine-5-oxides. Sulfones were usually formed by H<sub>2</sub>O<sub>2</sub>/AcOH (7) which generates *in situ* peracetic acid. Similarly, oxidation with organic peracids, such as *m*-chloroperoxybenzoic acid (8), provided sulfones. Sulfone formation was also achieved with sodium perborate (9) or NaOCl (10). Sulfoxide formation by peracid systems was only reported for ammonium persulfate (11) or H<sub>2</sub>O<sub>2</sub>/oxalic acid (12). Transformation of phenothiazine derivatives into sulfoxides was achieved by several inorganic nitrogen oxides. Phenothiazines substituted with electron withdrawing groups were converted to sulfoxides by nitric acid (1). Sulfoxides were obtained by using nitrous acid generated *in situ* from NaNO<sub>2</sub> by acids like AcOH (13) or aqueous HCl (14). Phenothiazines were converted into sulfoxides by nitrogen dioxide/O<sub>2</sub> (15) or NO<sup>(+)</sup>BF<sub>4</sub><sup>(-)</sup>/O<sub>2</sub> (16). Heavy metal-based systems were also used for *S*-oxidation of phenothiazines. KMnO<sub>4</sub>/H<sub>2</sub>O (17), Et<sub>3</sub>PhN<sup>(+)</sup>MnO<sub>4</sub><sup>(-)</sup> in organic

\* To whom correspondence should be addressed.

media (18), or  $\text{OsO}_4$ /*N*-methylmorpholine *N*-oxide (19) were applied for oxidation of phenothiazines into sulfones. On the contrary, the milder chromium based oxidizing agents like  $\text{CrO}_3$ /pyridine (20), resulted in sulfoxide formation. Ozone (21), electrochemical methods (22), tetramethyl oxirane (23) or perfluoroalkyl-oxaziridines (24) were also used for oxidizing phenothiazines to sulfoxides.

All of these methods, however, required either expensive/unusual reagents, or afforded the desired products only in moderate yields *via* tedious purification. In this work, new methods applying convenient oxidizing agents, requiring mild conditions and simple work up and providing good yields for selective synthesis of phenothiazine-5-oxides **2a-h** or 5,5-dioxides **3a-h** from 10-alkylphenothiazines **1a-h** are reported (Scheme 1).



Scheme 1. Preparation and selective oxidations of 10-alkylphenothiazines **1a-h**

## Experimental

Phenothiazine and reagents: Aldrich or Fluka.  $^1\text{H}$ -/ $^{13}\text{C}$ -NMR:  $\text{CDCl}_3$ /TMS, Bruker DRX-500 equipment at 500/125 MHz, ppm ( $\delta$ ). IR: Specord 2000 spectrometer, KBr ( $\text{cm}^{-1}$ ). TLC: Merck Kieselgel 60 F<sub>254</sub> alumina sheets. Vacuum-chromatography: Merck Kieselgel 60 (0.063-0.200  $\mu\text{m}$ ). M.P.: uncorrected. Solvents: purified and dried by standard methods. All new compounds gave proper analysis data.

### I. General procedure for alkylation of phenothiazine with alkyl iodides

Phenothiazine (20 g, 100 mmol) was suspended in 100 ml alkyl iodide and  $\text{NaNH}_2$  (6 g, 150 mmol, 95 %) powder was added portionwise and the reaction was stirred at reflux temperature. After the reaction was completed (by TLC), the excess of alkyl iodide was distilled off in vacuo and unreacted  $\text{NaNH}_2$  was quenched (water, 50 ml). Followed an extraction with toluene (50 ml, twice), drying ( $\text{MgSO}_4$ ) and removal of solvent, the residual crude 10-alkylphenothiazines **1a-h** were purified by vacuum-distillation for **1c-h** and by recrystallization from ethanol for **1a-b** (Table 1).

Table 1. Preparation of 10-alkylphenothiazines **1a-h**

| Compnd.                | T<br>[°C] | Time<br>[h] | Yield<br>[%] | M.p./lit. M.p.<br>[°C] | B.p. /lit. B.p.<br>[°C; mm Hg] |
|------------------------|-----------|-------------|--------------|------------------------|--------------------------------|
| <b>1a</b>              | 43-44     | 12          | 91           | 100 /100 (25)          |                                |
| <b>1b</b>              | 72-73     | 8           | 95           | 106 /103-4 (26)        |                                |
| <b>1c</b>              | 102       | 7           | 96           | 50 /49-50 (26)         | 164;0.05 /162-5;0.02 (27)      |
| <b>1d</b>              | 131       | 7           | 95           |                        | 145;1 /143-6;0.1 (28)          |
| <b>1e</b>              | 121       | 6           | 96           | 128 /127-9 (28)        | 156;1                          |
| <b>1f</b>              | 148       | 7           | 96           | 67 /66-8 (33)          | 169;0.5 /230-2;13 (33)         |
| <b>1g</b> <sup>1</sup> | 156       | 5           | 94           |                        | 166;0.07                       |
| <b>1h</b> <sup>2</sup> | 205       | 5           | 95           |                        | 172;0.05                       |

<sup>1</sup>**1g**: <sup>1</sup>H-NMR: 0.96 (3H, t), 1.33-1.47 (m, 4H), 1.83 (2H, m), 3.86 (2H, t), 6.87-6.99 (4H, m), 7.15-7.20 (4H, m); <sup>13</sup>C-NMR: 14.09, 22.41, 26.66, 29.22, 47.44, 115.42, 122.34, 124.93, 127.19, 127.44, 145.35; IR (CCl<sub>4</sub>): 1592, 1460, 1336, 1288, 1256, 748; <sup>2</sup> **1h**: <sup>1</sup>H-NMR: 0.89 (3H, t), 1.33 (4H, m), 1.45 (2H, m), 1.82 (2H, m), 3.85 (2H, t), 6.87-6.94 (4H, m), 7.15-7.18 (4H, m); <sup>13</sup>C-NMR: 14.10, 22.60, 26.97, 26.99, 28.98, 31.80, 47.47, 115.42, 122.33, 124.94, 127.19, 127.44, 145.37; IR (CCl<sub>4</sub>): 1592, 1460, 1336, 1288, 1248, 748.

Table 2. Yields and melting points for 10-alkylphenothiazine-5-oxides **2a-h**

| No.       | Method<br>Yield [%] |    | M.p./lit. M.p.<br>[°C] | IR  | NMR   |
|-----------|---------------------|----|------------------------|---|---|
|           | A                   | B  |                        |   |   |
| <b>2a</b> | 85                  | 75 | 201-2/194-6 (29)       | 1584, 1480, 1456, 1356, 1264, 1048, 1020, 764       | <sup>1</sup> H-: 3.78 (3H, s), 7.29 (2H, t), 7.40 (2H, d), 7.64 (2H, t), 7.95 (2H, d)   |
| <b>2b</b> | 91                  | 78 | 165-6/162-4 (30)       | 1584, 1488, 1456, 1376, 1256, 1056, 1024, 756       | <sup>1</sup> H-: 1.58 (3H, t), 4.36 (2H, q), 7.25 (2H, t), 7.46 (2H, d), 7.63 (2H, t), 7.94 (2H, d)   |
| <b>2c</b> | 82                  | 78 | 121-2/138-40 (31)      | 1584, 1488, 1460, 1376, 1252, 1052, 1028, 752       | <sup>1</sup> H-: 1.12 (3H, t), 1.97 (2H, m), 4.18 (2H, t), 7.24 (2H, t), 7.41 (2H, t), 7.62 (2H, t), 7.95 (2H, d)   |
| <b>2d</b> | 88                  | 76 | 130-1/-                | 1580, 1480, 1452, 1376, 1248, 1052, 1040, 1032, 768 | <sup>1</sup> H-: 1.08 (3H, t), 1.59 (2H, m), 1.95 (2H, m), 4.22 (2H, t), 7.24 (2H, t), 7.41 (2H, d), 7.62 (2H, t), 7.94 (2H, d)   |
| <b>2e</b> | 90                  | 81 | 181-3/-                | 1584, 1460, 1360, 1248, 1040, 1024, 752             | <sup>1</sup> H-: 0.95 (6H, d), 2.33 (1H, m), 4.09 (2H, d), 7.20 (2H, t), 7.44 (2H, d), 7.55 (2H, t), 7.89 (2H, d); <sup>13</sup> C-: 20.27, 27.07, 54.04, 116.80, 121.96, 125.88, 130.88, 132.44, 139.56  |
| <b>2f</b> | 91                  | 80 | 116-8/-                | 1584, 1460, 1372, 1248, 1176, 1056, 1040, 1024, 768 | <sup>1</sup> H-: 1.08 (d, 6H), 1.87 (3H, m), 4.25 (2H, t), 7.23 (2H, t), 7.41 (2H, d), 7.62 (2H, t), 7.97 (2H, d); <sup>13</sup> C-: 22.50, 26.61, 34.61, 46.70, 115.57, 121.63, 123.99, 131.71, 132.84, 138.27   |
| <b>2g</b> | 87                  | 79 | 113-4/-                | 1580, 1480, 1376, 1248, 1160, 1056, 1040, 1032, 764 | <sup>1</sup> H-: 1.01 (3H, m), 1.47-1.57 (4H, m), 1.97 (2H, m), 4.23 (2H, t), 7.26 (2H, t), 7.43 (2H, d), 7.65 (2H, t), 7.96 (2H, d); <sup>13</sup> C-: 14.07, 24.40, 25.92, 28.97, 48.19, 115.67, 121.67, 123.90, 131.74, 132.88, 138.33   |
| <b>2h</b> | 89                  | 79 | 108-10/-               | 1584, 1456, 1376, 1252, 1160, 1060, 1044, 1032, 768 | <sup>1</sup> H-: 0.93 (3H, t), 1.33-1.38 (4H, m), 1.44 (2H, m), 1.53 (2H, m), 1.96 (2H, m), 4.21 (2H, t), 7.24 (2H, t), 7.40 (2H, d), 7.62 (2H, t), 7.93 (2H, d); <sup>13</sup> C-: 14.19, 22.68, 26.35, 26.94, 31.07, 31.91, 48.34, 115.79, 121.78, 124.04, 131.83, 132.99, 138.45 |

**Table 3.** Yields and melting points for 10-alkylphenothiazine-5,5-dioxides **3a-h**

| No.       | Method Yield [%] |    | M.p./lit. M.p. [°C] | IR   | NMR   |
|-----------|------------------|----|---------------------|--|---|
|           | C                | D  |                     |  |   |
| <b>3a</b> | 89               | 93 | 219-20/220-1 (32)   | 1592, 1480, 1468, 1380, 1280, 1248, 1168, 1156, 1144, 1088, 748, 576 | <sup>1</sup> H-: 3.73 (3H, s), 7.31 (4H, m), 7.64 (2H, t), 8.12 (2H, d)   |
| <b>3b</b> | 93               | 91 | 163-4/162-64 (32)   | 1592, 1480, 1468, 1380, 1280, 1248, 1168, 1156, 1144, 1088, 748, 576 | <sup>1</sup> H-: 1.58 (3H, t), 4.29 (2H, q), 7.29 (2H, t), 7.40 (2H, d), 7.65 (2H, t), 8.15 (2H, d)   |
| <b>3c</b> | 91               | 90 | 193-4/193-95 (32)   | 1592, 1464, 1372, 1288, 1248, 1168, 1152, 1140, 1080, 752, 572       | <sup>1</sup> H-: 1.09 (3H, t), 1.96 (2H, m), 4.13 (2H, t), 7.25 (2H, t), 7.35 (2H, d), 7.63 (2H, t), 8.14 (2H, d)   |
| <b>3d</b> | 89               | 90 | 148-9/149-50 (33)   | 1592, 1480, 1468, 1372, 1288, 1248, 1168, 1152, 1140, 1080, 752, 572 | <sup>1</sup> H-: 1.06 (3H, t), 1.53 (2H, m), 1.92 (2H, m), 4.18 (2H, t), 7.28 (2H, t), 7.36 (2H, d), 7.65 (2H, t), 8.13 (2H, d)   |
| <b>3e</b> | 93               | 92 | 138-9/143-44 (33)   | 1592, 1464, 1336, 1288, 1240, 1164, 1152, 1140, 1104, 1064, 760, 568 | <sup>1</sup> H-: 0.95 (6H, d), 2.26 (1H, m), 4.08 (2H, d), 7.27 (2H, t), 7.39 (2H, d), 7.62 (2H, t), 8.10 (2H, d); <sup>13</sup> C-: 20.38, 27.25, 55.05, 117.36, 122.32, 122.97, 125.97, 133.13, 142.36  |
| <b>3f</b> | 95               | 94 | 140/-               | 1592, 1576, 1464, 1372, 1280, 1248, 1168, 1152, 1140, 1080, 752, 572 | <sup>1</sup> H-: 1.05 (6H, d), 1.83 (3H, m), 4.19 (2H, t), 7.26 (2H, t), 7.37 (2H, d), 7.62 (2H, t), 8.11 (2H, d)   |
| <b>3g</b> | 92               | 91 | 95-6/-              | 1592, 1576, 1464, 1372, 1280, 1248, 1168, 1152, 1140, 1080, 752, 572 | <sup>1</sup> H-: 0.97 (3H, t), 1.45 (4H, m), 1.94 (2H, m), 4.16 (2H, t), 7.27 (2H, t), 7.35 (2H, d), 7.63 (2H, t), 8.13 (2H, d); <sup>13</sup> C-: 14.42, 22.73, 26.84, 29.22, 48.82, 116.35, 122.12, 124.02, 124.62, 133.51, 141.22                    |
| <b>3h</b> | 92               | 92 | 75/-                | 1592, 1464, 1376, 1292, 1248, 1164, 1152, 1144, 1084, 756, 575       | <sup>1</sup> H-: 0.92 (3H, d), 1.34-1.50 (8H, m), 1.94 (2H, m), 4.16 (2H, t), 7.26 (2H, t), 7.35 (2H, d), 7.63 (2H, t), 8.13 (2H, d); <sup>13</sup> C-: 14.05, 22.54, 26.67, 26.76, 28.88, 31.73, 48.46, 115.96, 121.72, 123.62, 124.24, 133.11, 140.83 |

**II. Oxidation of 10-alkylphenothiazines 1a-h to sulfoxides 2a-h**

**Method A:** To a stirred solution of 10-alkylphenothiazine 1a-h (5 mmol) in 25 ml of CH<sub>2</sub>Cl<sub>2</sub>, a solution of *m*-CPBA (5 mmol, ~1.23 g, ~70%, freshly titrated iodometrically) dissolved in 20 ml anhydrous CH<sub>2</sub>Cl<sub>2</sub> was added dropwise (30 min, 0-5 °C). After the reaction was completed (checked by TLC, 2-4 h), the mixture was washed with 10% KOH solution (15 ml, twice), 5% HCl solution (10 ml), and satd. NaHCO<sub>3</sub> solution (10 ml) and dried (MgSO<sub>4</sub>). The solvent was distilled off and residue was separated by column chromatography (CH<sub>2</sub>Cl<sub>2</sub>-acetone, 9:1). Finally, the products were recrystallized from hexane. Yields and melting points for the resulting sulfoxides 2a-h are given in Table 2.

**Method B:** To a cooled solution of 10-alkylphenothiazines 1a-h (5 mmol) in 5 ml CCl<sub>4</sub>, a mixture of 97% fuming nitric acid (5 mmol, 0.22 ml) and glacial acetic acid (0.1 ml) was added dropwise. After a few minutes, a white precipitate was formed which was removed by filtration, washed with saturated NaHCO<sub>3</sub> solution, dried and recrystallized from anhydrous DMF. Characteristics of sulfoxides 2a-h obtained by this method, including the melting points, were identical to those obtained by Method A.

### III. Oxidation of 10-Alkylphenothiazines **1a-h** to 10-alkylphenothiazine-5,5-dioxides **3a-h**

**Method C:** To a solution of 10-alkylphenothiazine **1a-h** (5 mmol) in 25 ml of  $\text{CHCl}_3$ , aqueous solution (50 ml) of  $\text{KMnO}_4$  (7.5 mmol) and  $\text{C}_{16}\text{H}_{35}\text{N}(\text{CH}_3)_3^+(\cdot) \text{Cl}^{(\cdot)}$  (0.5 g) were added and the resulting mixture was vigorously stirred at R.T. for 4–6 h. Then the reaction mixture was filtered on white quartz sand. The organic layer was separated, washed with water (15 ml, four times), dried ( $\text{MgSO}_4$ ) and concentrated in vacuum. The residue was purified by column chromatography (eluent  $\text{CH}_2\text{Cl}_2$ ) and recrystallized from dry ethanol. Yields and melting points for the resulting sulfones **3a-h** are given in Table 3.

**Method D:** To a stirred solution of **1a-h** (5 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 ml) *m*-CPBA (0.01 mmol, ~2.5 g, 70%) was added in small portions at room temperature and the resulting mixture was stirred at R.T. for 2 h. Further work up was carried out as described for Method A, except that ethanol was used for recrystallization. Characteristics of sulfones **3a-h** obtained by this method were identical to those obtained by using Method C.

### Results and Discussion

10-Alkylphenothiazines **1a-h** were obtained from phenothiazine by reaction with the corresponding alkyl iodides in the presence of  $\text{NaNH}_2$  without using solvent (Scheme 1, Table 1).

10-Alkylphenothiazine-5-oxides **2a-h** were prepared from the appropriate 10-alkylphenothiazines **1a-h** by two alternative methods (Scheme 1, Table 2). Sulfoxide formation was carried out by dropwise addition of *m*-CPBA acid dissolved in anhydrous  $\text{CH}_2\text{Cl}_2$  into a solution of the *N*-alkylphenothiazines **1a-h** in  $\text{CH}_2\text{Cl}_2$  at 0 °C (Method A). The same 10-alkylphenothiazine-5-oxides **2a-h** were obtained by addition of glacial  $\text{AcOH}$  solution of fuming  $\text{HNO}_3$  into a solution of *N*-alkylphenothiazines **1a-h** in  $\text{CCl}_4$  (Method B), also at 0 °C. Similarly, two alternative methods were elaborated for oxidation of 10-alkylphenothiazines **1a-h** into the corresponding 10-alkylphenothiazine-5,5-dioxides **3a-h** (Scheme 1, Table 3). The 10-alkylphenothiazines **1a-h** were transformed into their sulfones **3a-h** at R.T. by  $\text{KMnO}_4$  in a water-chloroform two-phase system using cetyltrimethylammonium chloride phase-transfer catalyst (Method C). Alternatively, portionwise addition of solid *m*-CPBA acid into  $\text{CH}_2\text{Cl}_2$  solution of 10-alkylphenothiazines **1a-h** at R.T. resulted in smooth formation of the 10-alkylphenothiazine-5,5-sulfones **3a-h** (Method D). Selective formation of 10-alkylphenothiazine-5-oxides **2a-h** (Method A) was achieved at low temperature (0 °C) in highly diluted system by using exact equivalency of precisely titrated *m*-CPBA. Twofold amount of *m*-CPBA at R.T., however, resulted in the formation of 10-alkylphenothiazine-5,5-dioxides **3a-h** (Method D). Since sulfoxides **2a-h** are poorly soluble in  $\text{CCl}_4$ , they precipitated in the case of Method B and no formation of nitro-compounds were detected. These results confirm that *S*-oxidation of *N*-alkylphenothiazines by nitric acid precedes nitration in formation of nitrophenothiazine sulfoxides (34). Efficiency of the  $\text{KMnO}_4$  oxidation was enhanced by using phase-transfer catalysis (Method C). By this modification, oxidation of the *N*-alkylphenothiazines **1a-h** to the corresponding sulfones **3a-h** could be performed at R.T. within 2 h in good yields.

### Acknowledgements

Cs. P. and C. M. thank Domus Hungarica Foundation for short term fellowships.

### References

- (1) R.R. Gupta, Phenothiazines and 1,4-Benzothiazines. Chemical and biomedical aspects, Elsevier, Amsterdam, 1988.
- (2) C. Bodea and I. Silberg, *Adv. Het. Chem.* **9**, 321 (1968).
- (3) M. Aravagiri, S. R. Marder, A. Yuwiler, K. K. Midha, N. S. Kula, and R. J. Baldessarini, *Neuropsychopharmacology* **13**, 235 (1995).
- (4) M. Chetty, V. L. Pillay, S. V. Moodley and R. Miller, *Eur. Neuropsychopharmacol.* **6**, 85 (1996).
- (5) H. Gilman and D. Nelson, *J. Am. Chem. Soc.* **75**, 5422 (1953).
- (6) H. Gilman and R. Rank, *J. Org. Chem.* **23**, 1014 (1958).
- (7) H. Wunderlich and A. Stark, *Pharmazie* **20**, 519 (1965).
- (8) Y. Girard, P. Hamel, M. Therien, J. P. Springer and J. Hirshfield, *J. Org. Chem.* **52**, 4000 (1987).
- (9) A. McKillop and D. Kemp, *Tetrahedron* **45**, 3299 (1989).
- (10) G. Cauquil and A. Casadevall, *Compt. Rend. Acad. Sci.* **240**, 1997 (1955).
- (11) G. M. Nano and P. Sancic, *Pharm. Acta Helv.* **38**, 623 (1963).
- (12) C. Kaiser, D.H. Tedeschi, P.J. Fowler, A.M. Pavloff, B.M. Lester and C.L. Zirkle, *J. Med. Chem.* **14**, 176 (1971).
- (13) M. V. Jovanovic and E. R. Biehl, *J. Heterocycl. Chem.* **21**, 1589 (1984).
- (14) M. L. Owens, E. C. Juenge and A. Poklis, *J. Pharm. Sci.* **78**, 334 (1989).
- (15) E. Bosch and J. K. Kochi, *J. Chem. Soc. Perkin Trans. 1* **8**, 1057 (1995).
- (16) E. Bosch and J. K. Kochi, *J. Org. Chem.* **60**, 3171 (1995).
- (17) D. A. Simov, *Izv. Khim. Inst. Bulgar. Acad. Nauk* **5**, 51 (1957); *Chem. Abstr.* **55**, 16556d (1961).
- (18) D. Scholz *Monatsh. Chem.* **241** (1981).
- (19) W. Priebe and G. Grynkiewicz, *Tetrahedron. Lett.* **32**, 1749 (1991).
- (20) E. A. Nodiff, *J. Heterocycl. Chem.* **8**, 321 (1971).
- (21) M. Nakapishi and N. Kuriyama, *Jap. Patent* 15617 (1961); *Chem. Abstr.* **56**, 12908i (1962).
- (22) K. Yasukouchi, I. Taniguchi, H. Yamaguchi, J. Ayukawa, K. Ohtsuka, Y. Tsuruta, *J. Org. Chem.* **46**, 1679 (1981).
- (23) W. Adam, S. Hueckmann and F. Vargas, *Tetrahedron Lett.* **30**, 6315 (1989).
- (24) D.D. DesMarteau, V.A. Petrov, V. Montarini, M. Pregnotato, G. Resnati *J. Org. Chem.* **59**, 2769 (1994).
- (25) H. Fujita and Y. Yamuchi, *J. Het. Comm.* **17**, 1053 (1980).
- (26) C. Lespagnol, *Bull. Soc. Chim. Fr.* 112 (1960).
- (27) R. Dahlbam, *Acta Chem. Scand.* **3**, 247 (1949).
- (28) D. Clarke, B. Gilbert and P. Hanson, *J. Chem. Soc. Perkin Trans. 2* 1103 (1978).
- (29) H. J. Shine, *J. Het. Chem.* **4**, 517 (1969).
- (30) E. H. Biehl and V. Patrizi, *J. Het. Chem.* **11**, 247 (1974).
- (31) E. Gilman, *J. Org. Chem.* **19**, 560 (1954).
- (32) E. Gilman, *J. Am. Chem. Soc.* **75**, 5422 (1953).
- (33) H. Wunderlich, *Pharmazie* **21**, 57 (1966).
- (34) C. Schmalz and A. Burger, *J. Am. Chem. Soc.* **76**, 5455 (1954).

Received on April 2, 2001