


This article was downloaded by: [University of Miami]

On: 29 December 2014, At: 06:42

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

AN EFFICIENT SYNTHESIS OF 4,4'-DIAMINODIPHENIC ACID

Arkadiy Rubezhov^a

^a Laboratory for SR&D, Wyeth, 401 N Middletown Rd, Pearl River, NY, 10965-1299, USA

Published online: 11 Feb 2009.

To cite this article: Arkadiy Rubezhov (2006) AN EFFICIENT SYNTHESIS OF 4,4'-DIAMINODIPHENIC ACID, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 38:4, 401-404

To link to this article: <http://dx.doi.org/10.1080/00304940609356000>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

OPPI BRIEFS

AN EFFICIENT SYNTHESIS OF 4,4'-DIAMINODIPHENIC ACID

Submitted
(11/07/05)


Arkadiy Rubezhov

Laboratory for SR&D, Wyeth
401 N Middletown Rd, Pearl River, NY 10965-1299, USA
e-mail: rubezha@wyeth.com

4,4'-Diaminodiphenic acid (DDPA, **4**) has received attention in recent years due to its importance for the synthesis of biologically active compounds¹ and in material science.² In particular, it is a key intermediate in the synthesis of polyamides,³ polyesters,⁴ photo-resists,⁵ liquid crystals,⁶ pigments and printing inks,⁷ and also as a reagent in analytical chemistry.⁸ Of the several reported DDPA preparations,^{9,10} the direct nitration of diphenic acid **1**, followed by catalytic hydrogenation, appears acceptable.¹⁰ However, the chromatography step and low overall yield of DDPA (25-27%) make this approach unsuitable for large scale syntheses. This article describes a novel, practical, and economical synthesis of **4** by a two-step procedure.

It was believed that, in the hydrogenation process, the amino group of 4,6'-diamino isomer (a minor component in the mixture) would undergo cyclization with the neighboring 2-carboxy group to give the intermolecular lactam **5** which should be easily separable from the desired 4,4'-diamino derivative **4** by careful control of the pH.

As a matter of fact, hydrogenation and chemical separation of the products were performed in a telescopic process. All steps may be carried out without isolation and purification of intermediates. After nitration, the mixture was poured into ice and allowed to warm to ambient temperature. The resulting suspension was neutralized with 10N aqueous NaOH to give



a solution of sodium salts which was hydrogenated. After hydrogenation, the basic reaction mixture was filtered through Magnesol®, and the mother liquor was titrated with glacial AcOH. At pH 5.29-6.06, the zwitterion **6** of the 4,4'-diaminodiphenic acid (**4**) precipitated as a grayish-white crystals while the zwitterion **7** of lactam **5** settled from the mother liquor as white solid at pH 3-4 (titration with 1N aqueous HCl). Sixty-two grams of 4,4'-diaminodiphenic acid zwitterion (**6**) and 18 g of the lactam zwitterion (**7**) were isolated starting from 100 g of diphenic acid **1**. The overall conversion to **6** and **7** is 94%. The structure of the 4,4'-diaminodiphenic zwitterion **6** was established by conversion to the corresponding dimethyl ester and comparison with an authentic sample.¹¹

EXPERIMENTAL SECTION

2,2'-Diphenic acid (98% purity) and 10 wt. % Pd/C, Degussa type E101 (water ~50%) were purchased from Aldrich Chemical Co. 30/40 Magnesol® is available from BNL. Hydrogenation was carried out in 2 L Parr reactor; TLC was performed on Kieselgel 60 F₂₅₄ (Merck) plate using EtOAc-heptane (1:1 v/v) as eluent. Melting points were determined on a Thomas Electro thermometer. Elemental analyses were carried out by the Analytical Laboratory, Wyeth, Pearl River, NY.

4,4'-Dinitro-2,2'-diphenic Acid (2) and 4, 6'-Dinitro-2, 2'-diphenic Acid (3).- To a mixture of 90% HNO₃ (750 mL) and conc. H₂SO₄ (250 mL) were added at 0°C the small portions of 2,2'-diphenic acid (**1**) (100.0 g, 0.412 mol) over about 2 h. At this point TLC showed no starting material but only the 4, 4'-dinitro isomer (R_f 0.63) and the 4, 6'-dinitro isomer (R_f 0.36). The reaction mixture was poured onto crushed ice (1.5 kg). The white precipitate was collected, washed with water, acetone and dried to give 125.7 g (98%) of mixture **2** and **3** which was used for hydrogenation without further purification.

4,4'-Diamino-2,2'-diphenic Acid (4) and 3-Amino-5,6-dihydrobenzo[c]quinolin-10-carboxylic Acid (5).- A solution of **2** and **3** (125.7 g) in 1 L of water (pH 2.78) was basified with 10 N NaOH to pH 8.5. This solution was hydrogenated at 50 psi of H₂ for 1.5 h in the presence of 10% Pd/C. When the reduction was completed (TLC control), the mixture was filtered through Magnesol. The filtrate (pH 9.36) was used for the isolation of **6** and **7** as described below.

4,4'-Diamino-2,2'-diphenic Acid (6, zwitterion) and Lactam (7, zwitterion).- The filtrate from the previous experiment was acidified very slowly by titration with AcOH. At pH 5.29, light grey crystals began to appear. The mixture was stirred for 1 h, after which time the pH rose to 6.06. More AcOH was added to adjust pH to 5.3, and again the mixture was stirred for 1 h. The crystals were collected, washed with water and acetone and then dried to give 62.0 g (55%) of light grey solid of 99.3% purity (HPLC area%), mp. > 230°C (dec.), *lit.*¹¹ mp. 265°C.

Anal. Calcd for C₁₄H₁₂N₂O₄: C, 61.76, H, 4.41, N, 10.29. Found: C, 61.97, H, 4.43, N, 10.13.

The clear filtrate from above was acidified to pH ~3 with 1N HCl. The resulting white solid (**7**) was collected, washed with water, acetone and dried to give 18.0 g (17%) of colorless solid, mp. > 300°C (dec.), *lit.*¹⁰ mp. 330-335°C.

Anal. Calcd for $C_{14}H_{10}N_2O_3$: C, 66.14, H, 3.94, N, 11.02. Found: C 65.94, H, 3.87, N, 10.86.

Dimethyl 4,4'-Diamino-2,2'-diphenolate.- A mixture of **6** (5.0 g, 0.018 mol), MeOH (25 mL) and conc. H_2SO_4 was refluxed for 48 h with TLC monitoring of the disappearance of starting material, appearance and disappearance of monoester and accumulation of the desired diester. When reaction was complete, the reaction mixture was poured into water, basified to pH ~10 with 20% aqueous NH_4OH and extracted with CH_2Cl_2 . The organic phase was dried over $CaSO_4$ and evaporated to give 4.5 g (82%) of dimethyl 4,4'-diamino-2,2'-diphenolate as a colorless solid, mp. 103-106°C, undepressed upon admixture with an authentic sample.¹²

Anal. Calcd for $C_{16}H_{16}N_2O_4$: C, 64.00, H, 5.33, N, 9.33. Found: C, 64.34, H, 5.42, N, 9.05.

Acknowledgments.- Drs J. Potoski and A. Mohan and all the RSV team are acknowledged for mutually beneficial discussions and Dr W. McMahon is acknowledged for friendly support.

REFERENCES

1. W.-D. Ding, B. Mitsner, G. Krishnamurthy, A. Aulabaugh, C. D. Hess, J. Zaccardi, M. Cutler, B. Feld, A. Gazumyan, Y. Raifeld, A. Nikitenko, S. A. Lang, Y. Gluzman, B. O'Hara and G. A. Ellestad, *J. Med. Chem.*, **41**, 2671 (1998).
2. D. L. Singleton, G. Paraskevopoulos, A. D. Buckthought, R. S. Irwin and I. T. Emesh, *Proceeding of SPIE-The International Society for Optical Engineering*, 48 (1990); *Chem. Abstr.*, **113**, 450956 (1990).
3. S. Irwin, *US Patent 5039785* (1991); *Chem. Abstr.*, **115**, 159430 (1991).
4. H. E. Hinderer, R. W. Smith and J. Preston, *J. Applied Polymer Symposia*, 1 (1973); *Chem. Abstr.*, **82**, 82480 (1975).
5. K. Yasuda, Y. Yokoyama and N. Besso, *W. O. Patent 2004048462* (2004); *Chem. Abstr.*, **141**, 24506 (2004).
6. W. Kreuder, H. Ringsdorf, O. Herrmann-Schoenherr and J. H. Wendorff, *Angew. Chem.*, **99**, 1300 (1987).
7. J. A Stirling, *US Patent 5306342* (1994); *Chem. Abstr.*, **121**, 111473 (1994).
8. V. I Agafonova and I. P. Ryazanov, *Izvestiya Vysshikh Uchebnykh Zavedenii, Khimiya i Khimicheskaya Tekhnologiya*, **12**, 1326 (1969); *Chem. Abstr.*, **72**, 96256 (1970).
9. H. R. Patel, D. W. Blackburn and G. I. Jenkin, *J. Amer. Pharm. Assoc.*, **46**, 51 (1957); *Chem. Abstr.*, **82**, 87480 (1975).
10. G. I. Migachev, A. M. Andrievskii and N. S. Dokunikhin, *Zh. Org. Khimii*, **13**, 463 (1977); *Chem. Abstr.*, **85**, 23105 (1977).
11. H. Adkins, E. F. Steinbring and E. Pickering, *J. Am. Chem. Soc.*, **46**, 1917 (1924).

12. A sample of dimethyl 4,4'-diaminodiphenolate for comparison was kindly supplied by S. Strunk of Wyeth.

A CONVENIENT, ONE-POT SYNTHESIS OF THIOCARBAMATES USING *bis*(TRICHLOROMETHYL) CARBONATE

Submitted by
(11/11/05)

W. K. Su*, J. P. Zhang and X. R. Liang

*Zhejiang Key Laboratory of Pharmaceutical Engineering
College of Pharmaceutical Sciences
Zhejiang University of Technology, Hangzhou, 310014, P. R. CHINA
E-mail: suweike@zjut.edu.cn*

Thiocarbamates are an important class of compounds due to their biological activity¹ and have found wide applications in the chemical industry such as in the production of commodity chemicals such as herbicides,² pesticides,³ bactericides⁴ and antiviral agents.⁵ Therefore the development of methods for the synthesis of thiocarbamates is important.

Several methods for the preparation of thiocarbamates have been reported in the literature⁶ such as the reaction of amines with phosgene and thiols.⁷ In addition, the direct condensation of thiols with carbamoyl chlorides⁸ or isocyanates⁹ has also been reported. Unfortunately, both carbamoyl chlorides and isocyanates are typically prepared from phosgene and some of them are difficult to store, as they are sensitive to water. There are a variety of other methods including (i) the reaction of carbon monoxide, amines with thiols¹⁰ or disulfides¹¹ in the presence of metal catalysts, (ii) the use of sulfur¹² with amines and carbon monoxide followed by addition of alkyl halides, (iii) coupling of thiocyanates and alcohols in the presence of sulfuric acid,¹³⁻¹⁴ (iv) the conversion of various *o*-substituted thiocarbamates *via* intramolecular rearrangement.¹⁵ Recently, Wynne reported a synthesis of thiocarbamates from thiols¹⁶ in which two equivalents of trichloroacetyl chloride and of amines were required. Despite of the numerous routes, most methods involve unstable reagents, multi-step procedures, harsh reaction conditions, expensive catalyst or complex materials. Base on these premises, we now report a mild and convenient one-pot procedure for the synthesis of thiocarbamates using *bis*(trichloromethyl) carbonate (BTC).

BTC is well known as an important equivalent for phosgene and has emerged as a versatile synthetic auxiliary in the preparation of various organic compounds.¹⁷ Reactions with BTC usually proceed under relatively mild conditions and often afford good to excellent yields. The present work reports the reaction of BTC for the synthesis of thiocarbamates from thiophenols and amines.