

Synthesis and Characterization of the Novel Copper(I) Anion $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$ †

Graham A. Bowmaker*

Department of Chemistry, University of Auckland, Private Bag, Auckland, New Zealand

Brian W. Skelton and Allan H. White

Department of Physical and Inorganic Chemistry, University of Western Australia, Nedlands, Western Australia 6009, Australia

Peter C. Healy

Department of Science and Technology, Griffith University, Nathan, Queensland 4111, Australia

Solid-state ^{31}P n.m.r., far-i.r. spectroscopy and single-crystal X-ray diffraction techniques have been used to characterize a new type of ionic copper(I) compound: $[\text{PPh}_3\text{Me}][\text{Cu}(\text{PPh}_3)\text{Br}_2]$. X-Ray analysis shows the PPh_3 ligand to be co-ordinated to the copper atom of the CuBr_2 unit in a trigonal planar arrangement, with Cu–P 2.210(3), Cu–Br 2.367(2), 2.386(2) Å and Br–Cu–Br 114.9(1)° {monoclinic, space group Pn , $a = 11.779(5)$, $b = 9.914(5)$, $c = 14.705(10)$ Å, $\beta = 94.92(5)^\circ$, $Z = 2$, $R = 0.037$ for 1 570 ‘observed’ [$I > 3\sigma(I)$] reflections}. The far-i.r. spectrum shows a strong band at 195 cm^{-1} assigned to the antisymmetric Cu–Br stretching mode of the CuBr_2 unit, and a weaker band at 150 cm^{-1} assigned to the symmetric Cu–Br stretching mode. This assignment is supported by an approximate normal-co-ordinate analysis, and the resulting Cu–Br force constant is compared with values obtained in similar analyses of related bromocopper(I) complexes. The cross-polarization magic angle spinning ^{31}P n.m.r. spectrum shows a strong signal at 21 p.p.m. (relative to 85% H_3PO_4) due to the cation, and a weaker quartet centred at –14 p.p.m. due to the co-ordinated PPh_3 in the anion, with line spacings due to Cu–P coupling of 1.37, 1.69, and 1.92 kHz.

Copper(I) generally shows a preference for a co-ordination number of four in its complexes, although an increasing number of complexes have been reported in which it has co-ordination numbers of two or three.¹ The factors which determine the preferred co-ordination number are not entirely clear, however, and further data relevant to this question should be obtainable by studying related series of complexes to determine the range of possible co-ordination numbers in the series, and by measuring physical properties of the complexes which yield information about the changes in bond strength and valence-electron distribution which occur with change in co-ordination number. Thus, for the series $\text{CuX}_n^{(n-1)-}$ with $\text{X} = \text{Br}$ or I , complexes up to $n = 3$ are known, whereas for $\text{X} = \text{Cl}$ only up to $n = 2$. The metal–halogen force constants obtained from the $\nu(\text{Cu–X})$ vibrational frequencies for the $\text{X} = \text{Br}$ or I complexes decrease rapidly with n , and these results suggest that the $n = 4$ complexes should not exist for this series.² In the case of the series $\text{Cu}(\text{PPh}_3)_n\text{X}$, mononuclear complexes with $n = 2$ or 3 are known for $\text{X} = \text{Cl}$, Br , or I .^{3,4} Thus, the replacement of a halide ligand X^- by PPh_3 appears to favour higher co-ordination numbers in copper(I) complexes.

In a development of these studies, we are examining the ability of neutral ligands to co-ordinate to the two-co-ordinate anions CuX_2^- ($\text{X} = \text{Cl}$, Br , or I). In the present communication we report the preparation and characterization by X-ray diffraction, i.r., and solid-state cross-polarization magic angle spinning (c.p.m.a.s.) n.m.r. spectroscopy of the novel anion $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$ in the compound $[\text{PPh}_3\text{Me}][\text{Cu}(\text{PPh}_3)\text{Br}_2]$.

Experimental

Preparation of Compounds.—Methyltriphenylphosphonium dibromotriphenylphosphinecuprate(I), $[\text{PPh}_3\text{Me}][\text{Cu}(\text{PPh}_3)\text{Br}_2]$ —

$\text{Br}_2]$. Copper(I) bromide (0.36 g, 2.5 mmol), triphenylphosphine (0.66 g, 2.5 mmol), and methyltriphenylphosphonium bromide (0.89 g, 2.5 mmol) were added to dichloromethane (15 cm^3) and the mixture was stirred until all the solid material had dissolved. The solution was heated on a water-bath until it boiled, and then diethyl ether was slowly added until the first permanent precipitate formed. On cooling to room temperature the solution deposited white crystals which were collected by filtration and dried in a vacuum, m.p. 167–168.5 °C (Found: C, 58.1; H, 4.9. Calc. for $\text{C}_{37}\text{H}_{33}\text{Br}_2\text{CuP}_2$: C, 58.3; H, 4.4%).

The unsolvated adducts of $[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$ and the ‘step’ isomer of $[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ were prepared as previously described.⁵

Structure Determination.—Crystal data. $\text{C}_{37}\text{H}_{33}\text{Br}_2\text{CuP}_2$, $M = 763.0$, monoclinic, space group Pn [C_2^2 , no. 7 (variant)], $a = 11.779(5)$, $b = 9.914(5)$, $c = 14.705(10)$ Å, $\beta = 94.92(5)^\circ$, $U = 1.711(2)$ Å³, $D_c(Z = 2) = 1.48$ g cm^{-3} , $F(000) = 768$, $\mu(\text{Mo–K}_\alpha) = 30.3$ cm^{-1} , specimen $0.51 \times 0.40 \times 0.04$ mm (capillary), $A^*_{\text{min.,max.}} = 1.13, 2.50$ (Gaussian correction).

A unique data set was measured to $2\theta_{\text{max.}} = 50^\circ$ on a Syntex $P2_1$ four-circle diffractometer (monochromatic Mo- K_α radiation source, $\lambda = 0.71069$ Å) in conventional 2θ – θ scan mode at 295 K. 2 243 Independent reflections were measured, 1 570 with $I > 3\sigma(I)$ being considered ‘observed’ and used in the full-matrix least-squares refinement after solution of the structure by vector methods. ($x, y, z, U_{\text{iso}})_\text{H}$ were included at estimated values. Residuals on $|F|$ at convergence were $R = 0.037$, $R' = 0.024$ (preferred hand); statistical reflection weights were used, derived from $\sigma^2(I) = \sigma^2(I)_{\text{diff}} + 0.0001\sigma^4(I)_{\text{diff}}$. The structure refined smoothly in the chosen space group with no anomalies with respect to features such as thermal envelopes or disorder. The residuals were lowest for the chosen enantiomer. Neutral atom complex scattering factors were used.⁶ No significant extinction effects were observed. Computation used the XTAL83 program system⁷ implemented by S. R. Hall on a Perkin-Elmer 3240

† Supplementary data available: see Instructions for Authors, *J. Chem. Soc., Dalton Trans.*, 1988, Issue 1, pp. xvii–xx.

Table 1. Non-hydrogen atom co-ordinates for $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$

Atom	$l = 1$			$l = 2$		
	x	y	z	x	y	z
Cu(1)*	0	0.264 3(1)	0	—	—	—
Br(1)	0.173 7(1)	0.144 7(1)	0.036 5(1)	—	—	—
Br(2)	−0.158 8(1)	0.180 1(1)	0.070 5(1)	—	—	—
P(1)	−0.022 1(2)	0.449 1(3)	−0.083 8(3)	0.454 5(2)	0.783 4(3)	0.017 0(2)
C(11)	0.096 6(8)	0.500 0(10)	−0.148 6(6)	0.490 8(7)	0.859 4(10)	0.126 7(6)
C(12)	0.205 1(8)	0.450 9(10)	−0.119 1(7)	0.443 6(9)	0.982 7(11)	0.146 5(7)
C(13)	0.296 5(8)	0.487 6(11)	−0.164 4(6)	0.473 9(10)	1.045 2(12)	0.228 7(8)
C(14)	0.286 6(8)	0.574 0(11)	−0.235 8(7)	0.552 7(10)	0.983 2(13)	0.290 6(7)
C(15)	0.180 6(9)	0.625 1(12)	−0.266 6(7)	0.600 3(9)	0.859 1(11)	0.271 9(7)
C(16)	0.086 8(9)	0.587 0(11)	−0.220 6(7)	0.567 2(8)	0.798 6(11)	0.189 1(7)
C(21)	−0.048 4(8)	0.594 4(9)	−0.012 6(7)	0.486 9(8)	0.605 4(9)	0.021 6(6)
C(22)	−0.120 7(8)	0.578 9(10)	0.055 3(7)	0.416 3(9)	0.517 2(11)	0.062 0(7)
C(23)	−0.142 7(8)	0.684 6(10)	0.112 2(7)	0.435 2(8)	0.380 3(11)	0.060 7(7)
C(24)	−0.092 8(9)	0.806 8(10)	0.102 5(7)	0.525 5(9)	0.331 8(10)	0.015 6(6)
C(25)	−0.021 0(9)	0.827 2(10)	0.037 6(8)	0.596 2(8)	0.419 9(11)	−0.023 9(7)
C(26)	0.002 4(9)	0.723 1(10)	−0.021 7(7)	0.577 6(8)	0.554 5(11)	−0.024 8(7)
C(31)	−0.148 2(7)	0.448 0(10)	−0.167 8(6)	0.532 2(8)	0.861 6(11)	−0.070 1(6)
C(32)	−0.187 1(8)	0.328 6(10)	−0.201 3(6)	0.489 0(9)	0.854 4(11)	−0.159 9(7)
C(33)	−0.279 4(8)	0.321 9(12)	−0.263 0(7)	0.554 9(10)	0.903 4(13)	−0.225 4(7)
C(34)	−0.335 5(8)	0.439 3(13)	−0.293 6(7)	0.659 9(11)	0.953 4(12)	−0.201 5(8)
C(35)	−0.297 7(8)	0.561 3(11)	−0.260 9(7)	0.703 0(9)	0.966 5(13)	−0.113 0(8)
C(36)	−0.203 3(8)	0.567 3(10)	−0.197 9(9)	0.637 0(10)	0.918 8(12)	−0.046 2(7)
C	—	—	—	0.304 8(8)	0.801 1(11)	−0.014 4(7)

* Defines origin.

computer and the CHEM-X program package (Chemical Designs Ltd.) implemented on a VAX-780 computer. Non-hydrogen atom co-ordinates are listed in Table 1; P(1) refers to the phosphorus atom in the anion and P(2) to the phosphorus atom in the cation. The un-numbered C is the methyl carbon in the $[\text{PPh}_3\text{Me}]^+$ cation. The atom labelling scheme adopted for each PPh_3 moiety is such that the *ortho* carbon atoms C(*m*2), where *m* is the ring number, are defined as those atoms above the PC_3 pyramid of the ligand.

Additional material available from the Cambridge Crystallographic Data Centre comprises H-atom co-ordinates, thermal parameters, and remaining bond distances and angles.

Spectroscopy.—Solid-state c.p.m.a.s. ^{31}P n.m.r. spectra were obtained at room temperature on a Bruker CXP-300 spectrometer at 121.47 MHz using ^1H – ^{31}P cross polarization with radiofrequency fields of 8 and 20 G ($\text{G} = 10^{-4}$ T) respectively.

Far-i.r. spectra were recorded at *ca.* 298 K as petroleum jelly mulls between Polythene plates on a Digilab FTS-60 Fourier-transform i.r. spectrometer employing a FTS-60V vacuum optical bench and a 6.25- μm mylar beam splitter.

Results and Discussion

Copper(I) bromide, bromide ion, and triphenylphosphine combine in a 1:1:1 ratio to form the $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$ ion which was crystallized from dichloromethane–diethyl ether as its $[\text{PPh}_3\text{Me}]^+$ salt. Attempts to prepare the 2:1 complex of PPh_3 with CuBr_2^- by using an excess of PPh_3 resulted only in the formation of the 1:1 complex.

X-Ray Structure Determination.—The structure of the $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$ anion is shown in Figure 1. The PPh_3 ligand is co-ordinated to the copper atom of the CuBr_2 unit in a three-co-ordinate trigonal-planar arrangement, with Cu–P 2.210(3), Cu–Br 2.367(2), 2.386(2) Å and Br–Cu–Br 114.9(1), P–Cu–Br






Figure 1. The anion, projected normal to the PCuBr_2 plane; 20% thermal ellipsoids and atom numbering are given for the non-hydrogen atoms. Hydrogen atoms have an arbitrary radius of 0.1 Å

118.5(1), 126.5(1)°. The Cu–P distance is somewhat surprisingly similar to the distances of 2.191(2) Å found for $[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$ and 2.206(5) Å found in the 'step' isomer of $[\text{Cu}(\text{PPh}_3)\text{Br}_4]^{5-}$, both of which contain the $(\text{PPh}_3)\text{CuBr}_2$ unit (Table 2). The Cu–Br distances increase on co-ordination of the PPh_3 molecule to CuBr_2^- from an average value of 2.22 Å found in compounds containing uncomplexed linear CuBr_2^- ions⁸ to a mean value of 2.377 Å in $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$, with the Br–Cu–Br angle decreasing from linearity to a value less than 120°. In the complex $[\text{tst}]^+[\text{CuBr}_2]^-$ (tst = 'tetraselenotetracene', naph-


Table 2. Core geometries for molecules containing the $(\text{PPh}_3)\text{CuBr}_2$ fragment

Parameter ^a	$[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$	$[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]^b$	$[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ 'step' ^b
Cu-Br(1)	2.386(2)	2.409(2)	2.427(2)
Cu-Br(2)	2.367(2)	2.364(2)	2.371(3)
Cu-P	2.210(3)	2.191(2)	2.206(5)
P-Cu-Br(1)	126.5(1)	129.7(1)	119.2(1)
P-Cu-Br(2)	118.5(1)	117.8(1)	127.7(1)
Br(1)-Cu-Br(2)	114.9(1)	111.5(1)	112.9(1)

^a Distances (Å), angles (°). ^b Ref. 5.**Figure 2.** Plot of Br-Cu-Br angle vs. Cu-Br distance in compounds containing a CuBr_2 unit**Scheme.**

thaceno[5,6-*cd*:11,12-*c'd'*]bis[1,2]diselenole),⁹ a weak $\text{Cu} \cdots \text{Br}$ bridging interaction between the CuBr_2 units results in a decrease of the Br-Cu-Br angle to $153.8(1)^\circ$, and intermediate Cu-Br distances of $2.267(2)$ and $2.282(3)$ Å (Figure 2). The Cu-Br distances and Br-Cu-Br angles in the trigonal $(\text{PPh}_3)\text{CuBr}_2$ fragment of the aggregated multinuclear complex $[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$ and the 'step' isomer of $[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ are also quite similar to those in the present compound (Table 2). Consideration of the data for the bridged complexes alone might have led to the conclusion that the reduction of the Br-Cu-Br angle to less than 120° is dictated by the bonding requirements of the bridging bromine atoms. It is significant to note, therefore, that this also occurs in $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$, where no such bridging bonds occur.

Structurally, the present compound is a member of the series $[\text{Cu}(\text{PPh}_3)_n\text{Br}_{3-n}]^{(2-n)-}$ (I)–(IV). In a range of structures containing the $[\text{Cu}(\text{PPh}_3)_3]^+$ cation (I),¹⁰ the counter anion is co-ordinated to the copper atom and the P_3Cu core deviates significantly from planarity. For complexes with weakly co-ordinating anions such as ClO_4^- or BF_4^- the average P-Cu-P angle is 116° and Cu-P 2.31 Å. For the neutral planar compound $[\text{Cu}(\text{PPh}_3)_2\text{Br}]$, (II), the Cu-P distances decrease to an average value of 2.27 Å and Cu-Br $2.346(2)$ Å.³ In the present compound (III), as already discussed, Cu-P $2.210(3)$ Å and the average Cu-Br distance is 2.37 Å. Structural data are, as yet, not available for the $[\text{CuBr}_3]^{2-}$ anion, (IV).² These results are

**Figure 3.** Far-i.r. spectrum of $[\text{PPh}_3\text{Me}][\text{Cu}(\text{PPh}_3)\text{Br}_2]$

consistent with a greater σ -donor capacity for the PPh_3 ligand relative to the Br^- anion, leading to strong Cu-P bonds in which the Cu-P distance is determined primarily by the packing requirements of one, two, or three PPh_3 ligands around the relatively small Cu^+ ion. It is possible to argue from these results that the Cu-Br bond is relatively ionic, and in this context the similarity in the PCuBr_2 core geometries of $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$, $[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$, and the 'step' isomer of $[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ is an important result.

Far-i.r. Spectrum.—The far i.r. spectrum of $[\text{PPh}_3\text{Me}][\text{Cu}(\text{PPh}_3)\text{Br}_2]$ is shown in Figure 3. By comparison with the spectrum of $[\text{PPh}_3\text{Me}]\text{Br}$, the band at 376 cm^{-1} can be assigned to the cation and the bands at 195 and 150 cm^{-1} to the $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$ complex. By analogy with the results for other triphenylphosphine bromocopper(I) complexes,¹¹ the strongest bands in the far-i.r. are expected to be the Cu-Br stretching modes, $\nu(\text{CuBr})$. Thus the two bands at 195 and 150 cm^{-1} are assigned to the two expected $\nu(\text{CuBr})$ modes of the CuBr_2 unit. We have found previously that the metal-halogen vibrational frequencies of the metal-halogen core in phosphine metal halide complexes can be calculated by carrying out a normal-coordinate analysis on the metal-halogen core only.¹² We have used this approach here to calculate the frequencies of a CuBr_2 unit with two equivalent bonds and a Br-Cu-Br bond angle of 115° . A simple diagonal force field was used. This involves a Cu-Br bond stretching force constant f_r and a Br-Cu-Br angle-bending force constant f_a . With the assumption $f_a = 0.1f_r$ which we have used in this type of calculation previously,¹² f_r was varied and the vibrational frequencies calculated to give the best fit between the observed and experimental frequencies. The results are shown in Table 3. The agreement between the observed and

Table 3. Observed and calculated $\nu(\text{CuBr})$ wavenumbers, and Cu–Br bond-stretching force constants for some bromocopper(I) complexes


Complex	ν/cm^{-1}		$f_t/\text{N m}^{-1}$	Ref.
	obs.	calc.		
$[\text{CuBr}_2]^-$	321	321(Σ_u^+)	157	13
	191	191(Σ_g^+)		
$[\text{Cu}(\text{PPh}_3)_2\text{Br}_2]^-$	195	189(B_1)	60	*
	150	156(A_1)		
$[\text{Cu}(\text{PPh}_3)_2\text{Br}]$	221	221(A_1)	102	11,*
$[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$	180	180(B_1),	55	14,*
		153(A_1)	(trigonal Cu)	
	128	129(B_1),	32	
		127(A_1)	(tetrahedral Cu)	
$[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ 'step'	174			12
	135			
$[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ 'cubane'	144	145(T_2)	33	12
	119	123(T_2)		

* This work.

calculated frequencies is sufficiently good to support the above assignment, and shows that the higher-frequency band is due to the antisymmetric stretch (B_1 symmetry) and the lower-frequency band to the symmetric stretch (A_1 symmetry).

As expected from the comparison of bond lengths discussed above, the CuBr force constant in $[\text{Cu}(\text{PPh}_3)_2\text{Br}_2]^-$ is significantly lower than the value found for CuBr_2^- .¹³ This result parallels that found previously for $[\text{M}(\text{PPh}_3)_n\text{X}]$ ($\text{M} = \text{Cu}, \text{Ag}, \text{or Au}$; $\text{X} = \text{Cl}, \text{Br}, \text{or I}$) where the $\nu(\text{MX})$ frequency and the MX force constant are found to be strongly dependent on co-ordination number and MX bond length.^{3,11}

Table 3 contains other data for comparison with the above results. Thus, the force constant f_t for the terminal CuBr bond in $[\text{Cu}(\text{PPh}_3)_2\text{Br}]$ is significantly greater than that for the corresponding bonds in $[\text{Cu}(\text{PPh}_3)_2\text{Br}_2]^-$. This is in agreement with the trends in the CuBr bond lengths in these complexes, as discussed above. A similar normal-co-ordinate calculation was carried out on the Cu_2Br_2 core of $[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$. This analysis shows that the i.r. bands at 180 and 128 cm^{-1} can be satisfactorily accounted for only if the force constant for the Cu–Br bonds involving the trigonally co-ordinated copper atom is significantly greater than that involving the tetrahedrally co-ordinated copper atoms. From the results in Table 3 it can be seen that the force constant for the trigonal Cu–Br bonds in this complex is only slightly smaller than that for the corresponding bonds in $[\text{Cu}(\text{PPh}_3)_2\text{Br}_2]^-$. Again, this is in good agreement with observations made above concerning the relative bond lengths in these species. Likewise, the force constant for the Cu–Br bonds involving the tetrahedral copper atom in $[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$ is almost identical to that for the CuBr bonds in $[\text{Cu}(\text{PPh}_3)\text{Br}]_4$, which also involves tetrahedrally co-ordinated copper(I). Analysis of the potential-energy distribution for the four calculated $\nu(\text{CuBr})$ modes of $[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$ shows that the two higher-frequency modes are associated mainly with stretching of the CuBr bonds on the trigonal copper atom, while the two lower-frequency modes are associated with the tetrahedral copper atom. This result suggests an alternative explanation for the presence of two main bands in the far-i.r. spectrum of the 'step' $[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ (Table 3). This is that these two bands are due to the presence of trigonally and tetrahedrally co-ordinated copper atoms in the structure, rather than to the presence of doubly and triply bridging bromine atoms as postulated previously.^{12,14} These two descriptions are obviously related, but are not equivalent, as evidenced by the fact that $[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$ shows two well separated $\nu(\text{CuBr})$

**Figure 4.** Phosphorus-31 c.p.m.a.s. n.m.r. spectrum of $[\text{PPh}_3\text{Me}][\text{Cu}(\text{PPh}_3)\text{Br}_2]$

bands, despite the presence of only doubly bridging bromine atoms in the structure.

Solid-state ^{31}P N.M.R. Spectrum.—The solid-state c.p.m.a.s. ^{31}P n.m.r. spectrum of $[\text{PPh}_3\text{Me}][\text{Cu}(\text{PPh}_3)\text{Br}_2]$ is shown in Figure 4. It consists of a strong singlet at 21 p.p.m. (relative to 85% H_3PO_4) due to the phosphonium cation, and a weaker quartet centred at -14 p.p.m. due to the co-ordinated PPh_3 in the anion. As described previously, the quartet structure is due to spin-spin coupling of the phosphorus nucleus to the copper nucleus (^{63}Cu , ^{65}Cu ; $I = \frac{3}{2}$). The line spacings in the quartet are unequal, due to the presence of a nuclear quadrupole coupling interaction between the copper nucleus and its surroundings.^{3,4,15} For a spherically symmetric charge distribution about the copper nucleus, the nuclear quadrupole coupling constant would be zero, and the components of the quartet would be equally spaced. The marked asymmetry observed in the present case is thus a reflection of the electrical asymmetry about the copper nucleus which results from the relatively low-symmetry trigonal-planar co-ordination environment.

In order to investigate possible relationships between the ^{31}P n.m.r. parameters and the co-ordination environment about the copper atom, the solid-state ^{31}P n.m.r. data obtained in the present study are compared with those for some related species in Table 4. The average chemical shift for $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$ is more negative than those for the other bromocopper(I) complexes listed in Table 4. Since the last two compounds in this table also contain trigonal $(\text{PPh}_3)\text{CuBr}_2$ units, it appears that the chemical shift is not a good indicator of the co-ordination environment. The average line spacing $\langle\Delta\nu_i\rangle$ in the quartets increases with decreasing number of co-ordinated PPh_3 ligands and is usually found in the range 0.9–1.0 kHz for $\text{Cu}(\text{PPh}_3)_3$ compounds, 1.2–1.3 kHz for $\text{Cu}(\text{PPh}_3)_2$ compounds, and 1.6–1.8 kHz for $\text{Cu}(\text{PPh}_3)$ compounds. This trend presumably reflects the decrease in Cu–P bond lengths with decreasing number of co-ordinated ligands and a correspondingly greater degree of copper 4s-orbital participation in the Cu–P bonding. It is also notable that the asymmetry, $\Delta\nu_{ij}$, in the quartet splittings is approximately the same for the two trigonal mononuclear complexes $[\text{Cu}(\text{PPh}_3)_2\text{Br}]$ and $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$, but is much smaller for the tetrahedral mononuclear complex. This is in agreement with the expectation that the copper quadrupole coupling constant will be smaller for the tetrahedral than for the trigonal environment, as has been observed in related complexes by direct measurement of the nuclear

Table 4. C.p.m.a.s. solid-state ^{31}P n.m.r. parameters for some bromocopper(I) complexes^a

	δ_1	δ_2	δ_3	δ_4	$\langle\delta\rangle$	$\Delta\nu_1$	$\Delta\nu_2$	$\Delta\nu_3$	$\langle\Delta\nu_i\rangle$	$\Delta\nu_{ij}$	Ref.
$[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$	4.9	-6.4	-20.3	-36.1	-14	1.37	1.69	1.92	1.66	0.55	<i>b</i>
$[\text{Cu}(\text{PPh}_3)_2\text{Br}]$	8.0	0.8	-9.7	-21.6	-6	0.88	1.28	1.44	1.20	0.56	3
$[\text{Cu}(\text{PPh}_3)_3\text{Br}]$	6.5	-1.0	-9.0	-16.8	-5	0.90	0.96	0.95	0.94	0.06	4
$[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$	18.1	5.9	-8.1	-24.7	-2	1.48	1.70	2.02	1.73	0.54	<i>b</i>
	10.5	0.2	-11.2	-24.7	-6	1.25	1.40	1.65	1.43	0.40	
	12.6	4.0	-8.1	-19.2	-3	1.05	1.45	1.35	1.28	0.30	
$[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ 'step'	12.7	-0.9	-16.6	-33.0	-9	1.65	1.91	1.99	1.85	0.34	<i>b</i>
	15.9	4.1	-11.4	-28.4	-5	1.43	1.88	2.06	1.79	0.63	

^a δ_i are the chemical shifts (± 0.5 p.p.m.) with respect to 85% H_3PO_4 (δ for solid $\text{PPh}_3 = -9.9$ p.p.m.); $\langle\delta\rangle$ is the average chemical shift for each quartet; $\Delta\nu_i$ is the splitting in kHz (± 0.03) between each of the four peaks of the quartet; $\langle\Delta\nu_i\rangle$ is the average of the $\Delta\nu_i$ for each quartet; $\Delta\nu_{ij} = \Delta\nu_i(\text{max.}) - \Delta\nu_i(\text{min.})$ is the asymmetry parameter of the quartet. ^b This work.

quadrupole transition frequencies.^{16,17} The mean line spacing $\langle\Delta\nu_i\rangle$ and the quartet asymmetry parameters $\Delta\nu_{ij}$ for the unsolvated dinuclear and the tetranuclear aggregated complexes $[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$ and 'step' $[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ in Table 4, both of which contain trigonally and tetrahedrally co-ordinated copper atoms, also reflect the difference between the two kinds of co-ordination environment, although for the dinuclear complex, overlapping bands from the three quartets makes accurate determination of the peak positions difficult. Copper nuclear quadrupole resonance frequencies have been measured for the trigonal and tetrahedral sites in $[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$, and these are in this ratio 2:1.^{16,17} This suggests that the copper nuclear quadrupole coupling constants are in approximately the same ratio. The quartet at -2 p.p.m. in the solid-state ^{31}P n.m.r. spectrum of this complex has $\Delta\nu_{ij} = 0.54$ kHz, which is almost the same as the values for the two trigonally co-ordinated mononuclear complexes $[\text{Cu}(\text{PPh}_3)_2\text{Br}]$ and $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$. The $\langle\Delta\nu_i\rangle$ of 1.73 kHz is in the range expected for the $\text{Cu}(\text{PPh}_3)$ fragment and this quartet is thus assigned to the phosphorus atom bound to the trigonal copper site in the complex. The other two quartets are assigned to the two phosphorus atoms co-ordinated to the tetrahedral copper site on the basis of the smaller values of $\langle\Delta\nu_i\rangle$ and $\Delta\nu_{ij}$. The $\Delta\nu_{ij}$ values are, however, larger than that for the phosphorus atoms in the tetrahedrally co-ordinated mononuclear complex $[\text{Cu}(\text{PPh}_3)_3\text{Br}]$. This suggests that the copper quadrupole coupling constants are quite different for the two different co-ordination environments (P_2CuBr_2 and P_3CuBr) involved, despite the fact that the co-ordination number is four in both cases. In the 'step' form of $[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ the nuclear quadrupole resonance frequency of the trigonally co-ordinated copper atom has been measured,¹⁶ and is almost identical to that of the trigonal copper atom in $[(\text{Ph}_3\text{P})\text{CuBr}_2\text{Cu}(\text{PPh}_3)_2]$ and one quartet for the unsolvated 'step' $[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ has a $\Delta\nu_{ij}$ which is comparable in magnitude to the values for the trigonal sites in the other compounds in Table 4 and the other to the tetrahedral PCuBr_3 site.

While these results provide a consistent relationship between the structural parameters of the $[\text{Cu}(\text{PPh}_3)_m\text{X}]_n$ system and the n.m.r. parameters $\Delta\nu_i$ and $\Delta\nu_{ij}$, as a cautionary note we report that for the spectra of solvated or partially solvated forms of the 'step' $[\text{Cu}(\text{PPh}_3)\text{Br}]_4$ ¹⁸ the differences in these parameters for the trigonal and tetrahedral copper sites is much less marked, and more work is still required to elucidate the origins of these effects.

Conclusions

The present study establishes the existence of the first anionic mixed-ligand complexes of copper(I) with halide and triphenyl-

phosphine. The isolation of $[\text{Cu}(\text{PPh}_3)\text{Br}_2]^-$ in the presence of excess of PPh_3 suggests that CuBr_2 is only able to accommodate one PPh_3 molecule in its co-ordination sphere. Trends in metal-ligand bond lengths, ^{31}P n.m.r. line spacings, and bond-stretching force constants between this and related species can be understood on the basis of a greater σ -donor capacity for PPh_3 relative to Br^- , and establish a close similarity between the $(\text{PPh}_3)\text{CuBr}_2$ unit in various structures, irrespective of whether the Br atoms are terminal or bridging.

Acknowledgements

We thank Dr. A. Camus for suggesting the investigation of the type of complex reported here, and Mrs. Catherine Hobbs for assistance with the preparative work. The Bruker CXP-300 spectrometer is operated by the Brisbane N.M.R. Centre and we thank the Centre for making instrument time available. We thank Mr. David Tucker for recording the solid-state ^{31}P spectra and Ms. Candiera Albert for assistance in the preparation of some of the neutral complexes used. We are grateful to the New Zealand Universities Grants Committee and the Australian Research Grants Scheme for financial assistance.

References

- G. A. Bowmaker, *Adv. Spectrosc.*, 1987, **14**, 1.
- G. A. Bowmaker, G. R. Clark, D. A. Rogers, A. Camus, and N. Marsich, *J. Chem. Soc., Dalton Trans.*, 1984, 37.
- G. A. Bowmaker, J. C. Dyason, P. C. Healy, L. M. Engelhardt, C. Pakawatchai, and A. H. White, *J. Chem. Soc., Dalton Trans.*, 1987, 1089.
- P. F. Barron, J. C. Dyason, P. C. Healy, L. M. Engelhardt, C. Pakawatchai, V. A. Patrick, and A. H. White, *J. Chem. Soc., Dalton Trans.*, 1987, 1099.
- J. C. Dyason, L. M. Engelhardt, C. Pakawatchai, P. C. Healy, and A. H. White, *Aust. J. Chem.*, 1985, **38**, 1243.
- 'International Tables for X-Ray Crystallography,' eds. J. A. Ibers and W. C. Hamilton, Kynoch Press, Birmingham, 1974, vol. 4.
- S. R. Hall, The XTAL User Manual, Technical Report of the Computer Science Centre, University of Maryland, U.S.A., 1983.
- S. Anderson and S. Jagner, *Acta Chem. Scand., Ser. A*, 1985, **39**, 297, 515, 577; 1986, **40**, 210; M. Asplund, S. Jagner, and M. Nilsson, *ibid.*, 1983, **37**, 57.
- R. P. Shibaeva, V. F. Kaminskii, E. B. Yagubskii, and L. A. Kushch, *Sov. Phys. Crystallogr.*, 1983, **28**, 48.
- J. C. Dyason, L. M. Engelhardt, P. C. Healy, H. L. Klich, and A. H. White, *Aust. J. Chem.*, 1986, **39**, 2003.
- G. A. Bowmaker and D. A. Rogers, *J. Chem. Soc., Dalton Trans.*, 1984, 1249.
- G. A. Bowmaker and P. C. Healy, *Spectrochim. Acta, Part A*, 1988, **44**, 115.

- 13 G. A. Bowmaker, L. D. Brockliss, and R. Whiting, *Aust. J. Chem.*, 1973, **26**, 29.
14 B-K. Teo and D. M. Barnes, *Inorg. Nucl. Chem. Lett.*, 1973, **12**, 681.
15 E. M. Menger and W. S. Veeman, *J. Magn. Reson.*, 1982, **46**, 257.
16 T. Okuda, M. Hiura, K. Yamada, and H. Negita, *Chem. Lett.*, 1977, 367.
17 H. Negita, M. Hiura, K. Yamada, and T. Okuda, *J. Mol. Struct.*, 1980, **58**, 205.
18 P. F. Barron, J. C. Dyason, L. M. Engelhardt, P. C. Healy, and A. H. White, *Inorg. Chem.*, 1984, **23**, 3766.

Received 22nd February 1988; Paper 8/00687C