

TWO CARBOCYCLIC LOCKED NUCLEIC ACID ANALOGUES GIVE STRUCTURAL INFORMATION ABOUT THE ROLE OF HYDRATION IN A-TYPE DUPLEXES

Nanna Albæk, Michael Petersen, and Poul Nielsen □ *Nucleic Acid Center,
Department of Chemistry and Physics, University of Southern Denmark, Odense M, Denmark*

□ *Two locked nucleic acid (LNA) analogues with three-carbon 2'-4' linkages, saturated or unsaturated, are synthesized using a ring-closing metathesis based strategy. Strongly stabilized duplexes with complementary RNA and slightly destabilized duplexes with complementary DNA are observed. CD-spectroscopy indicates a less pronounced shift toward A-type duplexes compared to LNA. These results combining a strong N-type conformation with the absence of a 2'-oxygen demonstrate a stronger importance of minor groove hydration in an intermediate duplex type than in an A-type duplex.*

Keywords Locked nucleic acid; RNA recognition; duplex hydration

Locked nucleic acid (LNA)^[1–3] and its analogue ENA^[4] are among the most intriguing nucleic acid analogues due to their unprecedented affinity for complementary RNA and DNA sequences. The LNA and ENA monomers (Figure 1) are both locked in the N-type conformation. A single incorporation of an LNA monomer in an oligodeoxynucleotide (ODN) leads to an increase in the thermal stability of 3–8°C of a DNA:RNA duplex and 3–5°C of a DNA:DNA duplex.^[2] This behavior is related to the ability of LNA-monomers to conformationally steer their neighboring 2'-deoxynucleotides into an N-type conformation and hereby to induce A-type or A-type like duplex conformation.^[3]

ENA demonstrates almost the same stabilisation of duplexes as LNA,^[4] whereas other analogues in which the 4'-2' linkage has another constitution (i.e., a “COC” or an even longer “COCO” or “CCCO”) demonstrates somewhat less pronounced stabilization of a DNA:RNA duplex and a neutral or even destabilizing effect on DNA:DNA duplexes.^[4,5] The 4'-2'-linkage with three carbon atoms, however, has been only briefly mentioned in a review^[6]

The Danish National Research Foundation, the Danish Natural Research Council, and the Danish Centre for Scientific Computing are thanked for financial support.

Address correspondence to Poul Nielsen, Nucleic Acid Center, Dept of Chemistry, University of Southern Denmark, DK-5230 Odense, Denmark. E-mail: pon@ifk.sdu.dk





FIGURE 1 LNA and ENA and two new carbocyclic analogues.

and reported to give a destabilized DNA:RNA duplex which is a surprising result compared to the results of the other analogues. We, therefore, decided to synthesise this analogue, **X**, by a ring-closing metathesis based strategy giving access also to the unsaturated analogue **Y** (Figure 1).

The synthesis started from uridine following a linear approach (Scheme 1) and took advantage of the reported selective 5',3'-silylation^[7]



SCHEME 1 a) TBSCl, DABCO, AgNO₃, THF, 90%, ref. 7; b) PhOCSCl, DMAP, CH₃CN, 79%; c) AllylSnBn₃, AIBN, toluene, 65%, ref. 8; d) aq. AcOH, 75%; e) i) Dess-Martin periodinane, CH₂Cl₂; ii) H₂CO, MeOH, aq. NaOH, dioxane then NaBH₄, 52%; f) i) BzCl, pyridine, CH₃CN, 65%; ii) TBSCl, Im., DMF; iii) NaOMe, MeOH, 71%; g) Dess-Martin periodinane, CH₂Cl₂, 93%; h) Ph₃PCH₃Br, BuLi, THF, 95%; i) **A**, CH₂Cl₂, 96%; j) TBAF, THF, 98%; k) H₂, PtO₂, MeOH, 100%; l) DMTCl, pyridine, CH₃CN, 78%; m) NC(CH₂)₂OPCIN(iPr)₂, DIPEA, CH₂Cl₂ 63%; n) DMTCl, Et₃N, CH₂Cl₂, 78%; o) NC(CH₂)₂OPCIN(iPr)₂, DIPEA, CH₂Cl₂ 57%.

TABLE 1 Hybridization data.

ODN sequence	$T_m (\Delta T_m \text{ per modification}) / ^\circ\text{C}$	
	Complementary DNA	Complementary RNA
5'-dGTGATATGC-3'	30.0	27.5
5'-dGTGAXATGC-3'	29.0 (−1.0)	32.0 (+4.5)
5'-dGXXGAXXGC-3'	27.0 (−1.0)	38.0 (+3.5)
5'-dGTGAYATGC-3'	29.0 (−1.0)	31.5 (+4.0)
5'-dGYGAYAYGC-3'	23.0 (−2.3)	34.5 (+2.3)

Buffer: 5 mM Na₂HPO₄, 10 mM NaH₂PO₄, 100 mM NaCl, 0.1 mM EDTA, pH 7.0; 1.5 μM of each strand.

and 2'-allylation,^[8] protecting group manipulations, a 4'-vinylation and a very efficient ring-closing metathesis using Grubbs second generation catalyst **A** to give the analogue **Y**. Hydrogenation gave **X** and both analogues were converted into DMT-protected phosphoramidites and incorporated into ODN's (Table 1)

The ODNs were mixed with complementary DNA and RNA sequences and the stability of the duplexes were determined. Significant increases in stability of the DNA:RNA duplexes were observed. These are only slightly smaller than the increases found for LNA and ENA. On the other hand, slight decreases in stability of the DNA:DNA duplexes were observed. This is different to the data obtained with LNA and ENA for which increased stability was found also for DNA:DNA sequences. CD-spectra (not shown) demonstrate that the shift toward A-type (for DNA:RNA) or A-type like duplexes (for DNA:DNA) is smaller for the duplexes containing **X** or **Y** than for comparable LNA sequences. In other words, the conformational steering on neighbouring 2'-deoxynucleotides is smaller for **X** and **Y** than it is for LNA (and probably also ENA) nucleotides. This might be due to the lack of a 2'-oxygen atom in the former hampering the hydration pattern in the minor groove of the duplexes. Apparently, this hydration pattern is more important in intermediate type DNA:DNA duplexes with a narrow minor groove than in the compact A-type DNA:RNA duplexes with a broader minor groove.

REFERENCES

1. The present results are also described in Albæk, N.; Petersen, M.; Nielsen, P. Analogues of locked nucleic acid with three carbon 2',4'-linkages; Synthesis by ring-closing metathesis and influence on nucleic acid duplex stability and structure. *J. Org. Chem.* **2006**, 71, 7731–7740.
2. a) Singh S.K.; Nielsen P.; Koshkin A.A.; Wengel J.; LNA (locked nucleic acids): synthesis and high-affinity nucleic acid recognition. *Chem. Comm.* **1998**, 455–456; b) Obika S.; Nanbu D.; Hari Y.; Andoh J.; Morio K.; Doi T.; Imanishi T. Stability and structural features of the duplexes containing nucleoside analogues with a fixed N-type conformation, 2'-O, 4'-C-methyleneribonucleosides. *Tetrahedron Lett.* **1998**, 39, 5401–5405.

3. Petersen, M.; Bondensgaard, K.; Wengel, J.; Jacobsen, J.P. Locked nucleic acid (LNA) recognition of RNA: NMR solution structures of LNA:RNA hybrids. *J. Am. Chem. Soc.* **2002**, 124, 5974–5982.
4. Morita, K.; Takagi, M.; Hasegawa, C.; Kaneko, M.; Tsutsumi, S.; Sone, J.; Ishikawa, T.; Imanishi, T.; Koizumi, M. Synthesis and properties of 2'-O,4'-C-Ethylene-Bridged nucleic acids (ENA) as effective antisense oligonucleotides. *Bioorg. Med. Chem.* **2003**, 11, 2211–2226.
5. a) Wang, G.; Gunic, E.; Girardet, J.-L.; Stoisavljevic, V.; Conformationally locked nucleosides. Synthesis and hybridization properties of oligodeoxynucleotides containing 2',4'-C-bridged 2'-deoxynucleosides, *Bioorg. Med. Chem. Lett.* **1999**, 9, 1147–1150; b) Hari, Y.; Obika, S.; Ohnishi, R.; Eguchi, K.; Osaki, T.; Ohishi, H.; Imanishi, T. Synthesis and properties of 2'-O, 4'-C-methyleneoxymethylene bridged nucleic acid. *Bioorg. Med. Chem.* **2006**, 14, 1029–1038.
6. Freier, S.M.; Altmann, K.-H. The ups and downs of nucleic acid duplex stability: structure-stability studies on chemically-modified DNA:RNA duplexes. *Nucleic Acids Res.* **1997**, 25, 4429–4443.
7. Hakimelahi, G.H.; Proba, Z.A.; Ogilvie, K.K. New catalysts and procedures for the dimethoxytritylation and selective silylation of ribonucleosides. *Can. J. Chem.* **1982**, 60, 1106–1113.
8. Tronchet, J.M.J.; Grand, E.; Zsely, M.; Giovannini, R.; Geoffroy, M. Spin labeled dinucleosides, *Carbohydrate Lett.* **1998**, 3, 161–168.

Copyright of Nucleosides, Nucleotides & Nucleic Acids is the property of Taylor & Francis Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.