

“Click”-Capture, Ring-Opening Metathesis Polymerization (ROMP), Release: Facile Triazolation Utilizing ROMP-Derived Oligomeric Phosphates


Toby R. Long,[†] Saqib Faisal,^{†,‡} Pradip K. Maity,[†] Alan Rolfe,[†] Ryan Kurtz,[†]
Sarra V. Klimberg,[†] Muhammad-Rabbie Najjar,[†] Fatima Z. Basha,[‡] and Paul R. Hanson^{*,†}

Department of Chemistry, University of Kansas, 1251 Wescoe Hall Drive, Lawrence, Kansas 66045, United States, The University of Kansas Center for Chemical Methodologies and Library Development (KU-CMLD), 2034 Becker Drive, Delbert M. Shankel Structural Biology Center, Lawrence, Kansas 66047, United States, and H. E. J. Research Institute of Chemistry, International Center for Chemical and Biological Science, University of Karachi, Karachi, Pakistan

phanson@ku.edu

Received February 16, 2011

ABSTRACT



Soluble, high-load ring-opening metathesis polymerization (ROMP)-derived oligomeric triazole phosphates (OTP) are reported for application as efficient triazolating reagents of nucleophilic species. Utilizing a “Click”-capture, ROMP, release protocol, the efficient and purification-free, direct triazolation of *N*-, *O*-, and *S*-nucleophilic species was successfully achieved. A variety of OTP derivatives were rapidly synthesized as free-flowing solids on a multigram scale from commercially available materials.

Rapid access to collections of diverse molecules in desirable quantities and purity for high-throughput

screening is an important challenge in drug discovery. In this regard, methods that integrate synthesis and purification have become powerful tools in the arena of facilitated synthesis.^{1,2} This has been driven in recent years by the development of an array of polymer-bound reagents and scavengers for utilization in streamlined protocols to access small molecule libraries.² Despite huge advances in this area, limitations in nonlinear reaction kinetics (heterogeneous reactions), low resin-load capacities, means of distributing reagents, and solution phase automation technology continue to warrant the development of designer polymers for library production.³ To this effect,

[†] University of Kansas.

[‡] University of Karachi.

(1) (a) Booth, R. J.; Hodges, J. C. *Acc. Chem. Res.* **1999**, *32*, 18–26. (b) Ley, S. V.; Baxendale, I. R.; Bream, R. N.; Jackson, P. S.; Leach, A. G.; Longbottom, D. A.; Nesi, M.; Scott, J. S.; Storer, R. I.; Taylor, S. J. *J. Chem. Soc., Perkin Trans. 1* **2000**, 3815–4195. (c) Kirschning, A.; Monenschein, H.; Wittenberg, R. *Angew. Chem., Int. Ed.* **2001**, *40*, 650–679. (d) Eames, J.; Watkinson, M. *Eur. J. Org. Chem.* **2001**, 1213–1224. (e) Strohmeier, G. A.; Kappe, C. O. *Angew. Chem., Int. Ed.* **2004**, *43*, 621–624. (f) Lesch, B.; Thomson, D. W.; Lindell, S. D. *Comb. Chem. High. Throughput Screening* **2008**, *11*, 31–36.

(2) For reviews concerning soluble polymers, see: (a) Gravert, D. J.; Janda, K. D. *Chem. Rev.* **1997**, *97*, 489–509. (b) Toy, P. H.; Janda, K. D. *Acc. Chem. Res.* **2000**, *33*, 546–554. (c) Dickerson, T. J.; Reed, N. N.; Janda, K. D. *Chem. Rev.* **2002**, *102*, 3325–3344. (d) Haag, R. *Chem.—Eur. J.* **2001**, *7*, 327–335. (e) Haag, R.; Sunder, A.; Hebel, A.; Roller, S. *J. Comb. Chem.* **2002**, *4*, 112–119. (f) Bergbreiter, D. E. *Chem. Rev.* **2002**, *102*, 3345–3384. (g) Bergbreiter, D. E.; Tian, J.; Hongfa, C. *Chem. Rev.* **2009**, *109*, 530–582.

(3) (a) Studer, A.; Curran, D. P. *Tetrahedron* **1997**, *53*, 6681–6696. (b) Hjerten, S.; Li, Y.-M.; Liao, J. L.; Mankazato, K.; Mohammad, J.; Pettersson, G. *Nature* **1992**, *356*, 810–811. (c) Baumann, M.; Baxendale, I. R.; Ley, S. V.; Nikbin, N.; Smith, C. D. *Org. Biomol. Chem.* **2008**, *6*, 1577–1586.

a variety of reagents and scavengers possessing tunable properties have emerged from ring-opening metathesis polymerization (ROMP) technology.^{4–6} We herein report the development of a new ROMP-derived oligomeric triazole phosphate (OTP) for application as a soluble, efficient triazolating reagent of nucleophilic species. Overall, these reagents are free-flowing solids that are easy to handle, nontoxic, soluble, and air stable. In addition, they are easily stored long-term at room temperature and are readily prepared, vide infra, from commercially available starting materials on a multigram scale.

The development of new methodologies for the diversification of biologically interesting core scaffolds with functional handles is of paramount importance. In this regard, triazoles and their derivatives have demonstrated a wide variety of biological activity, with many reports focusing on antifungal activity.⁷ Despite this activity, the utilization of solution phase or immobilized reagents to directly triazolate nucleophilic species in a one-step protocol has been limited to reports of a two-step, one-pot propargylation-click protocol.⁸

Oligomeric and polyphosphates are ideal immobilized leaving groups due to the inherent pK_a , stability and innate leaving group properties of phosphate anions.^{9–11} Recently, we reported the generation and application of

oligomeric benzyl phosphate (OBP) as an efficient benzylating reagent.¹⁰ We now report the synthesis of ROMP-derived triazolating reagents (OTP) for application in purification-free diversifications of nucleophilic species using the title method, termed “Click”-Capture, ROMP, Release. This method utilizes a propargyl-tagged norbornenyl-phosphate to capture an azide in a classical “click” reaction, followed by ROM polymerization to generate the desired soluble oligomeric triazole reagent (OTP) **4**. Subsequent release via S_N2 displacement with nucleophilic species yields triazolated products along with the spent oligomeric phosphate that is readily sequestered via precipitation (Figure 1).





Figure 1. Reaction of oligomeric triazole phosphate (OTP) **4**.

The synthesis of the oligomeric triazole phosphate bearing a 4-MeOPh group, OTP **4a**, starts with the exonorbornenyl tagged (Nb-tagged) phosphonyl chloride **1** utilized in the synthesis of previously reported ROMP-derived benzylating reagent OBP.^{10,12} Phosphorylation of propargyl alcohol with Nb-tagged phosphonyl chloride **1**, followed by a “Click”-capture event of the corresponding azide, yields the desired monomer **3a** in an efficient fashion. ROM polymerization of monomer **3a** was achieved with $RuCl_2(PCy_3)_2=CHPh$ (cat. **A**), followed by basic workup


Scheme 1. Synthesis of Oligomeric Triazole Phosphate (OTP) **4a**



- (4) (a) Barrett, A. G. M.; Hopkins, B. T.; Kobberling, J. *Chem. Rev.* **2002**, *102*, 3301–3324. (b) Harned, A. M.; Probst, D. A.; Hanson, P. R. In *Handbook of Metathesis*; Grubbs, R. H., Ed.; Wiley-VCH: Weinheim, 2003; Vol. 2, pp 361–402. (c) Flynn, D. L.; Hanson, P. R.; Berk, S. C.; Makara, G. M. *Curr. Opin. Drug Discovery Dev.* **2002**, *5*, 571–579. (d) Harned, A. M.; Zhang, M.; Vedantham, P.; Mukherjee, S.; Herpel, R. H.; Flynn, D. L.; Hanson, P. R. *Aldrichimica Acta* **2005**, *38*, 3–16.
- (5) (a) Rolfe, A.; Loh, J. K.; Maity, P. K.; Hanson, P. R. *Org. Lett.* **2011**, *1*, 4–7. (b) Maity, P.; Rolfe, A.; Samarakoon, T. B.; Faisal, S.; Kurtz, R.; Long, T. R.; Schätz, A.; Flynn, D.; Grass, R. N.; Stark, W. J.; Reiser, O.; Hanson, P. R. *Org. Lett.* **2011**, *1*, 8–10. (c) Rolfe, A.; Probst, D.; Volp, K.; Omar, I.; Flynn, D.; Hanson, P. R. *J. Org. Chem.* **2008**, *73*, 8785–8790. (d) Stoianova, D. S.; Yao, L.; Rolfe, A.; Samarakoon, T.; Hanson, P. R. *Tetrahedron Lett.* **2008**, *49*, 4553–4555. (e) Zhang, M.; Flynn, D. L.; Hanson, P. R. *J. Org. Chem.* **2007**, *72*, 3194–3198. (f) Roberts, R. S. *J. Comb. Chem.* **2005**, *7*, 21–32. (g) Harned, A. M.; Sherrill, W. M.; Flynn, D. L.; Hanson, P. R. *Tetrahedron* **2005**, *61*, 12093–12099. (h) Arstad, E.; Barrett, A. G. M.; Tedeschi, L. *Tetrahedron Lett.* **2003**, *44*, 2703–2707. (i) Barrett, A. G. M.; Hopkins, B. T.; Love, A. C.; Tedeschi, L. *Org. Lett.* **2004**, *6*, 835–837.
- (6) Vedantham, P.; Zhang, M.; Gor, P. J.; Huang, M.; Georg, G. I.; Lushington, G. H.; Mitscher, L. A.; Ye, Q.-Z.; Hanson, P. R. *J. Comb. Chem.* **2008**, *10*, 195–203.
- (7) (a) Xia, Y.; Qu, F.; Peng, L. *Rev. Med. Chem.* **2010**, *10*, 806–821. (b) Cronin, S.; Chandrasekar, P. H. *J. Antimicrob. Chemother.* **2010**, *65*, 410–416. (c) Nivoix, Y.; Ubeaud-Sequier, G.; Engel, P.; Leveque, D.; Herbrecht, R. *Curr. Drug Metab.* **2009**, *10*, 395–409.
- (8) (a) Inverarity, I. A.; Hulme, A. N. *Org. Biomol. Chem.* **2007**, *5*, 636–643. (b) Smith, G.; Glaser, M.; Perumal, M.; Nguyen, Q.-D.; Shan, B.; Aarstad, E.; Aboagye, E. O. *J. Med. Chem.* **2008**, *51*, 8057–8067.
- (9) (a) Manfredini, T.; Pellacani, G. C.; Pozzi, P.; Corradi, A. B. *Appl. Clay Sci.* **1990**, *5*, 193–201. (b) Best, M. D.; Zhang, H.; Prestwich, G. D. *Nat. Prod. Rep.* **2010**, *27*, 1403–1430. (c) Fylaktakidou, K. C.; Duarte, C. D.; Koumbis, A. E.; Nicolau, C.; Lehn, J.-M. *Chem. Med. Chem.* **2011**, *6*, 153–168 and references cited therein.
- (10) Long, T. R.; Maity, P. K.; Samarakoon, T. B.; Hanson, P. R. *Org. Lett.* **2010**, *12*, 2904–2907 and references cited therein.
- (11) (a) Nicolaou, K. C.; Shi, G. Q.; Gunzner, J. L.; Gaertner, P.; Yang, Z. *J. Am. Chem. Soc.* **1997**, *119*, 5467–5468. (b) La Cruz, T. E.; Rychnovsky, S. D. *J. Org. Chem.* **2007**, *72*, 2602–2611. (c) Lapointe, D.; Fagnou, K. *Org. Lett.* **2009**, *11*, 4160–4163. (d) Westheimer, F. H. *Science* **1987**, *236*, 1173–1178. (e) Thomas, C. D.; Hanson, P. R. In *Metathesis in Natural Product Synthesis*; Cossy, J.; Areniyadis, S.; Meyer, C., Ed.; Wiley-VCH: Weinheim, 2010; pp 129–144. (f) Beaucage, S. L.; Caruthers, M. H. *Bioorganic Chemistry: Nucleic Acids*; Hecht, S. M., Ed.; Oxford University Press: New York, 1996; Chapter 2, pp 36–74.


utilizing the Pederson protocol.^{13,14} Precipitation via drop-wise addition into anhydrous Et₂O afforded the corresponding oligomeric triazole phosphate (OTP₂₀) **4a** as a free-flowing white solid possessing a theoretical load of 2.4 mmol/g (Scheme 1).

Scheme 2. Triazolation of 2,4-Cl-PhOH Utilizing OTP **4a**



Investigation into the utilization of OTP **4a** as a direct triazolating reagent was next studied using reaction conditions reported for the application of OBP.¹⁰ After optimization of reaction conditions for the triazolation of 2,4-dichlorophenol utilizing OTP **4a**, the corresponding triazole ether, **5a**, was isolated in excellent yield (99%) and crude purity (>90%) using simple filtration through a Celite SPE (Scheme 2).

Table 1. Triazolation of *N*-, *O*-, and *S*-Nucleophiles Using OTP₂₀ **4a**^a




entry	nucleophile	pdt	yield (%) ^b
1	2,4-Cl-PhOH	5a	98
2	4-F-PhOH	5b	92
3	4- ^t Bu-PhOH	5c	90
4	naphthalene-1-ol	5d	69
5	4-SMe-PhSH	5e	60
6	morpholine	5f	72
7	thiomorpholine	5g	75
8	1-phenylpiperazine	5h	95
9	indoline	5i	88
10	<i>N</i> -ethylnaphthalen-1-amine	5j	62

^a OTP **4a** utilized as a 20-mer. ^b Purities >90% observed for all reactions using both GC and ¹H NMR.

The application of OTP **4a** as an efficient triazolating reagent was extended to a variety of *N*-, *O*-, and *S*-nucleophilic species (Table 1). Initially, a variety of phenols were utilized (Table 1, entries 1–3) though reduced yields were observed for sterically hindered naphthalene-1-ol.

In addition to phenols, thiophenols (Table 1, entry 5) and a variety of amines (Table 1, entries 6–10) were successfully utilized to release the corresponding triazole in >90%


Table 2. Synthesis of Various OTP Analogues **4**



monomer	R ¹	pdt	yield (%) ^a
3a	4-OMe-Ph	4a	82
3b	4-Me-Ph	4b	88
3c	2-Me-Ph	4c	77
3d	4-Cl-Ph	4d	71
3e	4-F-Ph	4e	74
3f	4-CF ₃ -Ph	4f	73
3g	Cyclohexyl	4g	70
3h	4-Br-Ph	4h	89
3i	Furfuryl	4i	56

^a Yields corresponding to metathesis of corresponding monomers.

Table 3. Triazolation Utilizing OTP Derivatives **4a–g**^a



entry	nucleophile	OTP	pdt	yield (%) ^b
1	naphthalene-1-ol	4a	6a	72
2	naphthalene-1-ol	4b	6b	90
3	naphthalene-1-ol	4c	6c	55
4	naphthalene-1-ol	4d	6d	68
5	naphthalene-1-ol	4e	6e	70
6	naphthalene-1-ol	4f	6f	65
7	naphthalene-1-ol	4g	6g	49
8	<i>N</i> -ethylnaphthalen-1-amine	4a	6h	62
9	<i>N</i> -ethylnaphthalen-1-amine	4b	6i	52
10	<i>N</i> -ethylnaphthalen-1-amine	4c	6j	63
11	<i>N</i> -ethylnaphthalen-1-amine	4d	6k	51
12	<i>N</i> -ethylnaphthalen-1-amine	4e	6l	50
13	<i>N</i> -ethylnaphthalen-1-amine	4f	6m	60
14	<i>N</i> -ethylnaphthalen-1-amine	4g	6n	66

^a OTP **4a–g** utilized as a 20-mer. ^b Purities >90% observed for all reactions using both GC and ¹H NMR.

crude purity. Building on the success of OTP **4a**, a variety of additional OTP derivatives **4b–4i** were synthesized as free-flowing powders on a gram scale from ROM polymerization of their corresponding monomers utilizing cat. **A** (Table 2).

With a variety of OTP **4** derivatives in hand, the triazolation of both naphthalene-1-ol and *N*-ethylnaphthalen-1-amine with OTP derivatives **4a–g** was investigated (Table 3). All reactions proceeded with good yields with >90% crude purity after Celite SPE to remove the spent oligomer.

In conclusion, we have developed and demonstrated the synthesis and utilization of oligomeric triazole phosphates for direct triazolation of *N*-, *O*- and *S*-nucleophilic species in a “Click”-capture, ROMP, release protocol. These oligomeric reagents are readily synthesized on a multigram scale from commercially available materials as soluble, high-load, free-flowing powders. The application of OTP

in the diversification of core scaffolds for the synthesis of diverse collections of small molecules is underway and will be reported in due time.

Acknowledgment. This investigation was generously supported by the National Institute of General Medical Sciences (NIH P050-GM069663 and NIH-STTR R41 GM076765) with additional funds from the State of Kansas. We would like to also thank Materia, Inc. for supplying the metathesis catalyst and helpful discussions.

Supporting Information Available. Detailed experimental procedures and tabulated ¹H NMR, ¹³C NMR, ³¹P NMR, FTIR, and mass data and ¹H NMR spectra of products obtained by the described methods. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(12) Craig, D. *J. Am. Chem. Soc.* **1951**, *73*, 4889–4892.

(13) (a) Schwab, P.; Grubbs, R. H.; Ziller, J. W. *J. Am. Chem. Soc.* **1996**, *118*, 100–110. (b) Schwab, P.; France, M. B.; Ziller, J. W.; Grubbs, R. H. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 2039–2041.

(14) Pederson, R. L.; Fellows, I. M.; Ung, T. A.; Ishihara, H.; Hajela, S. P. *Adv. Synth. Catal.* **2002**, *344*, 728–735.