

the relative energy of the metal filled d orbitals of the appropriate π symmetry from a position they would otherwise have. This closes the energy separation between the filled metal d orbitals and CO π^* orbitals, which in turn promotes extensive metal to carbonyl π bonding.¹⁴ The values of $\nu(\text{CO})$ and Mo-C and C-O distances are thus apparently inconsistent with what would be expected based solely on the formal oxidation state of the metal. A similar argument can and has been used to account for the exceedingly low values of $\nu(\text{NO})$ found in the complexes $\text{Mo}_2(\text{OPr}^i)_6(\text{NO})_2$ (1630 cm^{-1}) and $\text{W}(\text{O}-\text{Bu}^i)_3(\text{py})(\text{NO})$ (1560 cm^{-1}) which contain linear M-N-O groups.¹⁵ Here terminal RO⁻ groups may push up, by the presence of their filled π -type orbitals, the relative energies of the filled metal d_{xz}, d_{yz} orbitals. This in turn promotes metal (d_{xz}, d_{yz})⁴ to NO π^* bonding.

Acknowledgments. We thank the donors of the Petroleum Research Fund, administered by the American Chemical Society, the National Science Foundation, the Marshall H. Wrubel Computing Center, and the taxpayers of the State of Indiana for financial support of this work. We also appreciate the valuable and stimulating discussions we have had with Professor F. A. Cotton during the course of this work.

Supplementary Material Available: A table of observed and calculated structure factors (47 pages).¹⁶ Ordering information is given on any current masthead page.

References and Notes

- (1) M. H. Chisholm, F. A. Cotton, M. W. Extine, and R. L. Kelly, *J. Am. Chem. Soc.*, **100**, 2256 (1978); (b) *ibid.*, submitted for publication.
- (2) M. H. Chisholm, F. A. Cotton, M. W. Extine, and W. W. Reichert, *Inorg. Chem.*, **17**, 2944 (1978).
- (3) Satisfactory elemental analyses have been obtained for both of the new compounds reported herein.
- (4) By the method of direct insertion using an AEI MS9 spectrometer at +6 °C (7×10^{-8} Torr). We thank Mr. Peter Cook and Professor D. C. Bradley for kindly obtaining this valuable spectral data for us.
- (5) Or at least were totally surprising and had no precedent to our knowledge in metal carbonyl chemistry.
- (6) This compound is insoluble in hexane and, though soluble in toluene, reacts rapidly to give $\text{Mo}(\text{CO})_6$ and as yet unidentified oxidized molybdenum compounds. This has prevented any characterization of the solution properties of $\text{Mo}(\text{OBU}^i)_2(\text{py})_2(\text{CO})_2$. All reactions were carried out under a dried and O₂-free nitrogen atmosphere.
- (7) Crystal data collected at -175 °C using a gaseous nitrogen cold stream and locally constructed goniostat system: monoclinic; space group $P2_1/c$ with unit cell dimensions $a = 12.738$ (2), $b = 10.330$ (5), $c = 16.328$ (11) Å; $\beta = 93.86$ (3)°; $Z = 4$. A total of 4919 unique data points were collected of which 4653 had $I > \sigma(I)$ and were used in the refinement $R(F) = 0.0433$, $R_w(F) = 0.082$.
- (8) See values quoted for compounds containing the *cis*- $\text{Mo}^{2+}(\text{CO})_2$ moiety in G. J. J. Chen, R. O. Yelton, and J. W. McDonald, *Inorg. Chim. Acta*, **22**, 249 (1977), and references therein.
- (9) F. A. Cotton and C. S. Kraihanzel, *J. Am. Chem. Soc.*, **84**, 4432 (1962).
- (10) F. A. Cotton, *Inorg. Chem.*, **3**, 702 (1964). In this paper a number of related $(\text{dien})\text{Mo}(\text{CO})_3$ ($\text{M} = \text{Cr}, \text{Mo}, \text{W}$) are discussed. The data quoted for $(\text{dien})\text{Mo}(\text{CO})_3$ was originally reported by C. S. Kraihanzel and F. A. Cotton, *ibid.*, **2**, 533 (1963), and were obtained in CH_3NO_2 solution. The stretch-stretch coupling interaction in $\text{Mo}(\text{OBU}^i)_2(\text{CO})_2(\text{py})_2$, which yields a separation of the two bands by 140 cm^{-1} , is all the more remarkable when compared with the separation of the A and E IR bands in $(\text{dien})\text{Mo}(\text{CO})_3$ compounds since the former and latter are related by $2k_c$ and $3k_c$, respectively. See ref 9. A plausible rationale for this extremely large coupling interaction is that in the $\text{Mo}(\text{OBU}^i)_2(\text{CO})_2(\text{py})_2$ molecule, the angle between the carbonyl ligands is extremely small, 72°.
- (11) F. A. Cotton and R. M. Wing, *Inorg. Chem.*, **4**, 314 (1965).
- (12) Attempts to grow crystals suitable for detailed single X-ray crystallography are in progress.
- (13) If the two μ -OR bridges are split symmetrically, the formal oxidation number of the carbonyl containing molybdenum atom is +3. However, one resonance form for the molecule could be written as $\text{Mo}^{8+}(\mu\text{-OR})_2\text{Mo}^{2+}$.
- (14) A more detailed argument based on the specific geometry observed for $\text{Mo}(\text{OBU}^i)_2(\text{py})_2(\text{CO})_2$ will be presented in a full paper.
- (15) M. H. Chisholm, F. A. Cotton, M. W. Extine, and R. L. Kelly, *Inorg. Chem.*, **18**, 116 (1979).
- (16) The complete structural report, MSC Report 7950 is available upon request in microfiche from only the Indiana University Chemistry Library.

Malcolm H. Chisholm,* John C. Huffman, Raymond L. Kelly
Department of Chemistry and Molecular Structure Center
Indiana University, Bloomington, Indiana 47405

Received August 13, 1979


Independent Existence in the Gas Phase of the 1-Methoxycyclopropyl and 2-Methoxyallyl Cations, as Observed in an Ion Cyclotron Resonance Spectrometer¹

Sir:


Substitution reactions of cyclopropyl compounds which have a good leaving group X at C₁ occur in the condensed phase, often with a disrotatory cleavage of the C₂-C₃ bond.² However, this ring rupture can be suppressed or even prevented if an electron-donating group Y is present at C₁ since it can supply electron density to the cyclopropyl cation center at C₁ as the group X leaves. If this substituent effect predominates over the donation of electrons from the C₂-C₃ bond, then substitution of group X without ring cleavage can take place. This electronic stabilization of the intermediate cyclopropyl cation has been found for various electron-donating groups Y such as dialkylamino,^{3,4} phenylthio,^{5,6} *o*-hydroxyphenyl,⁷ cyclopropyl,^{8,9} vinyl,¹⁰ methylthio,¹¹ *p*-methoxyphenyl,¹² *p*-tolyl,¹³ methoxy,¹³ arylethynyl,^{14,15} and several aryl groups.^{16,17} For Y = N(CH₃)₂^{3,4} and NHCH₃¹⁸ the intermediate cyclopropyl cations have been observed in NMR.

These observations are in line with ab initio molecular orbital calculations, which have suggested that the stabilization of the cyclopropyl cation by amino (and also hydroxy) substituents is so great that these species are not only more stable than the substituted perpendicular allyl cations but should be significantly more stable than the corresponding planar allyl cations.¹⁹


Evidence is now presented to show that 1-methoxycyclopropyl cations **1** are stable species in the gas phase and can be distinguished from the isomeric 2-methoxyallyl cations **2**. These ions have been generated independently in an ion cyclotron resonance drift cell²⁰ by electron impact ionization of 1-bromo-1-methoxycyclopropane²¹ and 2-methoxyallyl bromide,²² respectively, as shown in reactions 1 and 2. In the ions



1 the charge can be spread over the oxygen and adjacent cyclopropyl carbon atom, whereas in ions of structure **2** the charge is delocalized over the terminal carbon atoms. Ions **1** may be regarded as methyl cation bonded cyclopropanone species and, in contradistinction to ions **2**, they are therefore expected to be able to act as methyl cation donors to n-donor bases which have higher methyl cation affinities²³ (MCA) than cyclopropanone. Although the available MCA data are limited, amines are known to have higher MCA's than ketones: for example MCA(NH₃), 106; MCA(CH₃NH₂), 117, and MCA(CH₃O), 74 kcal/mol.²⁴ Possible reactions of ions **1** and **2** with ammonia, methylamine, dimethylamine, and trimethylamine have therefore been investigated by double resonance²⁰ experiments. They have all shown that ions **1**, but not ions **2**, do indeed transfer a methyl group as cation to these bases as summarized in eq 3. Variation of the electron energy used in the generation of the ions **1** and **2** over the range from 12 to 50 eV does not affect this observation. Furthermore, replacement of the methyl group in the ions **1** by a trideuteriomethyl group^{25a} confirms the transfer of the original methyl group in reaction 3.



The double resonance experiments have also shown that ions **1**, but not ions **2**, are able to transfer a proton to ammonia, methylamine, dimethylamine, and trimethylamine. Deuterium-labeling experiments show that it is one of the ring methylene protons which is transferred in this process, as indicated in reaction 4. Substitution of one of the methylene



groups in the ions **1** by a dideuteriomethylene group^{25b} reveals the existence of a kinetic isotope effect in reaction 4; using dimethylamine as the base this isotope effect is 1.7 at an electron energy of 12 eV decreasing to 1.6 at 14 eV and 1.3 at 16 eV.²⁶

The ions **1** and **2** are studied under ICR conditions on a time scale of $\sim 10^{-3}$ s, but they might have differing internal energies, which could also lead to different reactivities. However, we feel confident that the experimental results obtained strongly support the view that at least part of the ions **1** have a structure different from that of ions **2**. This is reinforced by the observation of a composite metastable peak for the loss of the bromine atom from the molecular ions of 1-bromo-1-methoxycyclopropane, whereas a noncomposite metastable peak is observed for loss of the bromine atom from the molecular ions of 2-methoxyallyl bromide.²⁷ This observation of the composite metastable peak indicates²⁸ that the molecular ions of 1-bromo-1-methoxycyclopropane which decompose within 10^{-6} s do so to form two different $(\text{M} - \text{Br})^+$ ions, presumably with and without rupture of the ring to give ions **2** and **1**, respectively.

Acknowledgment. The authors thank the Netherlands Organisation for Pure Research (SON/ZWO) for the purchase of the basic ICR mass spectrometer.

References and Notes

- (1) Dedicated to Professor E. Havinga of the University of Leiden on the occasion of his 70th birthday.
- (2) C. H. DePuy, *Acc. Chem. Res.*, **1**, 33 (1968).
- (3) E. Jongejan, W. J. M. van Tilborg, Ch. H. V. Dusséau, H. Steinberg, and Th. J. de Boer, *Tetrahedron Lett.*, 2359 (1972).
- (4) E. Jongejan, H. Steinberg, and Th. J. de Boer, *Tetrahedron Lett.*, 397 (1976).
- (5) U. Schöllkopf, E. Ruban, P. Tonne, and K. Riedel, *Tetrahedron Lett.*, 5077 (1970).
- (6) W. J. M. van Tilborg, H. Steinberg, and Th. J. de Boer, *Recl. Trav. Chim. Pays-Bas*, **93**, 303 (1974).
- (7) W. J. M. van Tilborg, J. R. van der Vecht, H. Steinberg, and Th. J. de Boer, *Tetrahedron Lett.*, 1681 (1972).
- (8) R. A. Martin and J. A. Landgrebe, *J. Org. Chem.*, **37**, 1996 (1972).
- (9) B. A. Howell and J. G. Jewitt, *J. Am. Chem. Soc.*, **93**, 798 (1971).
- (10) M. L. Poutsma and P. A. Ibarria, *Tetrahedron Lett.*, 4967 (1970).
- (11) M. Braun and D. Seebach, *Chem. Ber.*, **109**, 669 (1976).
- (12) J. R. van der Vecht, R. J. Dirks, H. Steinberg, and Th. J. de Boer, *Recl. Trav. Chim. Pays-Bas*, **96**, 309 (1977).
- (13) J. R. van der Vecht, H. Steinberg, and Th. J. de Boer, *Recl. Trav. Chim. Pays-Bas*, **96**, 313 (1977).
- (14) J. Salaün, *J. Org. Chem.*, **41**, 1237 (1976).
- (15) J. Salaün, *J. Org. Chem.*, **42**, 28 (1977).
- (16) H. C. Brown, C. G. Rao, and M. Ravindranathan, *J. Am. Chem. Soc.*, **99**,

- 7663 (1977).
- (17) H. C. Brown, C. G. Rao, and M. Ravindranathan, *J. Am. Chem. Soc.*, **100**, 7946 (1978).
- (18) E. Jongejan, H. Steinberg, and Th. J. de Boer, *Recl. Trav. Chim. Pays-Bas*, **98**, 66 (1979).
- (19) L. Radom, J. A. Pople, and P. v. R. Schleyer, *J. Am. Chem. Soc.*, **95**, 8193 (1973).
- (20) T. A. Lehman and M. M. Bursey, "Ion Cyclotron Resonance Spectrometry", Wiley-Interscience, New York, 1976.
- (21) This compound was obtained from the reaction of the trimethylsilyl ether of 1-methoxycyclopropanol with phosphorus tribromide in pyridine at -30°C . The 1-methoxycyclopropanol was obtained from the addition of methanol to cyclopropanone, synthesized from ketene (which was generated from acetone) and diazomethane according to the method of S. E. Schaafsma, H. Steinberg, and Th. J. de Boer, *Recl. Trav. Chim. Pays-Bas*, **85**, 1170 (1966); **86**, 651 (1967).
- (22) Synthesized according to the method of G. Greenwood and H. M. R. Hoffman, *J. Org. Chem.*, **37**, 611 (1972).
- (23) This is the negative of the enthalpy change for the gas phase reaction $\text{M} + \text{CH}_3^+ \rightarrow \text{MCH}_3^+$.
- (24) J. L. Beauchamp in "Interactions between Ions and Molecules", P. Ausloos, Ed., Plenum Press, New York, 1975, p 417.
- (25) (a) Trideuteriomethanol and (b) hexadeuterioacetone were used in the synthesis described in ref 20.
- (26) A referee has suggested that ions with the structure $\text{CH}_3\text{CH}=\text{C}=\text{OCH}_3^+$ might also transfer a methyl cation or a proton to the bases studied. However, the formation of such ions from ions **1** requires cleavage of the $\text{C}_1\text{-C}_2$ or $\text{C}_1\text{-C}_3$ bond of the cyclopropane ring. This would only be possible after the migration of a hydrogen atom from C_2 or C_3 to C_1 . The heat of formation of the intermediate 2-methoxycyclopropyl cations is expected to be 30–50 kcal/mol higher than that of ions **1**; see R. D. Bowen, J. R. Kalman, and D. H. Williams, *J. Am. Chem. Soc.*, **99**, 5481 (1977); R. D. Bowen, D. H. Williams, G. Hvistendahl, and J. R. Kalman, *Org. Mass Spectrom.*, **13**, 721 (1978). This value is similar to or higher than the activation energy of ~ 32 kcal/mol, which we have measured for the loss of formaldehyde from ions **1**. Therefore, it is very unlikely that those ions **1** which do not lose formaldehyde have rearranged to structure $\text{CH}_3\text{CH}=\text{C}=\text{OCH}_3^+$.
- (27) Decompositions occurring in the region between the ion source and the electric sector of an AEI MS 902 double focussing mass spectrometer with a β slit width of 0.5 mm.
- (28) R. G. Cooks, J. H. Beynon, R. M. Caprioli, and G. R. Lester, "Metastable Ions", Elsevier, Amsterdam, 1973.


M. W. E. M. van Tilborg, R. van Doorn
N. M. M. Nibbering*

Laboratory of Organic Chemistry, University of Amsterdam
Nieuwe Achtergracht 129
1018 WS Amsterdam, The Netherlands
Received May 10, 1979

Synthesis and Structure of $(\eta\text{-C}_5\text{H}_5)\text{W}(\text{PPh}_3)(\text{CO})(\text{CSPH})$: Transformation of a Thiocarbonyl Ligand into a Carbyne Ligand

Sir:

It was previously established in this laboratory¹ that the thiocarbonyl ligand in certain thiocarbonyl complexes can be alkylated with MeOSO_2F or R_3O^+ to give the CSR ligand, e.g.,



For this reaction to occur, it is necessary for the starting complex to be sufficiently electron rich to exhibit a $\nu(\text{CS})$ value of $\sim 1200\text{ cm}^{-1}$ or below. We now report a new and unexpected route to CSR complexes together with the first X-ray diffraction study of this type of ligand.

The title compound, $(\eta\text{-C}_5\text{H}_5)\text{W}(\text{PPh}_3)(\text{CO})(\text{CSPH})$, **A**, was prepared by the following sequence of reactions:

