

## Stereoselective Total Synthesis of ( $\pm$ )-Ishwarone


By EDWARD PIERS\* and TSE-WAI HALL

(Department of Chemistry, University of British Columbia, 2075 Wesbrook Place, Vancouver, B.C., Canada V6T 1W5)

**Summary** An efficient, stereoselective total synthesis of the racemic modification of the tetracyclic sesquiterpenoid ishwarone (**1**) is described.

THE structurally novel ketone ishwarone (**1**),<sup>1</sup> along with the corresponding hydrocarbon ishwarane (**2**), represent two of a relatively small number of tetracyclic sesquiterpenoids.

Although racemic ishwarane (**2**) has been obtained by total synthesis,<sup>3</sup> the preparation of ishwarone (**1**) has not yet been reported. We describe herein an efficient and completely stereoselective total synthesis of ( $\pm$ )-ishwarone (**1**),



Conversion<sup>4</sup> of the known aldehyde (**3**)<sup>5</sup> into the dibromoalkene (**4**),<sup>†</sup> followed by treatment of the latter with 2 equiv. of *n*-butyl-lithium in tetrahydrofuran (THF) at  $-78^{\circ}\text{C}^4$  and trapping<sup>6</sup> of the resultant lithium acetylide with formaldehyde, afforded (55% overall yield) the prop-2-ynylic alcohol (**5**). Hydrolysis (hydrochloric acid, acetone) of the latter gave the ketone (**6**) (90%) (m.p.  $43\text{--}45^{\circ}\text{C}$ ) which was hydrogenated (Pd/BaSO<sub>4</sub>, ethanol containing purified quinoline) to afford (**7**) (95%). Mesylation<sup>7</sup> of (**7**), followed by intramolecular alkylation (Bu<sup>t</sup>OK, Bu<sup>t</sup>OH) of the

resultant keto-mesylate (**8**), resulted in efficient formation ( $> 70\%$ ) of the octalone (**9**). Since both (**9**) and the corresponding decalone [obtained by hydrogenation of (**9**)] were recovered unchanged from methanolic sodium methoxide, it was clear that these compounds contained a *trans*-fused ring system.

The olefinic acetal (**10**) (m.p.  $100.5\text{--}102.5^{\circ}\text{C}$ ), obtained from (**9**) by a standard procedure, reacted smoothly with dimethyl diazomalonate<sup>8</sup> in the presence of copper bronze to afford a single adduct (**11**) (80%) (m.p.  $132\text{--}133.5^{\circ}\text{C}$ ). The stereochemistry of the latter was assigned as shown, since, for steric reasons, cyclopropane ring formation should have occurred *trans* to the angular methyl group. Conversion of the diester (**11**) into the keto-dimesylate (**14**), *via* compounds (**12**) (m.p.  $168\text{--}170^{\circ}\text{C}$ ) and (**13**) (m.p.  $138\text{--}139.5^{\circ}\text{C}$ ), was effected smoothly (67% overall yield) by standard methods (LiAlH<sub>4</sub> reduction, hydrolysis with hydrochloric acid in acetone, mesylation<sup>7</sup>).

Attempted intramolecular alkylation of (**14**) under a variety of conditions failed to produce the desired product (**16**). Although it is highly probable that the desired transformation occurred, the initially formed product (**16**) (cyclopropylcarbonyl mesylate) was apparently not sufficiently stable under the reaction conditions to allow for isolation. This difficulty was avoided as follows. The dimesylate (**14**) was smoothly converted (LiCl, hexamethylphosphoric triamide-ether, room temp.,  $> 90\%$  yield) into the corresponding dichloride (**15**), m.p.  $74\text{--}76^{\circ}\text{C}$ . When the latter was treated with freshly prepared potassium *t*-butoxide in dry THF, the isolable keto chloride (**17**) was obtained in 90% yield. Reduction of (**17**) with lithium triethylborohydride<sup>9</sup> in THF, followed by oxidation (pyridinium chlorochromate<sup>10</sup> in CH<sub>2</sub>Cl<sub>2</sub>) of the resultant product (**18**), afforded (75% overall yield) ( $\pm$ )-ishwarone (**1**), m.p.  $80\text{--}81^{\circ}\text{C}$ . The latter exhibited i.r. and <sup>1</sup>H n.m.r. spectra identical with those reported.<sup>1</sup> Furthermore, Wolff-Kishner reduction<sup>2</sup> of synthetic ( $\pm$ )-ishwarone (**1**) gave ( $\pm$ )-ishwarane (**2**), identical with a sample of the same material previously prepared by Kelly and his co-workers.<sup>3</sup>

We are very grateful to Professor R. B. Kelly for a sample of ( $\pm$ )-ishwarane. Financial support from the National Research Council of Canada and a N.R.C.C. scholarship (to T. W. H.) are gratefully acknowledged.

(Received, 28th July 1977; Com. 778.)

<sup>†</sup> All new compounds exhibited the expected spectral properties and gave satisfactory elemental analyses and/or molecular weight determinations (high-resolution mass spectrometry).

<sup>1</sup> A. K. Ganguly, K. W. Gopinath, T. R. Govindachari, K. Nagarajan, B. R. Pai, and P. C. Parthasarathy, *Tetrahedron Letters*, 1969, 133; T. R. Govindachari, K. Nagarajan, and P. C. Parthasarathy, *Chem. Comm.*, 1969, 823; H. Fuhrer, A. K. Ganguly, K. W. Gopinath, T. R. Govindachari, K. Nagarajan, B. R. Pai, and P. C. Parthasarathy, *Tetrahedron*, 1970, **26**, 2371.

<sup>2</sup> T. R. Govindachari, P. A. Mohamed, and P. C. Parthasarathy, *Tetrahedron*, 1970, **26**, 615.

<sup>3</sup> R. B. Kelly, J. Zamecnik, and B. A. Beckett, *Chem. Comm.*, 1971, 479; *Canad. J. Chem.*, 1972, **50**, 3455; R. B. Kelly and S. J. Alward, *ibid.*, 1977, **55**, 1786; R. M. Cory and F. R. McLaren, *J. C. S. Chem. Comm.*, 1977, 587.

<sup>4</sup> E. J. Corey and P. L. Fuchs, *Tetrahedron Letters*, 1972, 3769.

<sup>5</sup> F. E. Ziegler and P. A. Wender, *Tetrahedron Letters*, 1974, 449; F. E. Ziegler, G. R. Reid, W. L. Studt, and P. A. Wender, *J. Org. Chem.*, 1977, **42**, 1991.

<sup>6</sup> S. S. Klioze and F. P. Darmory, *J. Org. Chem.*, 1975, **40**, 1588.

<sup>7</sup> R. K. Crossland and K. L. Servis, *J. Org. Chem.*, 1970, **35**, 3195.

<sup>8</sup> B. W. Peace and D. S. Wulfsberg, *Synthesis*, 1973, 137; R. K. Singh and S. Danishefsky, *J. Org. Chem.*, 1976, **41**, 1668.

<sup>9</sup> H. C. Brown and S. Krishnamurthy, *J. Amer. Chem. Soc.*, 1973, **95**, 1669.

<sup>10</sup> E. J. Corey and J. W. Suggs, *Tetrahedron Letters*, 1975, 2647.