

Solvent Dependency of the Steric Course of Dienolate Alkylation

By Y. NAKADAIIRA and J. HAYASHI

(*Department of Chemistry, Tohoku University, Sendai, 980, Japan*)







and H. SATO and K. NAKANISHI*

(*Department of Chemistry, Columbia University, New York, New York 10027*)

Summary The stereochemistry of methylation of α -methyl- $\alpha\beta$ -unsaturated enones is shown to be dependent on reaction media.

WE present evidence that alkylation of dienolate anions derived from enones (I) and (II) are markedly affected by the solvent system, and that this factor should be considered seriously in alkylation reactions. The enolate anion

of 17 β -acetoxy-4-methyl-19-testosterone (I)¹ formed with EtC(Me)₂OK-C₆H₆ was treated with MeI to give, after acetylation, the 4,4-dimethyl steroid (II)² (52% overall yield); c.d. (EtOH) $\Delta\epsilon_{295} + 2.58$; u.v. (EtOH) ϵ_{295} 90. The positive Cotton effect together with enhanced u.v. maximum (n, π^*) establishes a chair conformation for ring A.³ N.m.r. signals of the 4,4-dimethyl group appear in CDCl₃ at 1.20 and 1.25 p.p.m. and are not sufficiently separated for



intramolecular nuclear Overhauser effect (N.O.E.) studies.⁴ However, in C₆D₆ they are shifted to 1.12 and 1.32 p.p.m.; irradiation of the 1.32 p.p.m. signal caused a 10% increase in the integrated area of the olefinic 6-H signal at 5.51 p.p.m. No increment was observed upon irradiation of the 1.12 p.p.m. signal. The 1.32 and 1.12 p.p.m. signals can therefore be assigned unambiguously to the 4 α - and 4 β -methyl groups (see II'), respectively.

In order to study the steric factors involved in this reaction, the same dienolate alkylation was next carried out with CD₃I, which gave a mixture of (II α) and (II β).† The intensity ratio of n.m.r. signals at 1.12 and 1.32 p.p.m. in C₆D₆ (1.25 and 1.20 p.p.m. in CDCl₃, respectively) (see II') indicated that 4 α -methylation giving (II α) and 4 β -methylation giving (II β) had occurred in a ratio of 3:7. Namely, a β -attack was favoured in benzene.

The conversion of (I) into (II) [and also to (II α)/(II β)] had been carried out previously by treatment with Bu^tOK-Bu^tOH and then with MeI.⁵ On the basis of n.m.r. intensity measurements of the deuteriated species (II α)/(II β), it was concluded that this ratio was 1:10. However, assignments of the two methyl groups at 73 and 75 Hz (from Me₄Si, 60 MHz) were opposite‡ to those described above as derived from N.O.E. measurements. Accordingly, the previous conclusions⁵ also have to be reversed, *i.e.*, in Bu^tOH an α -attack is preferred over β -attack to the extent of 10:1. This is contrary to the present results carried out in benzene, and shows that the solvent§ has a profound effect on the stereochemical course of enolate alkylations.

Similar alkylation of (\pm)-decalone (III) with EtC-(Me)₂OK-C₆H₅ and MeI gave the 1,1-dimethyl product (IV) in 55% yield; u.v. (EtOH) ϵ_{295} 100. Again, the 1.28 p.p.m. peak (in C₆D₆) exhibited a 15% N.O.E. on the 5.50 p.p.m. 9-H signal and thus could be assigned to the 1 α -methyl group; the 1 β -methyl appeared at 1.08 p.p.m. Alkylation with CD₃I yielded a mixture of (IV α) and (IV β) in a ratio of 3:7 (1 α - and 1 β -methyl groups appeared at 1.21 and 1.26 p.p.m., respectively, in CDCl₃), which indicated that, as in the case of (I) \rightarrow (II), a β -attack was predominant.

The effect of a 5-methyl substituent on the direction of alkylation in benzene was investigated next with the angularly substituted decalone (V). Dienolate alkylation under the present conditions gave (VI) in 80% yield; u.v. (EtOH) ϵ_{290} 33; n.m.r. (CDCl₃) 1.22, 1.22, 0.99 (three methyls); (C₆D₆) 1.27, 1.27, 0.91 p.p.m. Due to overlap of the n.m.r. methyl peaks in enone (VI), this was converted into the known saturated ketone (VII)⁷ (and <10% of *cis*-isomer, as judged from n.m.r.) in practically quantitative yield, *via* (i) reduction with LiAlH₄, (ii) hydrogenation over Pd-C-EtOH, and (iii) Jones oxidation; the carbonyl-reduction step was carried out in order to ensure stereo-specific hydrogenation.¶ This saturated ketone (VII) had been obtained through reductive methylation of (V) by Stork, *et al.*,⁸ and the stereochemistry of the alkylating step has recently been clarified by Mathews, *et al.*^{7,8} As the n.m.r. methyl peaks (in CDCl₃) had already been assigned as depicted in (VII),⁷ it was possible to clarify the

† The physical constants of (II) and (II α)/(II β) were identical except for the appearance in the i.r. of the latter of C-D stretching bands at 2250, 2510, and 2070 cm⁻¹.

‡ The critical assignments of the peaks were erroneously based on the assumption that the chemical shift of a methyl in a 4,4-gem-dimethyl group would be the same as that of a methyl in a 4-methyl-4-ethyl group.

§ We have presumed that difference in solvent polarity exerts a larger effect than the reagent base.

¶ Direct hydrogenation of (VI) resulted in a *ca.* 1:1 mixture of *cis* and *trans* saturated ketones, which could not be separated under various t.l.c. conditions. However, in the three-step reaction, the *trans* isomer was formed predominantly (>90%).

steric preference of the $\text{EtC}(\text{Me})_2\text{OK}-\text{C}_6\text{H}_6-\text{MeI}$ alkylation reactions. Thus it was found that decalone (V) gave compounds (VII α) and (VII β) in a 9:1 ratio with $\text{EtC}(\text{Me})_2\text{OK}-\text{C}_6\text{H}_6-\text{CD}_3\text{I}$ which meant that the major course was now an α -attack as expected.⁹

The results of alkylation carried out in the present non-polar benzene system, especially those of (I) and (III), can be rationalized on the basis of a product-like transition

state (sp^3 configuration at alkylating site) rather than a dienolate-like transition state (sp^2). Specific solvation by Bu^tOH on the less hindered side (β) of the anion resulting in α attack of methyl is another possible explanation.

The work was supported by a P.H.S. grant.

(Received, 23rd December 1971; Com. 2179.)

¹ N. Atwater, *J. Amer. Chem. Soc.*, 1960, **83**, 2847.

² Cf. S. L. Mukherjee and P. C. Dutta, *J. Chem. Soc.*, 1960, 67.

³ M. Gorodetsky, A. Yagev, and Y. Mazur, *J. Org. Chem.*, 1966, **31**, 699.

⁴ F. A. L. Anet and A. J. R. Bourn, *J. Amer. Chem. Soc.*, 1965, **87**, 5250; M. C. Woods, I. Miura, Y. Nakadaira, A. Terahara, M. Maruyama, and K. Nakanishi, *Tetrahedron Letters*, 1967, 321.

⁵ V. Permutti and Y. Mazur, *J. Org. Chem.*, 1966, **31**, 705.

⁶ G. Stork, P. Rosen, N. Goldman, R. V. Coomb, and J. Tsuji, *J. Amer. Chem. Soc.*, 1965, **87**, 275; see following for applications to stereospecific total synthesis of lupeol: G. Stork, S. Uyeo, T. Wakamatsu, P. Grieco, and J. Labovitz, *ibid.*, 1971, **93**, 4945.

⁷ R. S. Matthews, P. D. Hyer, and E. A. Folkers, *Chem. Comm.*, 1970, 38.

⁸ R. S. Matthews, S. J. Girgenti, and E. A. Folkers, *Chem. Comm.*, 1970, 708.

⁹ G. Just and K. St. C. Richardson, *Canad. J. Chem.*, 1964, **42**, 464.