

PHOTOCHEMICAL REACTION OF N-ACYLSYDNONE IMINES


Akiko CHINONE and Masaki OHTA

Department of Chemistry, Faculty of Science, Tokyo Institute  
of Technology, Ookayama, Meguro-ku, Tokyo


Irradiation of a benzene-ethanol solution of N-acyl-3,4-di-phenylsydnone imine afforded 1,5-diphenyl-4-hydroxy-1,2,3-triazole and anhydro-3-acyl-1,5-diphenyl-4-hydroxy-1,2,3-triazolium hydroxide. Irradiation of N-acylsydnone imine in the presence of dimethyl acetylenedicarboxylate yielded an addition product.

As part of our continuing interest in photochemical reaction of mesoionic compounds, we reported<sup>1,2)</sup> the photolysis of sydnone. In this paper, we describe the photochemical reaction of N-acylsydnone imines.


Irradiation of a benzene-ethanol (14:1) solution of N-acetyl-3,4-diphenylsydnone imine (I) in a Pyrex vessel under nitrogen and with a high pressure mercury lamp (100W, 9 hr) yielded 1,5-diphenyl-4-hydroxy-1,2,3-triazole (II) (yield 10%; mp 211-213°C; NMR  $\tau$ (DMSO- $d_6$ ) 2.4-2.75 (m, 10H), -0.67 (s, 1H)) and anhydro-3-acetyl-1,5-diphenyl-4-hydroxy-1,2,3-triazolium hydroxide (III) (yield 11%; mp 150-151°C; Found: C, 68.62; H, 4.74; N, 15.24%. Calcd for  $C_{16}H_{13}N_3O_2$ : C, 68.80; H, .69; N, 15.05%; UV  $\lambda_{max}$ (EtOH) 247 m $\mu$  ( $\epsilon$  11300); IR  $\nu$ (KBr) 3050, 1790, 1770, 1610, 1370, 1190, 770, and 700  $cm^{-1}$ ; NMR  $\tau$ (CDCl<sub>3</sub>) 7.7 (s, 3H), 2.6-2.75 (m, 10H); mass spectrum  $M^+$  m/e 279 (4%), m/e 43 (54%,  $CH_3CO$ ), m/e 77 (79%,  $C_6H_5$ ), m/e 180 (base peak,  $C_6H_5C=NC_6H_5$ ), m/e 236 (78%,  $M^+-CH_3CO$ )). Structure of II was confirmed by mixed melting point measurement with an authentic sample which was prepared by treating I with methanolic potassium hydroxide.<sup>3)</sup> Treatment of III with methanolic potassium hydroxide afforded II at room temperature. Irradiation of N-benzoyl-3,4-diphenylsydnone imine (IV) under the same conditions afforded II (yield 35%) and anhydro-3-benzoyl-1,5-diphenyl-4-hydroxy-1,2,3-triazolium hydroxide (V) (yield 2%; mp 179-180°C; Found: C, 73.81; H, 4.35; N, 12.33%. Calcd for  $C_{21}H_{15}N_3O_2$ : C, 73.89; H, 4.43; N, 12.31%; UV  $\lambda_{max}$ (EtOH) 235 m $\mu$  ( $\epsilon$  28400); IR  $\nu$ (KBr) 1760, 1610, 1370, 1250, 1235, 1055, 760, and 700  $cm^{-1}$ ; Mass spectrum of IV did not show a molecular ion peak but strong peaks at m/e 105 (base peak,  $C_6H_5CO$ ) and at m/e 180 (85%,  $C_6H_5C=NC_6H_5$ )).



These results showed that in the photochemical reaction of N-acylsydnone imines rearrangement preferentially occurred to the nitrile imine formation which predominated in the photolysis of sydnones,<sup>1,2,4)</sup> as shown below.



1,3-Dipolar cycloaddition of sydnones under irradiation has been reported.<sup>1,2,4)</sup> The photochemical reaction of sydnone in the presence of dimethyl acetylenedicarboxylate (VI) afforded the addition products of nitrile imines derived from sydnones and VI. However, irradiation of I in the presence of VI yielded II, III, and a crystalline adduct,  $C_{20}H_{17}N_3O_5$  (II:VI=1:1) (yield 3.6%; mp 110-111°C; Found: C, 63.30; H, 4.55; N, 11.37%. Calcd for  $C_{20}H_{17}N_3O_5$ : C, 63.32; H, 4.52; N, 11.08%; IR  $\nu$ (KBr) 3425, 1730, 1665, 1595, 1280, 755, and 690  $cm^{-1}$ ). The structure of the adduct was supposed to be dimethyl 3-hydroxy-4,7-diphenyl-1,2,7-triazabicyclo[2,2,1]heptane-5,6-dicarboxylate (VII), though not conclusive. Irradiation of IV under the same conditions afforded II and V, and the adduct was not detected. Treatment of II with VI did not afford the adduct in the dark.



VII

- 1) Y. Huseya, A. Chinone, and M. Ohta, Bull. Chem. Soc. Japan, **44**, 1667 (1971).
- 2) Y. Huseya, A. Chinone, and M. Ohta, *ibid.*, **45**(11) (1972).
- 3) H. U. Daeniker and J. Druey, Helv. Chim. Acta, **45**, 2441 (1962).
- 4) G. S. Angadiyavar and M. V. George, J. Org. Chem., **36**, 1589 (1971).

( Received July 21, 1972 )