

The authors are greatly indebted to Professor I. Uritani, Department of Agriculture, Nagoya University, for originally suggesting this work and his continued interest, and to Dr. J. Corse, Western Research Laboratory, Albany, California, for sending us copies of their two manuscripts pertaining to "isochlorogenic acid c" prior to publication. This work has been supported by a Plant Phenolics Grant-in-Aid from the Ministry of Education.

Department of Chemistry,
Tohoku University,
Sendai, Japan

Yasuo Inoue (井上康男)
Shohei Aoyagi (青柳象平)
Koji Nakanishi (中西香爾)

Received November 2, 1964


[Chem. Pharm. Bull.]
13(1) 104~105 (1965)

UDC 547.91.05 : 582.38

21-Episerratenediol, Isolation and its Structure

This communication is concerned with the constitution and stereochemistry of a triterpene-diol, 21-episerratenediol, which we have isolated from *Lycopodium serratum* THUNB. var. *Thunbergii* MAKINO (Lycopodiaceae) along with serratenediol, serratenediol-3-acetate and two new triterpenoids.

The structure of serratenediol and its 3-acetate, two major triterpenoid constituents, were already elucidated as formulated by I and II, respectively.^{1,2)}



Separation of minor triterpenoids was conveniently effected by repeated chromatography of derived acetates over silica gel and alumina. The physical constants of the three new triterpenoids and their acetates are listed in the Table. The nuclear magnetic resonance spectrum*¹ of A-diacetate indicated the presence of a trisubstituted

	Alcohol		Acetate
(A) 21-Episerratenediol	C ₃₀ H ₅₂ O ₂ ,	m.p. 289~290°	C ₃₄ H ₅₆ O ₄ , m.p. 225~229° [α] _D -29° (c=2.06, CHCl ₃)
(B) Serratriol	C ₃₀ H ₅₀ O ₃ ,	m.p. 335~336°	C ₃₈ H ₅₆ O ₆ , m.p. 245~247° [α] _D +21° (c=1.9, CHCl ₃)
(C) Tohogenol	C ₃₀ H ₅₂ O ₃ · 1/2 H ₂ O,	m.p. 242~244°	C ₃₄ H ₅₆ O ₅ , m.p. 305~306° [α] _D +28° (c=1.6, CHCl ₃)

*¹ NMR spectra were taken on Varian A-60 machine in CDCl₃ with Me₄Si as an internal standard by Mr. T. Shingu, Kyoto University, to whom we are indebted.

1) Y. Inubushi, T. Sano, Y. Tsuda : Tetrahedron Letters, No. 21, 1303 (1964).

2) Y. Tsuda, T. Sano, K. Kawaguchi, Y. Inubushi : Ibid., No. 20, 1279 (1964).

double bond at 4.65τ (1H, broad). The C-methyl region of the spectrum was almost superimposable with that of serratenediol diacetate but the acetate methyls of the A-di-acetate were appeared separately at 7.91τ (3H) and 7.95τ (3H) in contrast to serratenediol diacetate in which the acetate methyls appeared as an overlapped single peak 7.97τ (6H). Moreover, a broad multiplet at 5.52τ (1H) and a sharp triplet at 5.31τ (1H), both are attributable to geminal protons to acetoxyl groups, suggest that the former corresponds to an axial proton, the latter corresponding to an equatorial one.³⁾

Chromium trioxide-pyridine complex oxidation of the diol (A) gave a diketone m.p. $206\sim 208^\circ$ which was identical with serratenedione in all respects. Hence the diol (A) must be an epimer of serratenediol in respect to the stereochemistry of either one or two hydroxyl groups. Then, the diol (A) was partially acetylated by heating the compound with acetic anhydride and pyridine for 8 minutes to a monoacetate, $C_{32}H_{52}O_3$, m.p. $317\sim 319^\circ$; $[\alpha]_D -44^\circ$ ($c=1.3$, $CHCl_3$), IR ν_{max}^{Nujol} cm^{-1} : 3584 (OH), 1730, 1239 (OAc), in about 10% yield, which was apparently different from serratenediol 3-acetate (II). In the nuclear magnetic resonance spectrum of this mono-acetate, a broad multiplet at 5.55τ (1H) suggested that the presence of an axial proton geminal to an acetoxyl group and a sharp triplet at 6.55τ (1H) showed the presence of an equatorial proton geminal to a hydroxyl group.

Oxidation of the mono-acetate with chromium trioxide-pyridine complex afforded a keto-acetate m.p. $306\sim 307^\circ$. The identity of this with serrateneolone acetate²⁾ (3 β -hydroxy-serrat-14-en-21-one acetate) was rigidly confirmed by direct comparisons of the both samples (m.p., mixed m.p., IR, ORD and TLC). Therefore, the diol (A) was established to be 21-epi-serratenediol (III) (serrat-14-ene-3 β ,21 β -diol).^{*2}

Faculty of Pharmaceutical Sciences,
Osaka University,
Toyonaka, Osaka

Yasuo Inubushi (犬伏康夫)
Yoshisuke Tsuda (津田喜典)
Takehiro Sano (佐野武弘)
Ritsuko Nakagawa (中川律子)

Received November 2, 1964

^{*2} According to the private communication, Dr. J. W. Rowe isolated episerratenediol (diacetate, m.p. $236\sim 237^\circ$, $[\alpha]_D -18^\circ$), but because of insufficient amounts available the stereochemistry of hydroxyl groups has not been established. The compound is probably identical with 21-episerratenediol, although the direct comparison was not available. cf. J. W. Rowe: Tetrahedron Letters, No. 34, 2347 (1964).

3) H. Vorbrüggen, S. C. Pakrashi, C. Djerassi: Ann., 668, 57 (1963).