

suffered practical difficulties with analogue filtration. The tailored detection method is a simpler and more effective way of regaining the lost sensitivity, since it may be set up to act as a matched digital filter for the designated frequency components. The requisite mask is made up to match selected lines in the noise decoupled spectrum and each incoming free induction decay  $E(t)$  is multiplied by the corresponding time domain reference  $A'(t)$ , and the integral  $\int E(t)A'(t)dt$  stored as one point on the interferogram  $S(t_1)$ . Fourier transformation of  $S(t_1)$  generates a  $J$ -scaled spectrum with optimum sensitivity. This is because a  $J$ -scaled spectrum is equivalent to a skew projection of a two-dimensional spectrum,<sup>10,11</sup> the process of tailored detection corresponding to projection of selected traces (those carrying the signals), intermediate noisy traces being suppressed.

Cholesteryl acetate provides an example of a reasonably complicated carbon-13 spectrum to illustrate the adaptation of tailored detection to the  $J$ -scaling technique. Spectra were obtained on a Varian CFT-20 spectrometer modified as described earlier.<sup>8</sup> Figure 1a shows the broad band decoupled spectrum of the aliphatic region, comprising 25 resonances, several of which fall in crowded groups of nearly degenerate frequencies.<sup>12</sup> Lines in the noise decoupled spectrum are selected for the masks  $A'(f)$  in such a way to avoid confusing overlap of multiplets in the  $J$ -scaled spectrum. For the case of cholesteryl acetate, four separate mask spectra were used, each with line broadening chosen to match the experimental line widths, and each concentrating as far as possible on well-separated chemical shifts. Careful choice of the scaling factor helps clarify the resulting  $J$ -scaled spectra; traces b and c of Figure 1 used a scaling factor of  $R = 6$  while d and e used  $R = 10$ . Several near degeneracies in chemical shifts make the assignment quite challenging. For example in Figure 1b, the experiment designed to pick out the response of site 22 (a triplet) cannot avoid some breakthrough of the sharp singlet resonance from site 10, and similarly the triplet from site 2 has a superimposed sharp doublet from site 25. However, even lines that are unresolved in the decoupled spectrum may be assigned if they have different multiplicities, for example the quartet from the acetate methyl group and the triplet from site 11 in Figure 1b.

Several types of NMR experiments involve monitoring the peak of a chosen resonance as a function of time; the examples include spin-lattice relaxation measurements, the study of transient intermediates or transient polarization effects, and the determination of chemical exchange rates by following magnetization transfer. By acting as a matched filter and calculating an integral, tailored detection gives a much better measure of intensity, less susceptible to noise and variations of instrument resolution; it can also operate on several lines at once. Applied to double Fourier transform experiments, this procedure should prove generally useful in optimizing the signal-to-noise ratio of projections and cross sections of two-dimensional spectra. As has been demonstrated, the method is also valuable in analogous one-dimensional experiments such as  $J$  scaling and  $J$  spectroscopy.<sup>13</sup>

**Acknowledgments.** This work was made possible by an equipment grant from the Science Research Council and was supported by a Research Studentship (G.A.M.).

## References and Notes

- (1) J. Jeener, Ampere International Summer School, Basko Polje, Yugoslavia, 1971; Second European Experimental NMR Conference, Enschede, Holland, 1975.
- (2) W. P. Aue, E. Bartholdi, and R. R. Ernst, *J. Chem. Phys.*, **64**, 2229 (1976).
- (3) G. Bodenhausen, R. Freeman, R. Niedermeyer, and D. L. Turner, *J. Magn. Reson.*, **26**, 133 (1977).
- (4) B. L. Tomlinson and H. D. W. Hill, *J. Chem. Phys.*, **59**, 1775 (1973).
- (5) R. Freeman, H. D. W. Hill, B. L. Tomlinson, and L. D. Hall, *J. Chem. Phys.*, **61**, 4466 (1974).

- (6) G. Bodenhausen, R. Freeman, and G. A. Morris, *J. Magn. Reson.*, **23**, 171 (1976); G. A. Morris and R. Freeman, *ibid.*, **29**, 433 (1978).
- (7) In practice this may be derived from another experimental spectrum by setting selected peaks to unit intensity and all other regions to zero intensity.
- (8) R. Freeman and G. A. Morris, *J. Magn. Reson.*, **29**, 173 (1978).
- (9) R. Freeman, J. B. Grutzner, G. A. Morris, and D. L. Turner, *J. Am. Chem. Soc.*, **100**, 5637 (1978).
- (10) W. P. Aue, J. Karhan, and R. R. Ernst, *J. Chem. Phys.*, **64**, 4226 (1976).
- (11) K. Nagayama, P. Bachman, K. Wüthrich, and R. R. Ernst, *J. Magn. Reson.*, **31**, 133 (1978).
- (12) The assignment is that of H. J. Reich, M. Jautelat, M. T. Messe, F. J. Weigert, and J. D. Roberts, *J. Am. Chem. Soc.*, **91**, 7445 (1969), except that C-12 and C-16 have been interchanged in accordance with the later work of J. W. ApSimon, H. Beierbeck, and J. K. Saunders, *Can. J. Chem.*, **51**, 3874 (1973), and of W. B. Smith, D. L. Deavenport, J. A. Swanzy, and G. A. Pate, *J. Magn. Reson.*, **12**, 15 (1973).
- (13) R. Freeman and H. D. W. Hill, *J. Chem. Phys.*, **54**, 301 (1971).

Gareth A. Morris, Ray Freeman\*


Physical Chemistry Laboratory  
South Parks Road, Oxford, England

Received June 5, 1978

## Angle and Torsional Strains as Tools for Enhancing the Kinetic Stability of $\beta$ -Hydroperoxy Ketones


Sir:

The alkaline peroxidation of enones, discovered by Weitz and Scheffer<sup>1</sup> and first kinetically scrutinized by Bunton and Minkoff,<sup>2</sup> has more recently been shown to be nonstereospecific but highly stereoselective in acyclic systems.<sup>3-7</sup> Irrespective of the geometry of the starting  $\alpha,\beta$ -unsaturated ketone, the product having the less hindered carbonyl group is kinetically favored, a phenomenon attributed to overlap control.<sup>7</sup> These observations have implicated  $C_\alpha$ - $C_\beta$  rotational equilibration during the lifetime of the enolate anion **1** prior to ejection of hydroxide ion. Evidence has been presented to in-




dicate not only that the rate-determining step is  $k_1$ , the attack of  $\text{HOO}^-$  on the  $\beta$  carbon,<sup>2</sup> but that reversal of hydroperoxide anion addition may compete with closure to the epoxy ketone ( $k_{-1} > k_2$ ).<sup>7</sup>

Greater appreciation of the above mechanism can be gained by considering several extreme possibilities. One of these is the condition where  $k_2 \gg k_P$  such that protonation of **1** by solvent is not encountered. The lone established example of this behavior has been reported by Pappas and Bao who observed that the epoxidation of phenalenone (**2a**) with alkaline deuterium peroxide afforded **4a** with no deuterium incorporation.<sup>8</sup> In their hands, the reverse experiment with **2b** and  $\text{HOO}^-$  gave only **4b**. Since the rate of proton transfer from water to an enolate anion has been estimated to be on the order of  $10^5 \text{ s}^{-1}$ ,<sup>8,9</sup> the rapid cyclization of **3** may be close to a limit for nucleophilic attack on a peroxide linkage.




We now describe novel experimental findings that are at the other extreme of the scale, that is where the barriers associated with  $k_{-1}$  and  $k_2$  have become so high that enolate **1** is relegated



exclusively to protonation by solvent. Alkali stable  $\beta$ -hydroperoxy ketones, a class of compounds which seemingly has escaped prior isolation,<sup>10-13</sup> are thereby produced.


The pivotal indenones **6** and **9b** were prepared by Diels-Alder cycloaddition of 3-bromoindenone<sup>14</sup> to 1,3-cyclohexadiene ( $\text{C}_6\text{H}_6$ ,  $\text{BF}_3\text{-Et}_2\text{O}$ , 5 °C, 2 h) and cyclopentadiene ( $\text{CH}_2\text{Cl}_2$ , 25 °C, 12 h), respectively, followed by dehydrobromination of the resulting **5** and **8b** with an amidine base. Bromo ketone **8a**, obtained either through sequential catalytic hydrogenation and benzylic bromination (NBS) of the simpler indenone adduct or preferably by reduction of **8b**, served as the starting material for **9a**. Whereas **6** and **9a** proved to be entirely stable, **9b** was found to be labile with respect to formation of a high melting (>225 °C) substance and was utilized immediately upon its generation.

The alkaline peroxidation of **6** at 25 °C led efficiently to hydroperoxide **7** (mp 123–124 °C; 69% after silica gel chromatography),<sup>15</sup> the stereochemical assignment of which is secured by its characteristic <sup>1</sup>H NMR spectrum and cyclization to **11** (mp 94–95 °C) when treated with 2,4,4,6-tetrabromocyclohexa-2,5-dienone.<sup>16</sup> The kinetic preference for endo




attack conforms to the lower heat of formation calculated for stereoisomer **7** relative to its exo counterpart.<sup>17</sup> Stannous chloride reduction of **7** afforded **12a** which proved spectroscopically different from **5**, but closely allied to **12b**, the product of direct methanol addition to **6**.

Entirely similar results were realized with **9a** and **9b** which furnished **10a** (mp 124–125.5 °C, 60%) and **10b** (unstable oil, 82%). These hydroperoxides were individually reduced to **14** and **15**, and shown in turn to be stereochemically related through catalytic hydrogenation.



While submission of **7** to more alkaline conditions (15% aqueous NaOH,  $\text{CH}_3\text{OH}$ , 25 °C) did result in epoxy ketone formation (55% of **13** after 5 h), **10a** and **10b** proved totally unreactive toward base. Their inability to cyclize is not due to the kinetic nonacidity of the  $\alpha$ -keto proton, for deuterium exchange experiments under less basic conditions ( $\text{Na}_2\text{CO}_3$ ,  $\text{CH}_3\text{OD}$ ,  $\text{D}_2\text{O}$ , 25 °C) led to isotopic exchange at both  $>\text{CHCO-}$  and  $-\text{OOH}$ , nor was reconversion of **10a** and **10b** to their precursor indenones observed in a variety of alkaline media.

As a test of the premise that angle and torsional strains actually control the kinetic properties of the intermediate enolate



anions, the susceptibility of naphthoquinones **16–18**<sup>18</sup> to alkaline peroxidation was also examined. Molecular models reveal that fusion of a six- rather than a five-membered Michael acceptor to a representative bicyclic framework results in a lessening of energetically unfavorable structural constraints during the incorporation of a three-membered epoxy ring across the central bond. In line with this a priori assessment, the epoxidations of **16–18** were found to proceed to completion during 45 min to 2 h at 25 °C (30%  $\text{H}_2\text{O}_2$ ,  $\text{Na}_2\text{CO}_3$ ,  $\text{C}_2\text{H}_5\text{OH}$ ); only in the case of **17** was a trace of residual hydroperoxy ketone observable (TLC analysis). Of some interest is the stereoselectivity of these reactions. Whereas the product ratio in the case of **16** was 14:1 in favor of the endo isomer (O syn to the isolated double bond), that in norbornadienyl example **17** was more equitable, although reversed (1 part endo:3 parts exo). Quinone **18** gave rise exclusively to the exo epoxide.<sup>19</sup>

Accordingly, the dominant forces underlying the unprecedented behavior of **6**, **9a**, and **9b** toward  $-\text{OOH}$  appear to be angle and torsional strain.<sup>20</sup> The incipient formation of a three-membered ring which incorporates the  $\text{C}_2$  and  $\text{C}_3$  atoms of a norbornyl system generates eclipsing interactions which are significantly increased through added fusion of a five-membered ring. In the present instance, prevailing kinetic and steric factors guide the initial stereoselectivities of nucleophilic hydroperoxide attack.<sup>21</sup> For indenones **9a** and **9b**, bonding occurs preferentially syn to the methylene bridge. The resulting enolate anions can neither reverse the Michael process nor cyclize to the epoxy ketone. For the enolate anion of **7**, the strain associated with ring closure is somewhat lessened owing to greater conformational flexibility and cyclization that can ultimately be achieved.

**Acknowledgment.** We are grateful for the partial financial support provided by the National Cancer Institute (Grant CA-12115).

## References and Notes

- (1) E. Weitz and A. Scheffer, *Chem. Ber.*, **54**, 2327 (1921).
- (2) C. A. Bunton and G. J. Minkoff, *J. Chem. Soc.*, 665 (1949).
- (3) W. G. Black and R. E. Lutz, *J. Am. Chem. Soc.*, **75**, 5990 (1953); C. R. Bauer and R. E. Lutz, *ibid.*, **75**, 5995 (1953); R. E. Lutz and J. O. Weiss, *ibid.*, **77**, 1814 (1955).
- (4) N. H. Cromwell and R. A. Setterquist, *J. Am. Chem. Soc.*, **76**, 5752 (1954).
- (5) H. H. Wasserman, N. E. Aubrey, and H. E. Zimmerman, *J. Am. Chem. Soc.*, **75**, 96 (1953); H. H. Wasserman and N. E. Aubrey, *ibid.*, **77**, 590 (1955).
- (6) H. O. House and D. J. Reif, *J. Am. Chem. Soc.*, **77**, 6525 (1955); H. O. House and R. S. Ro, *ibid.*, **80**, 2428 (1958).
- (7) H. E. Zimmerman, L. Singer, and B. S. Thyagarajan, *J. Am. Chem. Soc.*, **81**, 108 (1959).
- (8) S. P. Pappas and L. Q. Bao, *J. Am. Chem. Soc.*, **95**, 7906 (1973).
- (9) M. Eigen, *Angew. Chem., Int. Ed. Engl.*, **3**, 1 (1964).
- (10) For a brief tabular summary, see H. Kropf and M. Ball, *Tetrahedron*, **28**, 1391 (1972).
- (11) A cyclic peroxide by-product has been reported from the alkaline epoxidation of mesityl oxide: G. B. Payne, *J. Org. Chem.*, **23**, 310 (1958).
- (12) The base-promoted oxygenation of 2,5,6-tri-*tert*-butylcyclohex-2-ene-1,4-dione afforded a hydroperoxide which is  $\alpha$  to one carbonyl and  $\beta$  to the second: A. Nishinaga, T. Itahara, T. Shimizu, and T. Matsuura, *Tetrahedron Lett.*, 2467 (1976).
- (13) The reaction of singlet oxygen with  $\alpha,\beta$ -unsaturated ketones, a little studied process, can apparently yield unsaturated  $\beta$ -hydroperoxy ketones: H. E. Ensley and R. V. C. Carr, *Tetrahedron Lett.*, 513 (1977). Unfortunately, no isolation of these intermediates has been attempted.
- (14) S. Schlossberg, *Chem. Ber.*, **33**, 2425 (1900).
- (15) Satisfactory combustion analyses have been obtained for all stable key compounds reported herein.
- (16) V. Calò, F. Cimminale, L. Lopez, and P. E. Todesco, *J. Chem. Soc. C*, 3652 (1971); V. Calò, L. Lopez, G. Pesce, and P. E. Todesco, *Tetrahedron*, **29**, 1625 (1973).
- (17) R. Gleiter and M. Böhm, private communication.
- (18) Naphthoquinones **16** and **17** were prepared by air oxidation of alkaline

solutions of the corresponding hydroquinones (with vastly improved yields) rather than by the literature method: O. Diels and K. Alder, *Chem. Ber.*, **62**, 2337 (1929). Catalytic hydrogenation of **17** (5% Pd/C, C<sub>2</sub>H<sub>5</sub>OH, 1 Torr) furnished **18** quantitatively.


- (19) The epoxyquinones were separated by high pressure liquid chromatography and given their stereochemical assignments on the basis of their <sup>1</sup>H NMR spectra; see K. Tori, K. Kitahonoki, Y. Takano, H. Tanida, and T. Tsuji, *Tetrahedron Lett.*, 599 (1964).  
 (20) P. v. R. Schleyer, *J. Am. Chem. Soc.*, **89**, 699 (1967).  
 (21) This subject is to be discussed in more detail separately.  
 (22) NATO Postdoctoral Fellow with financial support provided by the CNRS.

Leo A. Paquette,\* Richard V. C. Carr, Francois Bellamy<sup>22</sup>  
 Evans Chemical Laboratories, The Ohio State University  
 Columbus, Ohio 43210  
 Received May 22, 1978

## $\alpha$ -Silyloxy and $\alpha$ -Hydroxy Manganese Alkyls. Generation via a New Five-Membered Metallocycle

Sir:


An ongoing research objective in our laboratory has been the generation of ligand types believed to be intermediate in the metal-catalyzed reduction of CO by H<sub>2</sub>.<sup>1-3</sup> Since the ultimate reduction products range from methane and methanol to higher alkanes and alcohols (Fischer-Tropsch process),<sup>4,5</sup> there is considerable interest in discerning the factors which control product *selectivity*, and in developing milder and homogeneous catalysts for accomplishing these transformations.<sup>6</sup> Communicated herein are our initial efforts at preparing complexes containing the  $\alpha$ -hydroxyalkyl ligand (**2**) and silylated derivatives thereof.  $\alpha$ -Hydroxyalkyl **2** has been proposed as a key mechanistic branch point in Fischer-Tropsch reactions,<sup>5,7</sup> and is virtually a required intermediate in the Union Carbide ethylene glycol synthesis.<sup>8</sup> We also wish to report the X-ray structural characterization of a novel metalocyclic system which serves as the fulcrum of our syntheses.



Benzoylformylmanganese pentacarbonyl (**3**, 10 mmol)<sup>1,9</sup> in THF (30 mL) was treated with Li(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>BH (12 mmol)<sup>10</sup> under nitrogen. Infrared monitoring<sup>11</sup> suggested that an octahedral cis-disubstituted manganese tetracarbonyl product was quantitatively formed. Although we previously reported that Li(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>BH attacks metal carbonyl acyls at carbon monoxide, yielding kinetically unstable anionic formyl complexes,<sup>2</sup> no characteristic formyl <sup>1</sup>H NMR resonance could be detected, even when this reaction was conducted at -60 °C in a NMR tube.

After 15 min, PPN<sup>+</sup> Cl<sup>-</sup> (10 mmol)<sup>12</sup> dissolved in CH<sub>2</sub>Cl<sub>2</sub> (50 mL) was added to the reaction mixture. Solvent was removed and the residue extracted with ether (400 mL). Concentration and cooling (0 °C) afforded a yellow precipitate (two crops, 5.73 g, 6.6 mmol, 66%), which could be rendered analytically pure by first washing with cold ether and then extracting into THF. Solvent removal yielded a gold powder (46% overall) for which analytical and spectral data suggested the structure PPN<sup>+</sup>-**4** (Scheme I).<sup>13</sup>

## Scheme I. Synthetic Routes to $\alpha$ -Silyloxy and $\alpha$ -Hydroxy Manganese Alkyls



The formation of **4** is proposed to occur via intermediate **5**. In light of this unprecedented mode of metalocycle closure, an X-ray crystal structure was undertaken.

Slow recrystallization from ether-hexane afforded suitable air-stable crystals (dec pt 142–144 °C) for study. X-ray data were obtained at -160 ± 5 °C with monochromated Mo K $\alpha$  (0.71069 Å) radiation on a Syntex PI automatic diffractometer. The general techniques employed have been previously described.<sup>14</sup> The unit cell was found to be triclinic, space group *P* $\bar{1}$  (*Z* = 2), with lattice parameters *a* = 10.016 (4), *b* = 15.772 (5), *c* = 15.698 (4) Å;  $\alpha$  = 86.55 (3),  $\beta$  = 89.09 (3),  $\gamma$  = 118.62 (3)°. Of 5697 reflections with  $2\theta < 40^\circ$ , 2994 with *I*  $\geq$  3 $\sigma$ (*I*) were used in the final refinement. All seven phenyl rings were refined as rigid groups (C–C = 1.39 Å, C–H = 1.00 Å). A series of least-squares refinements of positional parameters of all atoms and groups, anisotropic thermal parameters of manganese and phosphorous atoms, and isotropic thermal parameters of all other nonhydrogen atoms converged to *R* = 0.055 and *R*<sub>w</sub> = 0.058.<sup>15</sup> A difference Fourier map at this point revealed the location of the proton derived from Li(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>BH, which was included in two further cycles of refinement. A final difference Fourier map showed no peaks larger than 0.4 e/Å. The final standard deviation of an observation of unit weight was 1.28.

The arrangement of ligands about the manganese atom is depicted in Figure 1. The metalocycle is essentially planar, and undergoes clean thermolysis (THF, 65 °C, 6 days) to PPN[Mn(CO)<sub>5</sub>], benzaldehyde, and CO.

The metalocycle **4** undergoes two key ring-opening reactions (Scheme I). When in situ prepared Li<sup>+</sup>-**4** (1.01 mmol) was treated with (CH<sub>3</sub>)<sub>3</sub>SiCl (12 mmol), a product formed over a 2-h period. The solvent was removed and the residue column chromatographed (5% EtOAc in hexane). The product eluted rapidly and crystallized upon solvent removal (0.96 mmol, 95% yield, mp 51–53 °C). Based upon micro- and spectral analyses,<sup>16</sup> the silyloxyacyl structure **6** is assigned. This preparation may be executed equally well utilizing