

project was completed while one of us (P. N. C.) was associated with Smith, Kline and French Labs.

References

- (1) P. N. Craig, *J. Med. Chem.*, **15**, 144 (1972) (paper 1).
- (2) C. Hansch, *Accounts Chem. Res.*, **2**, 232 (1969).
- (3) T. S. Osdene, P. B. Russell, and L. Rane, *J. Med. Chem.*, **10**, 431 (1967).
- (4) A. Leo, C. Hansch, and D. Elkins, *Chem. Rev.*, **71**, 525 (1971).
- (5) C. G. Swain and E. C. Lupton, Jr., *J. Amer. Chem. Soc.*, **90**, 4328 (1968).
- (6) T. Yamamoto and T. Otsu, *Chem. Ind. (London)*, 787 (1967).
- (7) P. N. Craig, *J. Med. Chem.*, **14**, 680 (1971).
- (8) J. G. Topliss and R. J. Costello, *ibid.*, **15**, 1066 (1972).
- (9) T. Fujita, J. Iwasa, and C. Hansch, *J. Amer. Chem. Soc.*, **86**, 5175 (1964).
- (10) P. N. Craig, *Advan. Chem. Ser.*, No. 114 (1972).
- (11) G. W. Snedecor, "Statistical Methods," Iowa State University Press, Ames, Iowa, 1966.
- (12) A. Cammarata, *J. Med. Chem.*, **15**, 573 (1972).


Compounds Affecting the Central Nervous System. 3.¹ 3 β -Phenyltropan-2-ols

Sol J. Daum,* Mario D. Aceto, and Robert L. Clarke

Sterling-Winthrop Research Institute, Rensselaer, New York 12144. Received November 20, 1972

A group of 3 β -phenyltropanes bearing both axial and equatorial hydroxyl and acetoxyl groups at C-2 was prepared. The ability of these compounds to prevent and reverse reserpine-induced ptosis in mice and their effects on overt behavior are reported. The modification having a 2 β -hydroxyl group appeared to be at least as active as cocaine in the reserpine-induced ptosis screen and a more active stimulant. The optical antipodes of the more interesting compounds were prepared by resolving (\pm)-3-phenyltropidine. (\pm)-8a appears to be more active than the active enantiomer (–)-8a but no explanation is apparent. The ethylene bridge of the tropane structure was found necessary for activity.

The structure of cocaine (1) has been modified in a continued effort to obtain a variation of this molecule that would be a useful stimulant or antidepressant. This paper concerns that modification wherein a phenyl group is attached directly to the 3 β position of the tropane ring and the 2 β -carbomethoxy group is replaced by hydroxyl or acetoxyl in either the α or β configuration.



Shortly after we prepared the compound with the 2 α -hydroxyl configuration (2a), Lyle, *et al.*,² reported the synthesis of the same compound, but without biological data, by a similar procedure. Treatment of this 2 α -hydroxy compound 2a with ethyl chloroformate followed by saponification of the 2 α -carbonate 4 afforded the α -hydroxyurethane 5a. An alternate approach to 5a was through the 2 α -acetate 3a,³ prepared in our hands by the Ac₂O-pyridine method.[†] The acetate 3a was converted to the urethane 6a which was saponified to give the α -hydroxyurethane 5a. The 2-ketourethane 7a, obtained by Jones oxidation of 5a, was reduced with LiAlH₄ to afford 3 β -phenyltropan-2 β -ol (8a, 30%), 3 β -phenyltropan-2 α -ol (2a, 31%), and a third 3-phenyltropan-2-ol (9%) which we speculate has the 3 α -phenyl structure 9a. The axial 2 β -alcohol 8a had a band in its ir spectrum at 3455 cm⁻¹ that persisted even on dilution to a 0.001 M concentration, a characteristic of expected intramolecular hydrogen bonding with nitrogen.⁴ Acetylation of 8a with Ac₂O-pyridine afforded the 3 β -acetate 10 (Scheme I).

Similarly 2b was converted to the 2 α -acetate 3b.[†] Treatment of 3b with methyl chloroformate afforded the ure-

thane 6b which was saponified to give 5b. Jones oxidation followed by LiAlH₄ reduction gave equal amounts of the equatorial 2 α -alcohol 2b and the axial 2 β -alcohol 8b. We did not try to isolate the isomer 9b. The axial 2 β -alcohol 8b had a band in its ir spectrum at 3450 cm⁻¹ that persisted even on dilution to a 0.001 M concentration.

In order to gain insight into the role which chirality plays in this biological activity, the optical antipodes of 3a and 8a were prepared. 3-Phenyltropidine⁵ was resolved by means of its bitartrate salt and the resulting enantiomers were hydrolyzed using Lyle's² method to give (+)- and (–)-2a. These alcohols were then converted to (+)- and (–)-3a and -8a by the procedures described above. *trans*-1-Ethyl-4-phenyl-3-piperidinol acetate (11), a nonrigid analog of 3a, was prepared in order to relate rigidity factors with activity.

Biological Results. The compounds reported were evaluated by means of the reserpine-induced eyelid ptosis screen in mice⁶ and by observation of overt behavioral changes (see Table I). Cocaine was included in the study for comparative purposes.


In the ptosis prevention test, only compound (\pm)-8a showed activity, being perhaps slightly more active than cocaine. In the ptosis reversal test, compounds (\pm)-8a and (\pm)-3a were quite active; (\pm)-2a and (\pm)-8b were minimally active. It is interesting that conversion of the 2-equatorial hydroxyl of (\pm)-2a to the axial configuration (\pm)-8a resulted in a substantial increase in activity in the ptosis reversal test. Acetylation of this axial isomer [forming (\pm)-10] produced a drop in this activity. In sharp contrast, acetylation of the equatorial isomer (\pm)-2a caused an increase in ptosis reversal activity. Introduction of a *p*-methoxyl group into the aromatic ring of (\pm)-3a lowered its activity [see (\pm)-3b].

The signs of overt stimulation paralleled the results of the ptosis screen, with indications that stimulation lasted as long as 3 hr with (\pm)-8a and (\pm)-10. (\pm)-3a and (\pm)-8a appear to be more stimulative than cocaine.

Study of the optical antipodes of 3a and 8a revealed that only one enantiomer of each was active. These active enantiomers [(+)-3a and (–)-8a] differ in actual sign of optical rotation but belong to the same absolute configurational

[†]Ellefson³ reported failure in the preparation of 2 α -acetates 3a and 3b using Ac₂O; it required ketene. He reported mp 65–75° (petroleum ether) for 3a, mp 236.5–239° for 3a HBr salt, and mp 88–92° for 3b. We found mp 72–74° (pentane) for 3a, mp 259–261° for 3a HBr salt, and mp 100–102° (pentane) for 3b.

Scheme I



series with respect to asymmetric carbons 1 and 5 of the tropane nucleus; both are derived from (+)-phenyltropidine.

It is curious that *rac*-8a appears to be more active in both reversal and prevention of ptosis than *l*-8a. The presence of inactive *d*-8a in the racemate would normally be expected to act merely as a diluent. In order to confirm this observation, a concurrent test was run on (±)-8a and its separated optical antipodes, the results of which are shown in Table II. Again the racemic mixture appears to be significantly more active than the levo enantiomer. No explanation for this phenomenon is apparent.

Finally, the high level of activity of (±)-3a is somehow related to the presence of the ethylene bridge of the tropane moiety. The unbridged analog, (±)-11, is inactive. Whether this dramatic difference in activity is due to limitation of degrees of freedom with attendant differences in active-site complex stability or is simply a matter of steric inhibition at this site has yet to be determined.

Conclusions

(±)-3β-Phenyl-1αH,5H-tropan-2β-ol [(±)-8a] demonstrates about the same activity as does cocaine in the reserpine-induced ptosis test. It appears to be more stimulative than cocaine. This activity appears to reside in only one enantiomer of 8a but, curiously, the racemate seems to be more active than the active enantiomer alone. The ethylene bridge of the tropane system is required for activity.

Experimental Section[‡]

Ethyl (±)-2α-Hydroxy-3β-phenyl-1αH,5αH-nortropane-8-carboxylate (5a). Method A. Ethyl chloroformate (148 g, 1.4 mol) was

added dropwise over a 40-min period to a stirred solution of 5.92 g (0.27 mol) of (±)-3β-phenyl-1αH,5αH-tropan-2α-ol (2a)⁸ in 775 ml of refluxing C₆H₆. After being heated under reflux for an additional 3 hr, the reaction mixture remained at room temperature overnight. Et₂O and 6 N HCl were added. The organic layer was washed (saturated NaCl), dried (Na₂SO₄), and concentrated by heating *in vacuo* to afford 68.6 g of a straw-colored, viscous oil. The physical data indicated that this oil was a mixture of 4 and 5a: ir (CHCl₃) 3560 (m), 3380 (m), 1740 (ms), 1670 cm⁻¹ (vs); nmr δ 1.4 (t, 4 H), 1.5–3.2 (m, 8–9 H), 3.6–4.6 (m, 5 H), 7.3 ppm (m, 5 H). This mixture (61.6 g, 0.19 mol) in 408 ml of H₂O and 95°C/ml of EtOH containing 61.6 g of KHCO₃ was heated under reflux for 24 hr. Et₂O was added, separated, washed (saturated NaCl), dried (Na₂SO₄), and concentrated *in vacuo* to yield 56.5 g of a straw-colored, viscous oil. A short-path distillation of the product at 155–158° (0.01 mm) afforded the analytical sample 5a. *Anal.* (C₁₆H₂₁NO₃) C, H, N.

Method B. Ethyl chloroformate (54.2 g, 0.5 mol) in 100 ml of C₆H₆ was added over a 30-min period to a solution of 33.6 g (0.13 mol) of (±)-3β-phenyl-1αH,5αH-tropan-2α-ol acetate (3a)[†] in 200 ml of C₆H₆ being heated under reflux. After the mixture was refluxed for 3 hr and left overnight at room temperature, Et₂O and dilute HCl were added. The separated organic layer was washed (saturated NaCl), dried (Na₂SO₄), and concentrated *in vacuo* to yield 42 g of 6a as a viscous oil: ir (film) 1740 (vs), 1680 cm⁻¹ (vs); nmr δ 1.6–2.5 (m, 6 H + 3 H), 3.0 (m, 1 H, *ca.* *J* = 7, *J* = 10, and *J* = 10 Hz), 3.9–4.6 (m, 4 H), 5.1 (q, 1 H, *ca.* *J* = 3–4 and *J* = 10.5 Hz), 7.2 ppm (m, 5 H). *Anal.* (C₁₈H₂₃NO₄) OEt.

A solution of 40.4 g (0.13 mol) of 6a (used without further purification) in 250 ml of H₂O and 750 ml of EtOH containing 40 g of KHCO₃ was heated under reflux for 24 hr. The work-up was that used in method A and yielded 43.8 g of 5a.

Ethyl (±)-2-Oxo-3β-phenyl-1αH,5αH-nortropane-8-carboxylate (7a). A solution of 32.8 g (0.10 mol) of 5a in 250 ml of Me₂CO was treated with 8.7 N CrO₃–H₂SO₄ (Jones reagent) until the color remained orange (30 ml). Et₂O and H₂O were added and the Et₂O layer was separated, washed (saturated NaCl), dried (Na₂SO₄), and concentrated by warming *in vacuo* to afford 22.2 g of viscous oil 7a: *n*_D²⁵ 1.5380; ir (film) 1720 (s), 1680 cm⁻¹ (s); nmr δ 1.2 (t, 3 H), 1.6–2.5 (m, 6 H), 3.6 (q, 2 H), 4.6 (m, 2 H), 7.2 ppm (m, 5 H). This was used without further purification.

LiAlH₄ Reduction of Ethyl (±)-2-Oxo-3β-phenyl-1αH,5αH-nortropane-8-carboxylate (7a). A solution of 18.0 g (0.066 mol) of 7a in

[‡]All melting points are uncorrected. Nmr spectral measurements were made on Varian A-60 or HA-100 spectrophotometers using CDCl₃ as solvent unless otherwise indicated. (CH₃)₄Si was used as an internal standard. Infrared spectra were determined on a Model-21 Perkin-Elmer infrared spectrophotometer or, where dilution studies were done, a Beckmann IR-7 instrument. Spin-decoupling experiments were done with a Varian HA-100 instrument using a Hewlett-Packard audio oscillator 4204A. The mass spectra reported were measured with a Joelco JMS-1-OCS mass spectrograph. Where analyses are indicated only by symbols of the elements, analytical results obtained for these elements were within ±0.4% of the theoretical values. Optical rotations of salts were measured in H₂O, those of bases in CHCl₃.

⁸Lyle, Carle, Ellefson, and Spicer² reported 2a and its *p*-methoxy analog without melting points or purification procedures. In the Ellefson thesis,³ the unsubstituted 3-phenyl-2α-tropanol 2a was reported to melt at 120–122° with a footnote that some 3-phenyl-3-tropanol was present in the product. The *p*-methoxy analog 2b was reported with mp 100–102°. Following chromatographic purification of these substances, we found 2a to exhibit polymorphism: mp 120–122 and 129–131°; 2b melted at 124–126°.

Table I. Effect of Compounds on Reserpine-Induced Eyelid Ptosis and the Overt Behavior of Mice

Compd	Dose, ^a mg/kg ip	Ptosis prevention		Ptosis reversal		Overt behavior ^g
		MPS ^b	PV ^c	MPS ^b	PV ^c	
(±)-2a	30	3.5	0.574	3.3	0.574	30 + 50 mg/kg controls, questionable depression
	50	3.1	0.646	2.8	0.038	
	G.T. ^d	3.3		3.5		
(±)-3a ^e	0.6			3.1	0.278	0.6, 6, 18, 30 mg/kg controls, stimulation and running
	6			2.0	0.006	
	18	3.4	0.798	1.9	0.010	
	30	3.1	0.646	1.6	0.004	
	H ₂ O ^h	3.3		3.4, 3.5		
(±)-3a ^f	0.9			3.1	0.506	9 mg/kg controls, stimulation, running 0.5 hr; 26 and 44 mg/kg controls, depression, tremors at 0.5 hr, questionable stimulation at 3 hr
	9			2.5	0.038	
	26	2.9	0.328	2.1	0.014	
	44	3.0	0.506	2.1	0.014	
	H ₂ O ^h	3.3		3.4		
	26	3.3	0.960	2.8	0.234	
(±)-3a ^f	44	2.9	0.278	2.6	0.130	26 and 44 mg/kg controls, depression at 0.5 hr
	H ₂ O ^h	3.3		3.3		
	1			3.4	0.720	
(±)-3b	10			2.9	0.104	Drug controls at 10, 30, and 50 mg/kg, stimulation, running, gut spasms
	30	2.8	0.104	2.3	0.020	
	50	2.6	0.082	1.8	0.004	
	G.T. ^d	3.4		3.4, 3.5		
	0.1			3.5	1.0	
(±)-8a	0.5			3.0	0.104	10 mg/kg controls, mild stimulation at 0.5 hr; 30 mg/kg controls, stimulation, running at 0.5 hr, and mild stimulation at 3 hr; reversal at 0.5 hr; 30 + 50 mg/kg stimulation, running and jumping when touched
	1	2.9	0.328	2.8	0.020	
	10	2.8	0.194	2.5	0.002	
	30	2.4	0.050	1.8	0.002	
	50	2.1	0.010	1.8	0.002	
	G.T. ^d	3.3, 3.3		3.5, 3.5, 3.5		
	26	3.0	0.720	2.4	0.064	
	44	2.8	0.382	2.3	0.028	
	H ₂ O ^h	3.3		3.3		
	26	3.1	0.720	3.3	0.328	
(±)-8a ^f	44	2.9	0.328	2.9	0.574	26 and 44 mg/kg controls, depression, tremors at 0.5 hr, questionable stimulation at 3 hr
	H ₂ O ^h	3.3		3.6		
	30	3.3	1.00	2.8	0.194	
(±)-8b	50	3.1	0.720	2.6	0.082	Drug controls at 30 and 50 mg/kg, stimulation, hyperexcitable, jumping, squeaking
	G.T. ^d	3.3		3.4		
	26	2.9	0.234	2.5	0.064	
(±)-10 ^f	44	2.7	0.190	2.1	0.014	26 mg/kg controls, stimulation, running at 0.5 hr, and mild stimulation at 3 hr; 44 mg/kg controls, stimulation, running, some ataxia convulsions at 0.5 hr, and mild stimulation at 3 hr; reversal at 0.5 hr; 26 and 44 mg/kg stimulation, some squeaking, prevention 44 mg/kg, 1/8 dead before reserpine
	H ₂ O ^h	3.4		3.4		
	30	2.9	0.382	3.1	0.506	
(±)-11 ^f	50	2.6	0.104	3.1	0.506	Inactive
	H ₂ O ^h	3.3		3.4		
	1	3.5	0.960	3.4	0.960	
Cocaine (1)	10	3.4	0.646	2.1	0.010	50 mg/kg controls, excitement convulsions; 50-mg reversal, convulsions, complete recovery from reserpine, 1/8 dead at 100 mg at 0.5 hr; mild stimulation up to 5 hr
	30	2.9	0.160	1.5	0.002	
	50	2.4	0.010	0.05	0.000	
	H ₂ O ^h			3.4		

^aDoses calculated as free base. ^bMPS, mean ptotic score. ^cPV, probability value, significant if 0.05 or lower. ^d1% gum tragacanth mucilage controls. ^eTested as cyclohexanesulfamate. ^fTested as HCl salt. ^gEffect of compound alone unless otherwise specified as "reversal." ^hH₂O control.

300 ml of Et₂O was added dropwise to a suspension of 8 g of LiAlH₄ being stirred in 500 ml of Et₂O. After 2 hr of refluxing, 20 ml of H₂O was added with cooling. The salts were separated by filtration and the Et₂O was evaporated to afford 14.3 g of oil that was chromatographed on 750 g of silica gel using Et₂O-pentane-*i*-PrNH₂ (25:72:3) for elution. The fractions were combined on the basis of tlc analysis. The combined early fractions afforded 7.7 g of crude 8a contaminated with 9a. The later fractions afforded 5.7 g of crude 2a which, upon recrystallization from Et₂O, yielded 3.7 g of 2a, mp 119.5–122°.

The crude 8a combined with 1.7 g of mid-fraction material was chromatographed on 29 silica gel coated plates (Brinkmann PF 254 silica gel, 20 × 40 cm) having a 1-mm coating. The plates were developed with Et₂O-*i*-PrNH₂ (97:3). The least polar zone solid (5.5 g) was recrystallized from Et₂O to yield 4.2 g of (±)-3 β -phenyl-1 α H,5 α H-tropan-2 β -ol (8a), mp 96–98°. The analytical sample (from Et₂O) had mp 98–100°; ir (CCl₄) 3615, 3455 cm⁻¹; ir (CCl₄, 0.05–0.001 M) 3455 cm⁻¹. Anal. (C₁₄H₁₉NO) C, H, N.

The acetate 10 HCl salt (Ac₂O-pyridine) from acetone exhibited

polymorphism, forming a clear melt at 245–246°, resolidifying, and then decomposing at 269°. Anal. (C₁₆H₂₁NO₂·HCl) C, H, Cl.

A mid-zone solid (1.7 g) was recrystallized from Et₂O to afford 1.2 g of 9a, mp 135–137°. The analytical sample (from Et₂O) had mp 136–138°; ir (CCl₄, 0.05–0.001 M) 3605 cm⁻¹; nmr δ 1.2–2.2 (m, 7 H), 2.2 (s, 3 H), 3.3 (m, 3 H), 4.3 (m, 1 H), 7.2 ppm (m, 5 H). Anal. (C₁₄H₁₉NO) C, H, N.

The acetate of 9a showed poor crystallizing properties but gave only 1 spot by tlc analysis; nmr spin decoupling studies indicated *J*_{1,2} = ca. 7 Hz. The most polar zone solid (1.5 g) was recrystallized from Et₂O to yield another 0.7 g of 2a, mp 119–121°.

(±)-3 β -(*p*-Methoxyphenyl)-1 α H,5 α H-tropan-2 β -ol (8b). (±)-3 β -(*p*-Methoxyphenyl)-1 α H,5 α H-tropan-2 α -ol acetate (3b)[†] was converted to urethane 6b using methyl chloroformate as the reagent in method B above. Anal. (C₁₈H₂₃NO₃) C, H, N. Saponification of 6b by the described method afforded oily 5b. Anal. (C₁₆H₂₁NO₄) C, H, N.

Oxidation of alcohol 5b as described for 5a afforded the 2-oxo compound 7b as an oil which was used without purification. Reduction of 7b (12.3 g, 0.043 mol) with LiAlH₄ yielded 2.5 g (24%) of the

Table II. Results of Concurrent Testing of (±)-8a, (+)-8a, and (−)-8a in Reversal of Reserpine-Induced Eyelid Ptosis (Mice)

Compd	Dose ^a	MPS ^b	PV ^c	Overt behavior ^d
(±)-8a	1	3.1	0.506	Controls, 10 and 30, stimulation and running, some biting and squeaking at 0.5 hr
	10	1.9	0.010	
	30	1.8	0.002	
	Control	3.4		
(−)-8a	0.9	3.3	0.720	Controls, 9, ? mild stimulation at 0.5 hr, mild stimulation at 3 hr; controls, 26, mild stimulation and running at 0.5 hr, mild stimulation at 3 hr
	9	2.8	0.194	
	26	2.3	0.014	
	Control	3.4		
(+) -8a	0.9	3.3	0.720	Controls, 26, ? depression at 0.5 hr, ? stimulation at 3 hr
	9	3.0	0.382	
	26	2.8	0.104	
	Control	3.4		

^aDoses calculated as free base, mg/kg ip. ^bMPS, mean ptotic score. ^cPV, probability value, significant if 0.05 or lower. ^dEffect of compound alone.

β-alcohol 8b, mp 85.5–87.5°, which on recrystallization from Et₂O melted at 86–87.5°; ir (CCl₄, 0.05–0.001 M) 3450 cm^{−1}. Anal. (C₁₂H₂₁NO₂) C, H, N. Also obtained from the reduction was 2.4 g (23%) of the α-alcohol 2b, mp 124–126° (Et₂O), and an oil in intermediate polarity amounting to 0.3 g which probably contains 9b.

Resolution of (±)-3-Phenyltropidine. A solution of 92 g (0.61 mol) of (+)-tartaric acid in 240 ml of H₂O was treated with 122.2 g (0.61 mol) of 3-phenyltropidine in 240 ml of Me₂CO. The solution was cooled in an ice bath and the precipitated bitartrate salt was collected. Fractional crystallization from 1:1 H₂O–acetone (6 ml/g) afforded 38.6 g of an apparently hydrated bitartrate: mp 57–60°; [α]_D²⁵ +23.1°.

The mother liquors were combined and made alkaline with concentrated NH₄OH. Et₂O extraction afforded 97.4 g of basic material which was dissolved in 200 ml of acetone and added to 76 g (0.50 mol) of (−)-tartaric acid in 200 ml of H₂O. The solution was cooled in an ice bath and the precipitated bitartrate salt was collected. Fractional crystallization from 1:1 H₂O–acetone (6 ml/g) afforded 72.5 g of an apparently hydrated bitartrate: mp 56–61°; [α]_D²⁵ −23.6°.

After recycling the mother liquors, another 64 g of bitartrate, mp 57–59°, [α]_D²⁵ +21.6°, and 35.8 g of bitartrate, mp 55–62°, [α]_D²⁵ −21.5°, was obtained.

(+)-3β-Phenyltropidine. A solution of 103 g of (+)-phenyltropidine (+)-bitartrate·(H₂O)_x in 200 ml of H₂O was made alkaline with concentrated NH₄OH. The Et₂O extract was dried (Na₂SO₄) and concentrated to afford 44.4 g of crude (+)-3β-phenyltropidine. Distillation at 93–101° (0.3–0.7 mm) gave 42.4 g of (+)-3β-phenyltropidine: *n*_D²⁵ 1.5785; [α]_D²⁵ +37.9°. Anal. (C₁₄H₁₇N) C, H, N.

(−)-3β-Phenyltropidine. (−)-3β-Phenyltropidine (−)-bitartrate·(H₂O)_x (108.3 g) in a similar fashion afforded 50 g of crude (−)-3β-phenyltropidine. Distillation at 94–110° (0.3–0.4 mm) gave 43.5 g of (−)-3β-phenyltropidine: *n*_D²⁵ 1.5781; [α]_D²⁵ −39.6°. Anal. (C₁₄H₁₇N) C, H, N.

(+)-3β-Phenyl-1αH,5αH-tropan-2α-ol [(+)-2a]. A solution of 40.6 g (0.20 mol) of (+)-3β-phenyltropidine in 225 ml of THF was added dropwise to 460 ml of 1 M BH₃ in THF with stirring in an ice bath. After being heated for 5 hr under reflux and standing at room temperature overnight, the solution was carefully treated with 35 ml of H₂O added dropwise followed by 170 ml of 3 N NaOH. A 30% solution (65 ml) of H₂O₂ was added dropwise at a rate that maintained reflux. After being heated for an additional 1 hr under reflux, the reaction mixture was cooled. Et₂O was added and the layers were separated. The Et₂O was dried (Na₂SO₄) and evaporated. The residue was taken up in 300 ml of MeOH and was shaken with

Raney Ni for 1 hr. Filtration of the catalyst and evaporation afforded 44 g of crude product. Crystallization from Et₂O gave 21.4 g of (+)-2a: mp 147–148°; [α]_D²⁵ +42.4°. A second crop, recrystallized from Et₂O, gave (±)-2a: mp 120–123°; [α]_D²⁵ 0°. The analytical sample of (+)-2a** melted at 147.5–148.5°, [α]_D²⁵ +42.5°. Anal. (C₁₄H₁₉NO) C, H, N.

(+)-3β-Phenyl-1αH,5αH-tropan-2α-ol acetate [(+)-3a]** melted at 71.5–73.5° (pentane), [α]_D²⁵ +64.5°. Anal. (C₁₆H₂₁NO₂) C, H, N. HCl salt of (+)-3a: mp 263–264° dec (acetone); [α]_D²⁵ +33.5°. Anal. (C₁₆H₂₁NO₂·HCl) C, H, Cl.

(−)-3β-Phenyl-1αH,5αH-tropan-2α-ol [(−)-2a]. In a similar procedure 42 g of (−)-3β-phenyltropidine afforded 22.1 g of (−)-2a: mp 146.5–148.5°; [α]_D²⁵ −41.3°. A second crop, recrystallized from Et₂O, gave (±)-2a: mp 119–121°; [α]_D²⁵ −3.6°. The analytical sample of (−)-2a** melted at 147.5–148.5°, [α]_D²⁵ −42.0°. Anal. (C₁₄H₁₉NO) C, H, N.

(−)-3β-Phenyl-1αH,5αH-tropan-2α-ol acetate [(−)-3a]** melted at 71.5–73° (pentane), [α]_D²⁵ −63.8°. Anal. (C₁₆H₂₁NO₂) C, H, N. HCl salt of (−)-3a: mp 264–265° dec (acetone); [α]_D²⁵ −34°. Anal. (C₁₆H₂₁NO₂·HCl) C, H, Cl.

The following resolved compounds** were prepared by the above described procedures giving ultimately the enantiomers of 8a. Ethyl 2α-hydroxy-3β-phenyl-1αH,5αH-nortropane-8-carboxylate (optically active 5a) from (+)-2a was characterized spectrally.

Ethyl 2α-hydroxy-3β-phenyl-1αH,5αH-nortropane-8-carboxylate (optically active 5a) from (−)-2a was characterized spectrally.

Ethyl 2-oxo-3β-phenyl-1αH,5αH-nortropane-8-carboxylate (optically active 7a) from (+)-2a was characterized spectrally.

Ethyl 2-oxo-3β-phenyl-1αH,5αH-nortropane-8-carboxylate (optically active 7a) from (−)-2a was characterized spectrally.

(+)-3β-Phenyl-1αH,5αH-tropan-2β-ol [(+)-8a] from (−)-2a: mp 122–124° (Et₂O); [α]_D²⁵ +113.1°. Anal. (C₁₄H₁₉NO) C, H, N. HCl salt of (+)-8a: mp 297–298° dec (CH₃CN); [α]_D²⁵ +84.6°.

Anal. (C₁₄H₁₉NO·HCl) C, H, Cl.

(−)-3β-Phenyl-1αH,5αH-tropan-2β-ol [(−)-8a] from (+)-2a: mp 121–122° (Et₂O); [α]_D²⁵ −114.7°. Anal. (C₁₄H₁₉NO) C, H, N.

HCl salt of (−)-8a: mp 293–295° dec (CH₃CN); [α]_D²⁵ −84.0°. Anal. (C₁₄H₁₉NO·HCl) C, H, Cl.

trans-(±)-1-Ethyl-4-phenyl-3-piperindol-3-Acetate Hydrochloride (11). A mixture of 3.0 g of *trans*-(±)-1-ethyl-4-phenyl-3-piperidinol† in 7.5 ml of Ac₂O and 15 ml of pyridine was heated at 100° for 30 min and left overnight. The oily acetate, isolated in the usual manner, was converted to its HCl salt: heavy needles from acetone; mp 186.5–188°. Anal. (C₁₅H₂₁NO₂·HCl) C, H, Cl.

Acknowledgment. We wish to thank Mr. Harry C. Bentley for the ptosis screening and our entire Physical Department for their spectral interpretation.

References

- (1) S. J. Daum, A. J. Gambino, M. D. Aceto, and R. L. Clarke, *J. Med. Chem.*, **15**, 509 (1972) (paper 2).
- (2) R. E. Lyle, K. R. Carle, C. R. Ellefson, and C. K. Spicer, *J. Org. Chem.*, **35**, 802 (1970).
- (3) C. R. Ellefson, Ph.D. Thesis, University of New Hampshire, Durham, N. H., 1968.
- (4) W. A. M. Davies, J. B. Jones, and A. R. Pinder, *J. Chem. Soc.*, 3504 (1960).
- (5) A. C. Cope and A. D'Addieco, *J. Amer. Chem. Soc.*, **73**, 3419 (1951).
- (6) M. D. Aceto and L. S. Harris, *Toxicol. Appl. Pharmacol.*, **7**, 329 (1965).
- (7) R. E. Lyle, D. H. McMahon, W. E. Kruge, and C. K. Spicer, *J. Org. Chem.*, **31**, 4164 (1966).

*Nmr and ir spectra were identical with spectra of unresolved samples.

††The authors thank Dr. W. B. Dickinson of these laboratories for a sample of this unpublished compound. The 1-methyl homolog is described; see ref 7.

#This isomer is not quite optically pure. Complete purification was accomplished in the next step.