ficient regeneration of 9. The ultraviolet spectrum of 12, no maximum >190 nm, gives no clear evidence for an interaction between the neighboring, but nonconjugated, π systems. Photoelectron spectroscopy will provide a better probe of this important possibility.

Diene 12 is not thermally stable. At 90 °C, Cope rearrangement to 13 occurred quantitatively over a 2-day period.^{11,12} Apparently, the trans-skeletal interactions in 12, probably of simple steric origin, are of sufficient magnitude (and therein remarkable) to destabilize 12 relative to the all-cis tetrasubstituted cyclobutane 13.



Catalytic hydrogenation of **12** gave **1**, perhydro[0.0]paracyclophane: IR ν 2920, 1480, 1460 cm⁻¹. The high symmetry of **1** on the NMR time scale is apparent from its two line proton-decoupled ¹³C NMR spectrum: δ 30.2 (4 C, d, J = 130 Hz) and 24.6 ppm (8 C, t, J = 126 Hz). The ¹H NMR spectrum shows a significant downfield shift for the endo protons, presumably a result of steric compression: ¹³ δ 2.03 (8 H, br d, J= 9 Hz), 1.95 (4 H, br s), 1.48 ppm (8 H, br d, J = 9 Hz).

Although more conveniently drawn in the symmetric, double boat conformation 1a, 1 is better represented as the lower energy, double twist boat 1b and its mirror image 1c. It is interesting to speculate whether passage from 1b to 1c is by way of the high energy D_{2h} conformer 1a.



Application of our synthetic scheme to somewhat modified intermediates should permit us access to hexaprismane (14) and, via rhodium(I) induced opening of 14, to tetrahydro[0.0]paracyclophane (15), *p*-dibenzene. We shall report our progress with these systems in due course.



Acknowledgments. The research of the principal investigator is supported by the National Science Foundation (MPS-75-04123) and the National Cancer Institute (PHS-CA-12961). Funds for purchase of nmr instrumentation were provided, in part, by the National Cancer Institute (PHS-CA-14599) via The University of Chicago Cancer Research Center and by the National Science Foundation.

References and Notes

- See, for example, L. Cassar, P. E. Eaton, and J. Halpern, J. Am. Chem. Soc., 92, 6366 (1970); W. G. Dauben, C. H. Schallhorn, and D. L. Whalen, *ibid.*, 93, 1446 (1971); L. A. Paquette and J. C. Stowell, *ibid.*, 92, 2584 (1970), and 93, 2459 (1971); L. A. Paquette, R. S. Beckley, and W. B. Farnham, *ibid.*, 97, 1089 (1975); L. A. Paquette, J. S. Ward, R. A. Boggs, and W. B. Farnham, *ibid.*, 97, 1101 (1975); L. A. Paquette, Acc. Chem. Res., 4, 280 (1971); P. E. Eaton and D. R. Patterson, J. Am. Chem. Soc., 100, 2573 (1978).
- (2) See, for example, L. Cassar, P. E. Eaton, and J. Halpern, J. Am. Chem. Soc., 92, 3515 (1970); W. C. Dauben and A. J. Kielbania, Jr., *ibid.*, 93, 7345 (1971); P. E. Eaton, L. Cassar, R. A. Hudson, and D. R. Hwang, J. Org. Chem., 41, 1445 (1976).
- (3) N. C. Yang and Jacqueline Libman, J. Am. Chem. Soc., 94, 9228 (1972); N. C. Yang, C. V. Neywick, and K. Srinivasachar, Tetrahedron Lett., 4313 (1975).
- (4) For other attempts, see H. Babad, Ph.D. Thesis, University of Illinois, Urbana,

III., 1961.

- (5) J. S. Newcomer and E. T. McBee, *J. Am. Chem. Soc.*, **71**, 946 (1949); E. T. McBee, D. L. Crain, R. D. Crain, L. R. Belohlav, and H. P. Barendlin, *ibid.*, **84**, 3557 (1962). It is not generally appreciated that the pure ketal is crystalline, mp 29–31 °C; we are indebted to Victor Mark for supplying us long ago with a seed crystal.
- 6) I. A. Akhtar, G. I. Fray, and J. M. Yarrow, J. Chem. Soc. C, 812 (1968).
- (7) Satisfactory analytical and/or spectroscopic data were obtained for all new compounds reported here.
- (8) Cf. P. E. Eaton and T. W. Cole, Jr., J. Am. Chem. Soc., 86, 3157 (1964);
 K. V. Scherer, Jr., G. A. Ungefug, and R. S. Lunt III, *ibid.*, 88, 2859 (1966);
 R. J. Stedman, L. S. Miller, L. D. Davis, and J. R. E. Hoover, J. Org. Chem., 35, 4169 (1970).
- (9) Cf. K. V. Scherer, Jr., R. S. Lunt III, and G. A. Ungefug, *Tetrahedron Lett.*, 1199 (1965); N. B. Chapman, J. M. Key, and K. J. Toyne, *J. Org. Chem.*, **35**, 3860 (1970).
- (10) In analogy to [Rh(norbornadiene)Cl]₂, the new rhodium(I)-diene complex is taken to be a chloro bridged dimer.
- (11) D. Valentine, N. J. Turro, Jr., and G. S. Hammond, J. Am. Chem. Soc., 86, 5202 (1964). None of the Diels-Alder dimers of 1,3-cyclohexadiene was found.
- (12) This behavior is similar to that of syn-tricyclo [4.2.1.1^{2.5}]deca-3,7-diene; see C. W. Doecke, G. Klein, and L. A. Paquette, J. Am. Chem. Soc., 100, 1596 (1978).
- (13) B. V. Cheney, J. Am. Chem. Soc., 90, 5386 (1968), and ref 1-8 therein.

Philip E. Eaton,* Utpal R. Chakraborty

Searle Chemistry Laboratory, Department of Chemistry The University of Chicago, Chicago, Illinois 60637 Received November 21, 1977

A Novel Synthesis of (\pm) -4-Demethoxydaunomycinone

Sir:

The anthracyclines daunorubicin (1) and adriamycin (2) are of current interest in view of their activity against various experimental tumors as well as some types of human cancer.¹



The activity of these compounds can be improved by structural modification, as shown by the recent report that 4-demethoxydaunorubicin is four to eight times more active than daunorubicin itself.² We now wish to report a basically new, simple, and efficient synthesis of the dimethyl ether of the corresponding aglycone 4-demethoxydaunomycinone (3).

Our synthetic strategy centers on the construction of ring



© 1978 American Chemical Society

A by the Diels-Alder addition of a reactive *o*-quinodimethane intermediate to the olefinic portion of an α,β -unsaturated ketone as shown in eq 1.³ The generation of an *o*-quinodimethane from an *o*-xylene derivative and its trapping by a dienophile has ample precedence in the literature.⁴

Phthalic anhydride was condensed with the readily prepared 2,3-dimethylhydroquinone (4)⁵ (AlCl₃/NaCl, 190 °C, 2 min) to give, after heating with dilute hydrochloric acid, 2,3-dimethyl-1,4-dihydroxyanthraquinone (5) (81%, mp 252-253 °C).^{6,7} Methylation of 5 (Me₂SO₄/K₂CO₃, refluxing 2-bu-



tanone, 12 h) afforded the dimethoxy compound 6 (95%, mp 159–160 °C) which was brominated photochemically (NBS/CCl₄, reflux, 8 h) to give the key dibromide 7 (95%, mp 171–173 °C). Reaction of dibromide 7 (Zn dust, DMF, 25 °C, 6 h) in the presence of excess methyl vinyl ketone gave, after aqueous workup, the tetracyclic ketone 8 (52%, mp 145–147 °C). Oxidation of ketone 8 (KO-*t*-Bu/O₂, DMF, -20 °C, 1 h) followed by reduction ((EtO)₃P, DMF, -20 °C, 1 h)⁸ gave, after mild acid hydrolysis, the hydroxy ketone 9 (55%, mp 184–186 °C).⁹

Since the conversion of 9 to 4-demethoxydaunomycinone (3) has already been described, 2.9,10 our synthesis of 9 also constitutes a new synthesis of 3.

We are currently studying variations of this *o*-quinodimethane approach, in particular the use of oxy derivatives of methyl vinyl ketone in order to provide a direct route to tetracyclic ketones containing an oxygenated side chain.

References and Notes

- (a) F. Arcamone, G. Franceshi, P. Crezzi, G. Cassinelli, W. Barbiere, and R. Mondelli, J. Am. Chem. Soc., 86, 5334 (1964); (b) J. Bernard, R. Paul, M. Boiron, C. Jacquillat, and R. Maral, Ed., "Rubidomycin", Springer-Verlag, New York, N.Y., 1969; (c) R. B. Livingston and S. K. Carter, "Daunomycin", Chemotherapy Fact Sheet, National Cancer Institute, Bethesda, Md. 1970; (d) F. Arcamone, G. Francheschi, and S. Penco, *Tetrahedron Lett.*, 1007 (1969); (e) R. H. Blum and S. K. Carter, Ann. Intern. Med., 80, 249 (1974); (f) M. Ghione, *Cancer Chemother. Rep., Part 3*, 83 (1975).
- (2) F. Arcamone, L. Bernardi, P. Giardino, B. Patelli, A. DiMarco, A. M. Casazza, G. Pratesi, and P. Reggiani, *Cancer Treatment Rep.*, **60**, 829 (1976).
- (3) For alternative Diels-Alder approaches in which ring A is constructed from a diene rather than from a dienophile, see (a) T. R. Kelly, J. W. Gillard, and R. N. Goerner, Jr., *Tetrahedron Lett.*, 3873 (1976); (b) A. S. Kende, Y. Tsay, and J. E. Mills, J. Am. Chem. Soc., 98, 1967 (1976); (c) W. W. Lee, A. P. Martinez, T. H. Smith, and D. W. Henry, J. Org. Chem., 41, 2296 (1976).
- (4) (a) M. P. Cava, A. A. Deana, and K. Muth, J. Am. Chem. Soc., 81, 6458 (1959); (b) K. Alder and M. Fremery, *Tetrahedron*, 14, 190 (1961).
- (5) J. Lars, G. Nilsson, H. Sievertsson, and H. Selander, Acta Pharm. Suecica, 5, 215 (1968); Chem. Abstr., 69, 106132j (1968).
- (6) C. Marshalk, F. Koenig, and N. Ouroussoff, Bull. Soc. Chim. Fr., 3, 1545 (1936).
- (7) All new compounds gave satisfactory combustion and spectroscopic data consistent with assigned structures. Compounds 5–7 were recrystallized from glacial acetic acid; compounds 8–9 were chromatographed (SiO₂, MeOH/CHCl₃) and then recrystallized from methanol.
- (8) The procedure followed was that employed with representative 20-ketosteroids: J. N. Gardner, F. E. Carlon, and O. Gnoj. J. Org. Chem., 33, 3294 (1968).
- (9) C. M. Wong, D. Popien, R. Schwenk, and J. Te Raa, *Can. J. Chem.*, 49, 2712 (1971).
- (10) A. S. Kende, D. P. Curran, Y. Tsay, and J. E. Mills, *Tetrahedron Lett.*, 3537 (1977).

Francis A. J. Kerdesky, Michael P. Cava*

Department of Chemistry, University of Pennsylvania Philadelphia, Pennsylvania 19104 Received January 19, 1978

A General, Selective, and Facile Method for Ketone Synthesis from Acid Chlorides and Organotin Compounds Catalyzed by Palladium

Sir:

Transition metal catalyzed coupling reactions of organic halides with Grignard reagents or organolithium compounds generally are not applicable for ketone synthesis via the acid chlorides, since the organometallics react with the product ketone.¹ Alkylrhodium complexes, prepared from a rhodium complex and organolithium or Grignard reagents, however, may be used for alkylation of acid chlorides, giving alkyl ketones.^{2,3} Unfortunately, in addition to being a two-stage synthesis, this reaction is stoichiometric with respect to rhodium and will not tolerate a number of other functional groups on the acid chloride.

We have found that organotin compounds readily undergo a palladium catalyzed coupling with acid chlorides, thereby providing a general and simple method for preparation of ketones (eq 1).⁴ The reaction is general both with respect to the organotin compound and the acid chloride (Table I).

$$RCOCl + R'_4Sn \xrightarrow{PhCH_2Pd(PPh_3)_2Cl (1)}{HMPA}$$

$$RCOR' + R'_{3}SnCl$$
 (1)

The following features make this method synthetically attractive. (1) The yields are high, and in many cases virtually quantitative. (2) The reaction can be carried out in the presence

© 1978 American Chemical Society