

The Cleavage of t-Adamantyloxy-radicals

By R. M. BLACK and G. B. GILL*

(Department of Chemistry, The University, Nottingham NG7 2RD)

Summary Two-step reaction sequences to 4-substituted tricyclo[4,3,1,0^{3,8}]decanes and 3-substituted tricyclo[3,3,1,0^{3,7}]nonanes are reported: the key step in each case involves the scission of a t-adamantyloxy-radical.


INTEREST has been shown recently in the chemistry of tricyclo[3,3,1,0^{3,7}]nonanes^{†1} and of tricyclo[4,3,1,0^{3,8}]decanes.^{†2,3} In the second case, tricyclo[4,3,1,0^{3,8}]decan-4-one (X) was prepared using the adamantan-2-yl → tricyclo[4,3,1,0^{3,8}]decan-4-yl cation rearrangement.³ The migration appears to proceed in a synthetically useful yield only if it is of the pinacol-pinacolone type (*i.e.* an adamantan-1-ol-2-yl cation was employed).^{3b} Nevertheless, starting from commercially available materials (*i.e.* adamantane-1-carboxylic acid) the preparation of the tricyclic ketone

requires a ten-stage sequence in which the overall yield is at best 8.5%. The routes available to 3-substituted tricyclo[3,3,1,0^{3,7}]nonanes are scarcely better. We report here on superior routes to both systems by way of the scission of the t-adamantyloxy-radicals (II, VII) which were conveniently generated from the corresponding hypohalites (I, VI).

(i) *Hypochlorites*. The following illustrates the procedure for compounds (I). A solution of 2-methyladamantan-2-ol hypochlorite (I; X = Cl, R = Me) in dry CCl₄ was prepared by the action of chlorine monoxide on the parent alcohol⁴ at 0°. Thermolysis at the reflux temperature (8 h) or photolysis at 0° (30 min) yielded 3-chlorobicyclo[3,3,1]non-7-yl methyl ketone (IV; X = Cl, R = Me) in 75–80% yield, b.p. 110–116° (bath)/0.3 mm. The assigned structure follows from the analytical and spectral data: *i.r.* (film).

† Commonly (and ambiguously) referred to as noradamantanes and isoadamantanes (protoadamantanes), respectively.

1705, 1355, and 770 cm^{-1} ; 100 MHz n.m.r. (CCl_4) τ 5.60 (1H, t of t, $J_{\text{AX}} + J_{\text{BX}}$ 36 Hz; 3-H), 7.35 (1H, t of t, $J'_{\text{AX}} + J'_{\text{BX}}$ 37 Hz; 7-H), 8.01 (3H, s; Me), and 7.70–8.70 (12H, m); m/e (M^+) 202, and 200.




The thermolabile chloroketone was converted by base (e.g. KOH-MeOH) into methyl tricyclo[3,3,1,0^{3,7}]non-3-yl ketone (V; R = Me), b.p. 60–65° (bath)/0.7 mm. (80%); oxime m.p. 152–153° (mixture of isomers). The conversion (I) \rightarrow (V) could be reduced to a single stage merely by eluting the crude reaction product (in benzene) over basic alumina (66% yield based on I; X = H, R = Me). The product was not tricyclo[4,3,1,1^{3,8}]undecan-4-one,⁵ and the assigned structure follows from the physical data: i.r. (CCl_4) 1697, and 1355 cm^{-1} ; 100 MHz n.m.r. (CCl_4) τ 7.42 (1H, t of t, $J_{\text{AX}} + J_{\text{BX}}$ 14 Hz; 7-H), 7.71br (2H, m; 1-H and 5-H), 7.96 (3H, s; Me), and 8.00–8.46 (10H, m); m/e (M^+) 164.

Haloform degradation (KOH-Br₂-aq. dioxan) of (V; R = Me) gave tricyclo[3,3,1,0^{3,7}]nonane-1-carboxylic acid (V; R = OH) in 88% yield, m.p. 108–109° (lit.^{1a} m.p. 106–107°); i.r. (CCl_4) 3300–2600br, 1693, 1416, 1310, and 1293 cm^{-1} ; 60 MHz n.m.r. (CCl_4) τ -1.65 (1H, s; CO_2H), 7.28 (1H, t of t, $J_{\text{AX}} + J_{\text{BX}}$ 14 Hz; 7-H), 7.70–8.55 (12H, m).

In contrast, only a very low yield of tricyclo[4,3,1,0^{3,8}]decan-4-one (X) was obtained in the analogous sequence from (VI; X = Cl).

(ii) *Hypoiodites*. The hypoiodite (VI; X = I) was prepared *in situ* in dry benzene by the action of lead tetra-acetate and iodine (in the presence of dry calcium carbonate)⁶ on adamant-1-ol. Thermolysis at 55–60° (higher temperatures decomposed the thermolabile product) for 1.5 h gave a colourless crystalline product (40–45%), m.p. 78–79° (from MeOH and then light petroleum at -78°), identified as *endo*-3-iodomethylbicyclo[3,3,1]nonan-7-one (IX; X = I); i.r. (CCl_4) 1708 cm^{-1} ; 60 MHz n.m.r.

(CCl_4) τ 7.05 (2H, d, J 6.3 Hz; CH_2I), 7.45–8.60 (11H, m), 9.14br (2H, t, separation between outer lines *ca.* 22 Hz; boat axial 2-H and 4-H); m/e (M^+) 278. The data favour conformer (IXb).



The iodoketone was converted by base (e.g. KOH-MeOH, or pyridine) into tricyclo[4,3,1,0^{3,8}]decan-4-one (X; 70–75%), m.p. 212–214° (from aq. MeOH or sublimation; lit.^{3b} m.p. 205–208°).

Reduction of (X) by lithium aluminium hydride gave a 2 : 1 mixture of the epimeric alcohols (XI; 90%), m.p. 232–234° (cf. ref. 3b). Pyrolysis of (XI) xanthate gave tricyclo[4,3,1,0^{3,8}]dec-4-ene (XII; 34%), m.p. 183–185°; i.r. (CCl_4) 3035, 1633, 870, and 696 cm^{-1} ; 100 MHz n.m.r. (CCl_4) τ 3.89 (2H, m; 4-H and 5-H), 7.38 (2H, m; 3-H and 6-H), and 7.50–8.66 (10H, complex m). Attempted dehydration of mixture (XI) by the toluene-*p*-sulphonyl chloride-pyridine method⁵ gave instead mainly adamant-2-ol toluene-*p*-sulphonate (I; X = $\text{C}_7\text{H}_7\text{SO}_2$, R = H).

The analogous reactions of hypoiodites (I; X = I, R = Me, Et, or Ph) took an abnormal course; thus (I; X = I, R = Me) gave a yellow crystalline di-iodoether ($\text{C}_{11}\text{H}_{16}\text{I}_2\text{O}$).

We thank the S.R.C. for a studentship (to R. M. B.).

(Received, June 1st, 1970; Com. 844.)

¹ (a) B. R. Vogt and J. R. E. Hoover, *Tetrahedron Letters*, 1967, 2841; (b) P. von R. Schleyer and E. Wiscott, *ibid.*, p. 2845; (c) A. Nickon, G. D. Pandit, and R. O. Williams, *ibid.*, p. 2851; (d) B. R. Vogt, S. R. Suter, and J. R. E. Hoover, *ibid.*, 1968, 1609.

² B. R. Vogt, *Tetrahedron Letters*, 1968, 1575.

³ (a) M. L. Sinnott, H. J. Storesund, and M. C. Whiting, *Chem. Comm.*, 1969, 1000; (b) J. R. Alford and M. A. McKerver, *ibid.*, 1970, 615.

⁴ S. Landa, J. Vais, and J. Burkhard, *Coll. Czech. Chem. Comm.*, 1967, 570.

⁵ R. M. Black and G. B. Gill, *J. Chem. Soc. (C)*, 1970, 671.

⁶ For example see: von Ch. Meystre, K. Heusler, J. Kalvoda, P. Wieland, G. Anner, and A. Wettstein, *Helv. Chim. Acta*, 1962, 45, 1317.