

InBr₃-catalyzed stereoselective synthesis of trans-2,6-disubstituted 3,6-dihydro-2H-pyrans

J. S. Yadav^{*}, V. Sunitha, B. V. Subba Reddy, P. P. Das, E. Gyanchander

Division of Organic Chemistry-I, Indian Institute of Chemical Technology, Hyderabad 500 007, India

Received 21 September 2007; revised 19 November 2007; accepted 28 November 2007

Available online 4 December 2007

Abstract

A new method for the stereoselective synthesis of trans-2,6-disubstituted 3,6-dihydro-2H-pyrans with a variety of substitution patterns is described, involving Lewis acid induced tandem allylation or cyanation of δ -hydroxy- α,β -unsaturated aldehydes to produce dihydropyrans in good yields and with trans-selectivity. This method is very useful for the synthesis of trans-2,6-disubstituted dihydropyran ring-containing natural products such as laulimalide, scytophycin C and many others.

© 2007 Elsevier Ltd. All rights reserved.

Keywords: Dihydropyrans; δ -Hydroxy- α,β -unsaturated aldehydes; Allylation; Cyanation

The 2,6-disubstituted dihydropyran ring system is frequently found in various natural products, examples include swinholide,^{1,2} scytophycin C^{3,4} and laulimalide (Fig. 1).⁵

2,6-Disubstituted dihydropyrans are also synthetically useful intermediates in the preparation of polysubstituted tetrahydropyran ring systems, such as those found in the pseudomonic acids.⁶ As a result, several approaches^{7–13} have been reported for the preparation of dihydropyrans, and some of the more varied methodologies include electrophile-initiated alkylation of glycals,¹⁴ hetero-Diels–Alder cycloadditions,^{15–18} olefin metathesis,¹⁹ Prins-cyclizations of cyclopropyl carbinols,²⁰ or homoallylic alcohols,²¹ and an intramolecular silyl-modified Sakurai reaction (ISMS).^{22,23} Many of these reactions have limitations, such as the need for strictly anhydrous conditions, stoichiometric quantities of a Lewis acid initiator, or delivery of a strong Lewis acid at a low temperature. Recently, indium tribromide has received increasing attention as a water-tolerant, green Lewis acid catalyst for organic synthesis demonstrating highly chemo-, regio- and stereoselec-





Fig. 1. Structures of laulimalide and scytophycin C.

tive results.^{24,25} Compared to conventional Lewis acids, it has advantages of water stability, recyclability, operational simplicity, strong tolerance to oxygen and nitrogen-containing substrates and functional groups, and it can often be used in catalytic amounts.^{26–28} In continuation of our interest in the synthesis of scytophycin C,²⁹ we disclose

^{*} Corresponding author. Tel.: +91 40 27193535; fax: +91 40 27160512.
E-mail address: yadavpub@iict.res.in (J. S. Yadav).


herein a novel protocol for the construction of trans-2,6-disubstituted dihydropyrans.

Following our interest in the catalytic uses of indium tri-bromide,^{30–36} we describe an efficient method for the preparation of trans-2,6-disubstituted dihydropyrans through a tandem allylation/cyanation of δ -hydroxy- α,β -unsaturated aldehydes. To the best of our knowledge, no synthesis of C6-(allyl)- or C6-(cyano)-3, 6-dihydropyrans from δ -hydroxy- α,β -unsaturated aldehydes has been reported. In a model reaction, δ -hydroxy-enal **1** was treated with allyltrimethylsilane **2** using 5 mol % of indium(III) bromide as a catalyst. Interestingly, product 6-allyl-2-[2-(benzyloxy)ethyl]-3,6-dihydro-2H-pyran **3a** was isolated in 82% yield with trans-selectivity (Scheme 1).

The structure of product **3a** was established using NMR, the proton assignment was achieved with a QCOSY experiment. The presence of NOE's between the C-2 proton and the C-7 protons suggests that the stereochemistry at C-6 is *R* (a single enantiomer) and that the protons at C-2 and C-6 are trans to each other (Fig. 2).

No cis-diastereoisomer was observed in the ¹H NMR spectrum of the crude products obtained from the C-allylation of δ -hydroxy-enals. This result provided the incentive for a further study of reactions with various δ -hydroxy-enals. Interestingly, a diverse range of hydroxy-enals participated well in this reaction to afford the corresponding trans-2,6-disubstituted dihydropyrans in good yields (Table 1). Enantiomerically pure aldehydes gave optically pure products (Table 1, entries a–d). Trimethylsilyl cyanide also participated effectively in this reaction (Scheme 2).

However, *ortho*-hydroxy *trans*-cinnamaldehyde and allylsilane did not give the desired product under similar reaction conditions. Furthermore, α,β -unsaturated aldehydes without a δ -hydroxyl group failed to give the expected products, although, homoallylic alcohols were obtained in good yields. A possible reaction mechanism is illustrated in Scheme 3. The reaction proceeds with activation of aldehyde by indium(III) bromide and subsequent formation of an oxonium intermediate in which stereoelectronic and/or steric factors dictate the direction of the incoming nucleophile.



Scheme 1. A tandem allylation.





Fig. 2. Characteristic NOE's of product **3a**.

Table 1
InBr₃-catalyzed synthesis of trans-2,6-disubstituted 3,6-dihydro-2H-pyrans

Entry	Substrate 1	Product 3	Time (h)	Yield ^b (%)
a			1.0	82
b			0.5	85
c			0.5	88
d			0.5	70
e			2.0	85
f			2.0	70
g			1.5	85
h			1.0	75
i			1.5	70
j			2.0	70
k			1.0	82
l			1.0	72

^a The products were characterized ¹H NMR, IR and mass spectrometry.


^b Yield refers to pure products after chromatography.



Scheme 2. A tandem cyanation.

The scope and generality of this process is illustrated in Table 1.

Various indium(III) reagents such as InBr₃, InCl₃, In(OTf)₃ and In(ClO₄)₃ were screened for this transformation. Of these catalysts, indium tribromide was found to be



Scheme 3. A plausible reaction mechanism.

most effective in terms of conversion and selectivity. For example, treatment of aldehyde **1a** with allyltrimethylsilane in the presence of 5 mol % of InBr₃ and 5 mol % of InCl₃ for 1 h afforded **3a** in 82% and 70% yields, respectively. In addition, this method is useful for the direct synthesis of 2,3-dideoxy C-glycoside analogues (Table 1, entries c and d). This method facilitates the introduction of an allyl or cyano functionality on the pyran ring system in one-pot, making it an efficient pathway for producing 6-allyl- or 6-cyano-3,6-dihydro-2H-pyrans, which can be further modified to give biologically important natural products.

In summary, we have demonstrated an efficient protocol for the synthesis of trans-2,6-disubstituted dihydropyrans from δ -hydroxy- α,β -unsaturated aldehydes and allyltrimethylsilane in a highly regio- and stereoselective manner. We believe that this new method will find applications in the synthesis of biologically active natural products that contain a dihydropyran ring system.

General procedure: A mixture of the δ -hydroxy- α,β -unsaturated aldehyde (1 mmol), allyltrimethylsilane or trimethylsilyl cyanide (1.2 mmol) and indium tribromide (5 mol %) in dichloromethane (10 mL) was stirred at room temperature. After complete conversion, as indicated by TLC, the reaction mixture was diluted with water (10 mL) and extracted with dichloromethane (2 $\times$ 15 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and purified by column chromatography on silica gel (Merck, 100–200 mesh, ethyl acetate–hexane, 1:9) to afford pure dihydropyran.

Spectroscopic data for selected compounds: Compound **3a**: ¹H NMR (500 MHz, CDCl₃): δ 7.35–7.23 (m, 5H, Ar-H), 5.85 (m, 1H), 5.81 (m, 1H), 5.70 (m, 1H), 5.07 (m, 2H), 4.50 (2 $\times$ d, J = 12.0 Hz, 2H), 4.19 (m, 1H), 3.88 (tt, J = 4.4, 8.8 Hz, 1H), 3.62 (m, 1H), 3.59 (m, 1H), 2.40 (m, 1H), 2.24 (m, 1H), 2.05–1.74 (m, 4H). ¹³C NMR (75 MHz, CDCl₃): δ : 30.9, 35.7, 38.9, 64.9, 67.1, 72.5, 73.2, 116.9, 124.5, 127.7, 127.9, 128.5, 129.3, 135.3, 138.7. IR (KBr) Neat: ν_{max} : 3445, 2924, 2856, 1728, 1635, 1451, 1279, 1082, 760 cm⁻¹. ESI-HRMS [M+Na]⁺ found 281.1517, C₁₇H₂₂O₂Na requires 281.1508. Compound **3b**: ¹H NMR (300 MHz, CDCl₃): δ 7.34–7.22 (m, 5H, Ar-H), 5.89–5.64 (m, 3H), 5.10–5.03 (m, 2H), 4.48 (s, 2H), 4.18 (m, 1H), 3.56 (m, 1H), 3.42 (dd, J = 6.6, 9.0, 2H), 2.45–2.32 (m, 1H), 2.26–2.16 (m, 1H), 2.01–1.93 (m, 2H), 1.91–1.77 (dq, J = 4.4, 6.6 Hz, 1H) 0.98 (d, J = 6.6 Hz, 3H). ¹³C NMR (75 MHz, CDCl₃): δ 14.3, 29.9, 38.6, 38.8, 68.9, 72.9, 73.1, 116.8, 124.7, 125.0, 127.5, 127.7, 128.4, 129.4, 135.5, 139.0. IR (KBr) Neat: ν_{max} : 3448, 2924, 2854, 1638, 1456, 1215, 1082, 758 cm⁻¹. ESI-HRMS [M+Na]⁺ found 295.1673, C₁₈H₂₄O₂Na requires 295.1661. Compound **3c**: ¹H NMR (300 MHz, CDCl₃): δ 7.40–7.12 (m, 5H, Ar-H), 6.01–5.68 (m, 3H), 5.26–4.97 (m, 2H), 4.26 (m, 1H), 3.71 (tt, J = 4.0, 8.5 Hz, 1H), 2.88 (ddd, J = 5.2, 9.5, 13.7 Hz, 1H), 2.67 (ddd, J = 7.0, 9.5, 13.7 Hz, 1H), 2.46 (m, 1H), 2.35–2.21 (m, 1H), 2.12–1.95 (m, 2H), 1.95–1.86 (m, 1H), 1.86–1.70 (m, 1H). ¹³C

NMR (75 MHz, CDCl₃) δ 30.8, 32.2, 37.3, 39.1, 67.4, 72.4, 117.0, 124.4, 125.9, 128.5, 128.6, 129.4, 135.3, 142.5. IR (KBr) Neat: ν_{max} : 3446, 2925, 2855, 1637, 1457, 1216, 1082, 761 cm⁻¹. ESI-HRMS [M+Na]⁺ found 251.1411, C₁₆H₂₀ONa requires 251.1402.

Acknowledgement

V.S. thanks the CSIR, New Delhi, for the award of fellowship.

References and notes

- Hayakawa, H.; Miyashita, M. *Tetrahedron Lett.* **2000**, *41*, 707.
- Faulkner, D. J. *Nat. Prod. Rep.* **2000**, *17*, 7.
- Roush, W. R.; Dilley, G. J. *Synlett* **2001**, 955.
- Norcross, R. D.; Paterson, I. *Chem. Rev.* **1995**, *95*, 2041.
- Paterson, I.; De savi, C.; Tudge, M. *Org. Lett.* **2001**, *3*, 3149.
- Class, Y. J.; DeShong, P. *Chem. Rev.* **1995**, *95*, 1843.
- Morris, W. J.; Custar, D. W.; Scheidt, K. A. *Org. Lett.* **2005**, *7*, 1113.
- Lowe, J. T.; Panek, J. S. *Org. Lett.* **2005**, *7*, 1529.
- Evans, P. A.; Lawler, M. J. *Am. Chem. Soc.* **2004**, *126*, 8642.
- Viswanathan, G. S.; Yang, J.; Li, C. *Org. Lett.* **1999**, *1*, 993.
- Vares, L.; Rein, T. *J. Org. Chem.* **2002**, *67*, 7226.
- Brimble, M. A.; Pavia, G. S.; Stevenson, R. J. *Tetrahedron Lett.* **2002**, *43*, 1735.
- Semeyn, C.; Blaauiw, R. H.; Hiemstra, H.; Speckamp, W. N. *J. Org. Chem.* **1997**, *62*, 3426.
- Steinhuebel, D. P.; Fleming, J. J.; Du Bois, J. *Org. Lett.* **2002**, *4*, 293.
- Dosseter, A. G.; Jamison, T. F.; Jacobsen, E. N. *Angew. Chem., Int. Ed.* **1999**, *38*, 2398.
- Lubineau, A.; Auge, J.; Lubin, N. *Tetrahedron* **1993**, *49*, 4639.
- Danishefsky, S. J.; Selnick, H. G.; Zelle, R. E.; DeNinno, M. P. *J. Am. Chem. Soc.* **1988**, *110*, 4368.
- Danishefsky, S. J.; DeNinno, M. P. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 15.
- Wildemann, H.; Dunkelmann, P.; Muller, M.; Schmidt, B. *J. Org. Chem.* **2003**, *68*, 799.
- Yadav, V.; Kumar, N. V. *J. Am. Chem. Soc.* **2004**, *126*, 8652.
- Chan, K.-P.; Loh, T.-P. *Org. Lett.* **2005**, *7*, 4491.
- Marko, I. E.; Bayston, D. J. *Tetrahedron* **1994**, *50*, 7141.
- Marko, I. E.; Dobbs, A. P.; Scheirrmann, V.; Chelle, F.; Bayston, D. J. *Tetrahedron Lett.* **1997**, *34*, 2899.
- Zhang, Z.-H. *Synlett* **2005**, 711.
- Sakai, N.; Hirasawa, M.; Konakahara, T. *Tetrahedron Lett.* **2005**, *46*, 6407.
- Agnusdei, M.; Bandini, M.; Melloni, A.; Umani-Ronchi, A. *J. Org. Chem.* **2003**, *68*, 7126.
- Huang, J.-M.; Wong, C.-M.; Xu, F.-X.; Loh, T.-P. *Tetrahedron Lett.* **2007**, *48*, 3375.
- Harada, S.; Takita, R.; Ohshima, T.; Matsunaga, S.; Shibasaki, M. *Chem. Commun.* **2007**, 948.
- Yadav, J. S.; Ahmed, M. *Tetrahedron Lett.* **2002**, *43*, 7147.
- Yadav, J. S.; Reddy, B. V. S.; Rao, K. V.; Raj, K. S.; Prasad, A. R.; Kumar, S. K.; Kunwar, A. C.; Jayaprakash, P. J.; Jagannath, B. *Angew. Chem., Int. Ed.* **2003**, *42*, 5198.
- Yadav, J. S.; Reddy, B. V. S.; Kumar, G. M. *Synlett* **2001**, 1781.
- Yadav, J. S.; Reddy, B. V. S.; Baishya, G. *Synlett* **2003**, 396.
- Yadav, J. S.; Reddy, B. V. S.; Swamy, T. *Synthesis* **2004**, 106.
- Yadav, J. S.; Reddy, B. V. S.; Krishna, A. D.; Swamy, T. *Tetrahedron Lett.* **2003**, *44*, 6055.
- Yadav, J. S.; Reddy, B. V. S.; Gakul, B. *Green Chem.* **2003**, *5*, 264.
- Yadav, J. S.; Reddy, B. V. S.; Reddy, M. S.; Parimala, G. *Synthesis* **2003**, 2390.