

Medicinal Plants of Thailand. I Structures of Rheedeiosides A—D and *cis*-Entadamide A β -D-Glucopyranoside from the Seed Kernels of *Entada rheedei*

Sachiko SUGIMOTO, Katsuyoshi MATSUNAMI, and Hideaki OTSUKA*

Department of Pharmacognosy, Graduate School of Biomedical Sciences, Hiroshima University; 1–2–3 Kasumi, Minami-ku, Hiroshima 734–8553, Japan.

Received December 22, 2010; accepted February 1, 2011; published online February 3, 2011

Four new oleanane-type triterpene oligoglycosides, named rheedeiosides A, B, C and D, and one new thioamide glycoside, *cis*-entadamide A β -D-glucopyranoside, were isolated from the seed kernels of a Thai medicinal plant, *Entada rheedei* SPRENGEL. The rheedeiosides were found to contain *N*-acetylglucosamine as a sugar component. Their structures were elucidated based on spectral and chemical evidence.

Key words *Entada rheedei*; Fabaceae; rheedeioside; entadamide; Thai medicinal plant

Entada rheedei SPRENGEL is cultivated throughout the tropics. Its seeds have various medicinal uses including topical application as an ointment for the treatment of jaundice.¹⁾ Through phytochemical studies on *Entada* species, several triterpene glycosides have been isolated. The seed kernels of *E. rheedii*,¹⁾ and *E. pursora*,²⁾ and a bark of *E. phaseoloides*^{3,4)} were reported to contain a number of *N*-acetyl-D-glucosamine-containing entagenic acid saponins. Despite these investigations, the chemical constituents of *E. rheedei* as well as their biological activities have not yet been fully characterized. In this paper, we describe the isolation and structure elucidation of four new oleanane-type triterpene saponins, named rheedeiosides A—D, and a new thioamide glycoside, *cis*-entadamide A β -D-glucopyranoside, from the seed kernels of *E. rheedei*. The structures of the new compounds were elucidated on the basis of spectral and physicochemical evidence.

Isolation and Structural Elucidation Finely chopped seed kernels of *E. rheedei* were extracted with *n*-hexane and then the residue was extracted with MeOH. The MeOH was evaporated off under reduced pressure to yield a white residue. An aliquot of the residue was subjected to Diaion HP-20 column chromatography (CC) to give H₂O-, MeOH-, and acetone-eluted fractions. The MeOH-eluted fraction was then subjected to normal- and reversed-phase silica gel CC, and repetitive HPLC separation, which gave five new compounds, rheedeiosides A (**1**, 190 mg, 0.027%), B (**2**, 6.6 mg, 0.00093%), C (**3**, 22.1 mg, 0.0030%), and D (**4**, 44.5 mg, 0.0063%), and *cis*-entadamide A β -D-glucopyranoside (**5**, 4.4 mg, 0.0034%), together with two known compounds, entadamide A β -D-glucopyranoside⁵⁾ (**6**, 878 mg, 0.67%) and phenylpropanol β -D-glucopyranoside⁶⁾ (**7**, 9.2 mg, 0.0071%). Their structures were elucidated by extensive spectroscopic analyses, which yielded 1D- and 2D-NMR spectral data and electrospray ionization (ESI)-MS spectra, and by acid and alkaline hydrolyses.

Rheedeioside A (**1**) was isolated as an amorphous powder exhibiting negative optical rotation ($[\alpha]_D^{25}$ –27.1). The IR spectrum of **1** showed absorption bands at 3395, 1729, 1647 and 1078 cm^{–1}, ascribable to hydroxyl, ester carbonyl, amide carbonyl and ether functional groups. The molecular formula, C₇₂H₁₁₅NO₃₇, of **1** was determined by high-resolution (HR)-ESI-MS analysis. The ¹H-NMR spectrum of **1** showed

seven singlet methyl signals at δ_H 0.84, 0.87, 0.94, 1.07, 1.09, 1.10 and 1.74 (Table 1), and two fairly deshielded singlet methyl signals at δ_H 1.90 and 2.10 (Table 1). A typical signal for an axial proton attached to a hydroxylated carbon at δ_H 3.20 (dd, *J*=11, 4 Hz) (Table 1), and seven anomeric protons for sugar moieties [δ_H 4.75 (d, *J*=8 Hz), 4.98 (d, *J*=8 Hz), 5.16 (d, *J*=7 Hz), 5.32 (d, *J*=7 Hz), 5.40 (1H, d, *J*=8 Hz), 6.03 (d, *J*=7 Hz), and 6.11 (d, *J*=3 Hz)] were also observed (Table 1). The ¹³C-NMR spectral data indicated the presence of one terminal apiofuranose and three terminal xylopyranoses, and the aglycone moiety showed good resemblance to those of entagenic acid (**8**) isolated from *Entada phaseoloides* (Table 2).^{3,4)} Judging from the downfield shift of C-3 (δ_C 89.6) and the upfield shift value observed for C-28 (δ_C 175.8) (Table 2), **1** was presumed to be a bisdesmosidic glycoside of entagenic acid with sugar linkages at C-3 through a glycosidic bond and at C-28 through an ester bond. Alkaline hydrolysis of **1** liberated a monodesmoside (**1a**) and a triose, whose sugar composition was analyzed to be D-apiose, D-xylose and D-glucose. The NMR spectral data of **1a** showed that the monodesmoside (**1a**) possessed two terminal xylopyranoses and two inner sugars (Table 2). Acid hydrolysis of **1a** gave entagenic acid as an aglycone (**1b** = **8**), and sugar analysis of the hydrolyzate revealed the presence of D-xylose, D-glucose and D-glucosamine hydrochloride. The presence of an *N*-acetylglucosamine group was substantiated by the ¹H-NMR at δ_H 2.10 (3H, s) and the D₂O exchangeable proton at δ_H 8.91 (d, *J*=8 Hz, NH), and the ¹³C-NMR resonances at δ_C 23.5 and 170.4, together with the heteronuclear single quantum correlation (HSQC) spectroscopic data for C-2' of the hexopyranose moiety (δ_H 4.40 on δ_C 57.4), confirmed the presence of the 2-(acetylamino)-2-deoxyglucopyranose (GlcNAc) unit.⁷⁾ The connectivities of the sugar chains were confirmed by heteronuclear multiple bond correlation (HMBC) experiments (Fig. 1). The cross peak between δ_H 4.98 (GlcNAc-H-1') and δ_C 89.5 (C-3) showed that the GlcNAc moiety was linked with the hydroxy group at C-3 of the aglycone. Further correlation was also observed between H-1'' (δ_H 5.40, *J*=8 Hz) of xylopyranose and C-3' (δ_C 80.7) of GlcNAc, indicating the location of it at C-3'. The HMBC correlations between H-1''' [δ_H 5.16 (d, *J*=7 Hz)] of glucopyranose and C-6' (δ_C 68.1) of GlcNAc, and between H-1'''' [δ_H 4.75 (d, *J*=8 Hz)] of xylopyranose and C-3''' (δ_C 83.2) of glu-

* To whom correspondence should be addressed. e-mail: hotsuka@hiroshima-u.ac.jp

Table 1. ¹H-NMR Spectral Data for Rheedeiosides A–D (1–4) (600 MHz, Pyridine-*d*₅) δ in ppm and *J* in Hz

H	1	2 ^{a)}	3	4
1	0.89 m 1.47 br d 13	0.85 m 1.45 m	0.92 m 1.47 br d 13	1.10 m 1.64 br d 13
2	1.70 br d 13 2.07 m	1.67 m 2.10 m	1.70 br d 13 2.08 m	1.84 br d 13 2.24 dd 13, 4
3	3.20 dd 11, 4	3.24 dd 11, 4	3.22 dd 12, 4	3.35 dd 13, 4
5	0.77 d 12	0.79 m	0.78 d 12	0.86 d 12
6	1.27 br d 12 1.50 br dd 12, 2	1.25 m 1.49 m	1.28 br d 11 1.50 br dd 11, 3	1.32 br dd 12, 3 1.53 br dd 12, 3
7	1.93 m 2.12 m	1.96 m 2.13 m	1.95 m 2.13 br d 10	2.00 m 2.16 br d 11
9	1.68 d 9	1.73 m	1.71 d 9	1.76 d 10
11	1.94 2H m	1.99 2H m	1.97 2H m	2.02 2H m
12	5.62 br s	5.64 br s	5.64 br s	5.62 br s
15	4.47 d 3	4.50 m	4.50 d 3	4.52 m
16	4.95 br s	5.01 m	4.97 m	4.99 m
18	3.41 br d 11	3.43 m	3.39 dd 13, 5	3.40 dd 13, 5
19	1.26 m 2.69 dd 13, 13	1.32 m 2.74 m	1.27 dd 13, 5 2.72 dd 13, 13	1.30 dd 13, 5 2.72 dd 13, 13
21	1.20 dd 12, 2 2.34 dd 12, 5	1.13 m 2.39 m	1.23 br d 12 2.37 dd 12, 4	1.22 m 2.38 dd 13, 5
22	2.16 br d 12 2.28 dd 12, 5	2.20 m 2.24 m	2.17 br d 13 2.33 dd 13, 5	2.19 br d 13 2.31 dd 14, 4
23	1.10 s	1.13 s	1.09 s	1.14 s
24	0.87 s	0.89 s	0.88 s	0.94 s
25	0.84 s	0.76 s	0.84 s	0.89 s
26	1.07 s	1.11 s	1.10 s	1.12 s
27	1.74 s	1.79 s	1.79 s	1.78 s
29	0.94 s	0.99 s	0.96 s	0.97 s
30	1.09 s	1.08 s	1.11 s	1.11 s
1'	4.98 d 8	5.04 d 7	5.02 d 8	5.00 d 8
2'	4.40 dd 9, 8	4.41 m	4.35 dd 9, 8	4.51 m
3'	4.72 m	4.77 m	4.74 dd 9, 8	4.48 m
4'	4.06 m	4.10 m	4.07 m	4.35 m
5'	3.96 m	4.02 m	4.00 m	3.93 m
6'	4.55 dd 11, 3 4.65 dd 11, 5	4.62 m 4.72 m	4.59 dd 11, 3 4.70 dd 11, 5	4.57 dd 11, 2 4.62 dd 11, 5
1''	5.40 d 8	5.48 d 8	5.43 d 8	
2''	3.89 m	3.86 m	3.94 m	
3''	4.23 m	4.23 m	4.21 m	
4''	3.94 m	4.00 m	4.08 m	
5''	3.53 br d 11 4.33 dd 11, 5	3.55 br d 12 4.66 m	3.65 dd 11, 2 4.66 dd 11, 6	
1'''	5.16 d 7	5.23 d 8	5.19 d 7	5.10 d 5
2'''	4.37 m	4.40 m	4.44 m	4.47 m
3'''	4.38 m	4.48 m	4.46 m	4.69 m
4'''	4.17 m	4.21 m	4.21 m	4.35 m
5'''	4.14 m	4.28 m	4.20 m	3.70 dd 11, 2 4.24 dd 11, 5
6'''	4.58 dd 11, 3 4.66 dd 11, 5	4.27 m 4.43 m	4.62 m 4.69 dd 10, 5	
1''''	4.75 d 8	4.84 d 8	4.78 d 8	4.96 d 7
2''''	3.95 m	4.05 m	3.98 m	3.98 m
3''''	4.20 m	4.21 m	4.25 m	4.17 m
4''''	4.16 m	4.20 m	4.09 m	3.97 m
5''''	3.40 br d 11 4.59 dd 11, 5	3.48 br d 13 4.30 m	3.41 br d 11 4.27 dd 11, 4	3.54 dd 12, 2 4.08 dd 12, 5
1'''''	6.03 d 8	6.15 d 7	6.06 d 8	6.07 d 8
2'''''	4.15 m	4.22 m	4.20 m	4.22 m
3'''''	4.04 m	4.06 m	4.04 m	4.08 m
4'''''	4.14 m	4.16 m	4.19 m	4.09 m
5'''''	3.86 m	3.92 m	4.21 m	4.00 m
6'''''	4.21 dd 11, 5 4.42 m	4.33 m 4.45 m	4.24 dd 12, 4 4.44 dd 12, 5	4.64 dd 12, 2 4.69 dd 12, 6
1''''''	5.32 d 7	5.38 d 7	5.40 d 7	5.35 d 8
2''''''	3.99 m	4.08 m	4.06 m	4.02 m
3''''''	4.20 m	4.19 m	4.23 m	4.17 m
4''''''	4.24 m	4.10 m	4.12 m	4.15 m
5''''''	3.60 br d 11 4.17 dd 11, 5	3.62 br d 13 4.18 m	3.60 br d 10 4.18 m	3.60 dd 12, 2 4.18 dd 12, 5
1'''''''	6.11 d 3	6.17 d 3	5.16 d 8	6.16 d 3
2'''''''	4.70 d 3	4.68 d 3	3.96 m	4.74 d 3
3'''''''	—	—	4.10 m	—
4'''''''	4.68 d 10 4.26 d 10	4.74 d 10 4.30 d 10	3.94 m	4.69 d 10 4.29 d 10
5'''''''	4.12 d 13 4.09 d 13	4.12 2H br s	3.63 dd 11, 2 4.38 dd 11, 4	4.13 d 13 4.11 d 13
NHCOCH ₃	2.10 s	2.11 s	2.11 s	2.10 s
NHCOCH ₃	8.91 d 8	8.86 d 10	8.90 d 9	8.74 d 7
OCOCH ₃	1.90 s	—	—	1.92 s

a) At 500 MHz. m: multiplet or overlapped signals.

copyranose indicated the location of them at C-6' and C-3''', respectively. In addition, the correlation observed between H-1'''' [δ_H 6.03 (d, *J*=8 Hz)] of glucopyranose and the signal of the carbon at δ_C 175.8 (C-28) in the HMBC spectrum confirmed the ester linkage at C-28. Further correlations observed in the HMBC spectrum between the signal at H-1 [δ_H 5.32 (d, *J*=7 Hz)] of xylopyranose and δ_C 79.9 (C-2'''), and between H-1'''''' [δ_H 6.11 (d, *J*=3 Hz)] of apiofuranose and C-3 (δ_C 85.0) of glucopyranose established the sugar linkage of the ester moiety. The position of acetyl group was determined to be at the hydroxy group at C-6'', as judged from the significant downfield shift of the methylene protons (H₂-6''', δ_H 4.58, 4.66), and the HMBC correlation peaks between H₂-6''' and the carbonyl carbon (δ_C 170.7) of acetyl group. Acetylation of **1** with acetic anhydride, pyridine and *N,N*-dimethylaminopyridine (DMAP) gave a nonadecaacetate (**1c**), and protons assignable to H-3', H-6', H-3''', H-2'''' and H-3'''''' remained intact in the ¹H-NMR spectrum. The ¹H-¹H correlation spectroscopy (COSY) and HMBC spectral data for **1c** also supported ramification of the sugar chains. The mode of linkage of all sugars was determined to be β from the coupling constants of their anomeric protons. Thus, the structure of **1** was elucidated to be entagenic acid 3-*O*-β-D-xylopyranosyl-(1→3)-6-*O*-acetyl-β-D-glucopyranosyl(1→6)-*O*-[β-D-xylopyranosyl(1→3)]-2-acetyl-amino-2-deoxy-β-D-glucopyranoside 28-*O*-[β-D-xylopyranosyl(1→2), β-D-apiofuranosyl(1→3)]-β-D-glucopyranosyl ester (rheedeioside A), as shown in Chart 1.

Rheedeioside B (**2**) was also isolated as an amorphous powder exhibiting negative optical rotation ([α]_D²³ -28.5). The IR spectrum of **2** showed absorption bands at 3395 and 1077 cm⁻¹, suggestive of a glycosidic structure, together with absorption bands at 1647 and 1559 cm⁻¹ due to an amide function. NMR spectral data were essentially the same as those of **1**, except for the absence of an acetyl group. The elemental composition (C₇₀H₁₁₃NO₃₆), determined by HR-ESI-MS, was also 42 mass units less (CH₂CO) from that of **1**. Mild alkaline hydrolysis of **1** gave deacetylated **1**, whose spectroscopic data were identical with those of **2**. Consequently, the structure of **2** was elucidated to be entagenic acid 3-*O*-β-D-xylopyranosyl-(1→3)-β-D-glucopyranosyl(1→6)-*O*-[β-D-xylopyranosyl(1→3)]-2-acetyl-amino-2-deoxy-β-D-glucopyranoside 28-*O*-[β-D-xylopyranosyl(1→2), β-D-apiofuranosyl(1→3)]-β-D-glucopyranosyl ester, as shown in Chart 1.

Rheedeioside C (**3**) was also obtained as an amorphous powder exhibiting negative optical rotation ([α]_D²³ -20.5). The molecular formula, C₇₂H₁₁₅NO₃₇, of **3** was determined by positive-ion HR-ESI-MS. Its ¹H- and ¹³C-NMR spectra were similar to those of **1**, exhibiting seven singlet methyls, and seven anomeric proton and carbon signals. The ¹³C-NMR spectrum exhibited 30 signals for the aglycone of bis-desmoisde, and four terminal xylopyranoses. On acid hydrolysis of **3**, the presence of D-xylose, D-glucose and D-glucosamine was confirmed. Thus, the apiofuranose found in **1** was replaced by xylopyranose. The HMBC experiment showed that there was a long-range correlation between H-1'''''' (δ_H 5.16) and C-3'''''. On the basis of the above mentioned evidence, the structure of **3** was elucidated to be entagenic acid 3-*O*-β-D-xylopyranosyl-(1→3)-6-acetyl-β-D-glucopyranosyl-(1→6)-*O*-[β-D-xylopyranosyl(1→3)]-2-acetyl-amino-2-deoxy-β-D-glucopyranoside, 28-*O*-β-D-xylopyr-

Table 2. ^{13}C -NMR Spectroscopic Data for Rheedeiosides A—D (**1**—**4**), Entagenic Acid (**8**), and **1a** (150 MHz, Pyridine- d_5)

C	1	1a	2^{a)}	3	4	8^{b)}	C	1	1a	2^{a)}	3	4
1	38.8	38.8	38.9	39.1	39.0	39.5	1'	104.3	104.3	104.4	104.32	104.8
2	26.3	26.4	26.5	26.4	26.6	28.3	2'	57.4	57.6	57.7	57.6	57.8
3	89.5	89.5	89.6	89.5	89.1	78.6	3'	80.7	80.8	81.0	80.8	75.3
4	39.0	39.1	39.2	38.9	39.2	39.4	4'	73.6	73.0	73.3	73.7	72.5
5	55.6	55.7	55.8	55.7	55.7	56.0	5'	76.2	76.3	76.5	76.4	76.0
6	18.8	18.8	18.8	18.8	18.8	19.4	6'	67.9	68.0	68.1	68.0	69.44
7	36.9	36.9	37.1	37.1	37.0	37.2	1''	104.2	104.3	104.5	104.34	
8	41.3	41.2	41.5	41.1	41.5	41.6	2''	74.7	74.5	74.79	75.0	
9	47.6	47.3	47.4	47.4	47.4	47.8	3''	77.9	77.7	78.0	78.3	
10	37.0	37.1	37.0	37.0	37.1	37.9	4''	70.2	70.7	70.9	70.80	
11	23.8	23.9	24.0	23.9	24.0	34.2	5''	66.1	66.1	66.2	66.2	
12	124.8	124.0	124.8	124.9	124.9	124.3	1'''	103.2	103.2	103.4	103.3	102.2
13	144.4	145.4	144.7	144.4	144.5	145.8	2'''	73.0	73.5	73.8	73.1	80.1
14	47.3	47.8	47.3	47.7	47.7	48.2	3'''	82.9	83.0	83.2	83.1	72.9
15	68.1	68.3	68.3	68.2	68.2	68.6	4'''	68.6	68.6	68.6	68.7	67.4
16	78.7	78.0	79.0	78.8	78.8	79.7	5'''	77.7	79.0	78.8	77.8	64.2
17	48.5	48.4	48.6	48.6	48.6	48.8	6'''	64.0	62.3	62.4	64.1	
18	41.7	42.0	41.8	41.8	41.8	42.4	1''''	107.4	107.5	107.7	107.6	106.2
19	46.3	46.7	46.6	46.4	46.4	47.1	2''''	75.6	74.8	75.1	75.7	75.8
20	30.7	30.9	30.8	30.8	30.8	31.0	3''''	78.1	78.3	78.7	78.4	78.3
21	35.8	36.1	36.0	35.9	35.9	36.4	4''''	70.7	71.4	70.7	70.83	70.4
22	31.8	32.8	32.1	31.9	31.9	32.8	5''''	67.2	67.3	67.5	67.3	67.2
23	28.0	28.0	28.1	28.1	28.1	28.9	1'''''	93.1		93.5	93.2	93.2
24	16.9	17.0	17.0	17.0	17.0	16.5	2'''''	79.9		80.6	80.1	80.3
25	15.6	15.5	15.7	15.7	15.8	15.9	3'''''	85.0		85.2	87.2	85.0
26	17.9	18.0	18.1	18.0	18.0	18.3	4'''''	71.4		71.6	71.5	70.8
27	20.8	20.8	20.8	20.6	20.7	20.7	5'''''	78.2		78.5	78.3	77.8
28	175.8	179.6	175.8	175.9	175.9	180.0	6'''''	62.1		62.2	62.3	64.1
29	33.1	33.3	33.3	33.2	33.2	33.4	1''''''	105.4		105.7	105.3	105.6
30	24.4	24.5	24.5	24.5	24.5	25.0	2''''''	74.5		74.77	74.5	74.9
							3''''''	78.1		78.1	78.0	78.3
							4''''''	69.2		69.4	69.0	69.36
							5''''''	66.9		67.1	67.2	67.0
							1''''''	111.1		111.5	106.0	111.3
							2''''''	77.6		77.8	74.7	77.7
							3''''''	80.2		80.4	78.0	80.4
							4''''''	74.9		75.2	70.3	75.1
							5''''''	65.2		65.6	66.9	65.4
							NHCOCH ₃	23.5	23.4	23.7	23.6	23.6
							NHCOCH ₃	170.4	170.2	169.9	170.1	170.3
							OCOCH ₃	20.5			20.7	20.6
							OCOCH ₃	170.7			170.7	170.6

a) At 125 MHz. b) Data taken from ref. 4.

anosyl-(1→2)-[β -D-xylopyranosyl-(1→3)]- β -D-glucopyranosyl ester.

Rheedeioside D (**4**) was also obtained as an amorphous powder with negative optical rotation ($[\alpha]_{\text{D}}^{25} -24.0$). The IR spectrum of **4** showed absorption bands at 3388, 1730, 1648, and 1079 cm^{-1} , ascribable to hydroxy, ester carbonyl, amide and ether functional groups. The molecular formula, $\text{C}_{66}\text{H}_{105}\text{NO}_{32}$, of **4** was determined by positive-ion HR-ESI-MS. Acid hydrolysis of **4** liberated entagenic acid together with D-apiose, D-glucose, D-glucosamine, D-xylose and L-arabinose, which were identified by HPLC analysis. The ^1H - and ^{13}C -NMR spectra of **4** showed signals assignable to one terminal apiofuranoside [δ_{H} 6.16 (d, $J=3$ Hz)], two terminal xylopyranosides [δ_{H} 4.96 (d, $J=7$ Hz) and 5.35 (d, $J=8$ Hz)], and GlcNAc [H-1': δ_{H} 5.00 (d, $J=8$ Hz), C-2': δ_{C} 57.8, and NHCOCH_3 : δ_{C} 23.6 and 170.3 and δ_{H} 2.10 (3H) and 8.74 (D_2O exchangeable)], together with one acetyl group. The remaining anomeric proton resonating at δ_{H} 5.10 (d, $J=5$ Hz) was assigned as that of arabinopyranose, and that resonating at 6.07 (d, $J=8$ Hz) as that of ester linked glucopyranose. In

the HMBC spectrum, the anomeric proton of GlcNAc, showed a correlation peak with with C-3, the anomeric proton of arabinopyranose with C-6' (δ_{C} 69.44) of GlcNAc and the anomeric proton of one of the xylopyranoses (δ_{H} 4.96) with C-2''' (δ_{C} 80.1), thus establishing the sugar linkage was at the C-3 position. The ^{13}C -NMR spectral data for the ester linked oligosaccharide were superimposable on those of **1**—**3**, except for the downfield shift of the H_2 -6''' protons, as well as H_2 -6''' protons showing a correlation peak with the acetyl carbonyl carbon (δ_{C} 170.6). Consequently, the structure of **4** was determined to be entagenic acid 3-O- β -D-xylopyranosyl-(1→2)- α -L-arabinopyranosyl-(1→6)-2-acetylamino-2-deoxy- β -D-glucopyranoside, 28-O- β -D-xylopyranosyl-(1→2)-[β -D-apiofuranosyl-(1→3)]-6-acetyl- β -D-glucopyranosyl ester.

Compound **5** was obtained as an amorphous powder with negative optical rotation ($[\alpha]_{\text{D}}^{24} -12.0$ in MeOH). The molecular formula, $\text{C}_{12}\text{H}_{21}\text{NO}_7\text{S}$, of **5** was determined by positive-ion HR-ESI-MS. The IR spectrum showed absorption bands at 3335 cm^{-1} (br, OH and NH), 1638 cm^{-1} (CO-N),




Fig. 1. Significant ^1H – ^1H and HMBC Correlations of Rheedeioside A (1)

Chart 1.

and 1576 cm^{-1} ($\text{C}=\text{C}$). In the UV spectrum of **5**, an absorption maximum was observed at 266 ($\log \epsilon$ 4.87) nm. This compound exhibited UV, IR, ^1H - and ^{13}C -NMR spectral data similar to those of entadamide A- β -D-glucopyranoside (**6**),⁵ which co-occurred in this plant, except for the coupling constant of the olefinic bond. The ^1H -NMR spectrum indicated

Fig. 2. Significant ^1H – ^1H and HMBC Correlations of *cis*-Entadamide A β -D-Glucopyranoside (**5**)

the geometry of the disubstituted double bond of **5** to be in a *cisoid* form [δ_{H} 6.93 and 5.89 (each 1H, both d, $J=10$ Hz)], and then the structure of **5** was determined to be *cis*-entadamide A β -D-glucopyranoside (**5**).

In conclusion, four oleanane-type triterpene bisbesmosides, rheedeiosides A (**1**), B (**2**), C (**3**), and D (**4**), and a thioamide glycoside (**5**), together with two known compounds, were isolated from the seed kernels of *E. rheedei* collected in Thailand.

Experimental

The following instruments were used to obtain physical and spectroscopic data: mp, Yanagimoto micromelting point apparatus (uncorr.); optical rotations, JASCO P-1030 digital polarimeter; IR spectra, Shimadzu FT-710 spectrophotometer; HR-ESI mass spectra, LTQ Orbitrap XL; ^1H -NMR spectra, JEOL JNM-LA500 (500 MHz) and JEOL ECA-600K (600 MHz) spectrometers; ^{13}C -NMR spectra, JNM-LA500 (125 MHz) and JEOL ECA-600K (150 MHz) spectrometers with tetramethylsilane as internal standard and run at 35°C ; and HPLC detector, Shimadzu RID-6A refractive index detector. Highly-porous synthetic resin Diaion HP-20 was purchased from Mitsubishi Chemical Co., Ltd. (Tokyo, Japan). Silica gel CC was performed on silica gel 60 [E. Merck, Darmstadt, Germany, 70–230 mesh]. Reversed-phase octadecyl silanized (ODS) open CC (RPCC) was performed on Cosmosil 75C₁₈-OPN (Nacalai Tesque, Kyoto, Japan) [$\Phi=50$ mm, $L=25$ cm, linear gradient: MeOH–H₂O]. HPLC was performed on an ODS-3 column (Inertsil; GL Science, Tokyo, Japan; $\Phi=10$ mm, $L=25$ cm), and the eluate was monitored with a refractive index monitor. Precoated silica gel 60 F₂₅₄ plates (E. Merck; 0.25 mm in thickness) were used for TLC analyses, with visualization by spraying of a 10% solution of H₂SO₄ in ethanol and heated at around 150°C on a hotplate. Authentic D-apiose [$[\alpha]_{\text{D}}^{25} +9.4^\circ$ ($c=0.84$, H₂O)] was obtained by chromatographic separation of the hydrolyzate of apiin, isolated from commercial parsley (*Petroselinum crispum*). D-Apiose was identified by NMR spectroscopy.⁸⁾

Plant Material The seed kernels of *Entada rheedei* were purchased at a Thai market in 2009 and identified by Prof. Sorasak Lhieochaiphant of Faculty of Pharmacy, Chiang Mai University, Thailand.

Isolation of Compounds 1–7 The seed kernels of *E. rheedei* (1.86 kg) were finely chopped and extracted with hexane (41) for three days at room temperature. Evaporation of the solvent under reduced pressure provided a hexane-soluble extract (102 g, 5.49%). The dried-up residue was extracted three times with MeOH (41) at room temperature for 3 d each and then evaporated, providing a MeOH-soluble extract (248 g, 13.4%). The MeOH-soluble extract (133 g) was subjected to a Diaion HP-20 CC [2.0 kg, H₂O (151)→MeOH (151)→acetone (101)] to give H₂O (98.1 g, 9.86%), MeOH (30.0 g, 3.01%) and acetone (3.2 g, 0.32%) eluates.

The MeOH eluate (29.0 g) was subjected to normal-phase silica gel CC (900 g) {CHCl₃ (41)→CHCl₃:MeOH [9:1 (61)→4:1 (61)→7:3 (61)→3:2 (61)]→CHCl₃:MeOH:H₂O (6:4:1) (201)→MeOH (61)} to give ten fractions [Fr. 1 (274 mg), Fr. 2 (888 mg), Fr. 3 (6.1 g), Fr. 4 (859 mg), Fr. 5 (791 mg), Fr. 6 (871 mg), Fr. 7 (1.7 g), Fr. 8 (3.5 g), Fr. 9 (11.3 g), and Fr. 10 (908 mg)]. Fraction 3 (5.3 g) was separated by RPCC [120 g, MeOH:H₂O (1:9→3:7→1:1→7:3→MeOH)] to yield 11 fractions [Fr. 3-1 (37.9 mg), Fr. 3-2 (4.9 g), Fr. 3-3 (67.4 mg), Fr. 3-4 (9.6 mg), Fr. 3-5 (9.6 mg), Fr. 3-6 (20.1 mg), Fr. 3-7 (10.2 mg), Fr. 3-8 (51.7 mg), Fr. 3-9 (1.2 mg), Fr. 3-10 (20.6 mg), and Fr. 3-11 (8.4 mg)]. Fraction 3-2 (880 mg) was purified by

HPLC [MeOH–H₂O (1:4, v/v)] to give *cis*-entadamide A β -D-glucopyranoside (**5**, 4.4 mg), entadamide A β -D-glucopyranoside (**6**, 878 mg), and phenylpropanol β -D-glucopyranoside (**7**, 9.2 mg).

Fr. 5 (791 mg), Fr. 6 (871 mg), and Fr. 7 (1.70 g) were combined and subjected to RPCC [120 g, MeOH:H₂O (1:9→3:7→1:1→7:3→9:1→MeOH)] to give 22 fractions [Fr. 5,6,7-1 (527 mg), Fr. 5,6,7-2 (282 mg), Fr. 5,6,7-3 (789 mg), Fr. 5,6,7-4 (254 mg), Fr. 5,6,7-5 (53.0 mg), Fr. 5,6,7-6 (21.2 mg), Fr. 5,6,7-7 (103 mg), Fr. 5,6,7-8 (43.7 mg), Fr. 5,6,7-9 (17.6 mg), Fr. 5,6,7-10 (22.1 mg), Fr. 5,6,7-11 (8.0 mg), Fr. 5,6,7-12 (44.8 mg), Fr. 5,6,7-13 (20.0 mg), Fr. 5,6,7-14 (14.6 mg), Fr. 5,6,7-15 (16.6 mg), Fr. 5,6,7-16 (6.0 mg), Fr. 5,6,7-17 (24.4 mg), Fr. 5,6,7-18 (478 mg), Fr. 5,6,7-19 (78.6 mg), Fr. 5,6,7-20 (55.0 mg), Fr. 5,6,7-21 (20.0 mg), and Fr. 5,6,7-22 (800 mg)]. Fr. 5,6,7-18 (450 mg) was separated by HPLC [MeOH–acetonitrile–H₂O (3:2:5, v/v/v)] to give rheedeiosides A (**1**, 190 mg), B (**2**, 6.6 mg), C (**3**, 44.5 mg), and D (**4**, 21.2 mg).

The known compounds were identified by comparison of their physical data ($[\alpha]_D$, IR, ¹H- and ¹³C-NMR, and MS) with the reported values.

Rheedeioside A (1): Amorphous powder; $[\alpha]_D^{24}$ –27.1 (*c*=2.41, MeOH); IR (film) $\nu_{\max}$ 3395, 2944, 1729, 1647, 1559, 1078, 1048 cm^{–1}; ¹H-NMR (600 MHz, pyridine-*d*₅): Table 1; ¹³C-NMR (150 MHz, pyridine-*d*₅): Table 2; HR-ESI-MS (positive-ion mode): *m/z* 1608.6993 [M+Na]⁺ (Calcd for C₇₂H₁₁₅NO₃₇Na: 1608.7040).

Rheedeioside B (2): Amorphous powder; $[\alpha]_D^{23}$ –28.5 (*c*=0.61, MeOH); IR (film) $\nu_{\max}$ 3395, 2941, 1730, 1647, 1560, 1077, 1049 cm^{–1}; ¹H-NMR (500 MHz, pyridine-*d*₅): Table 1; ¹³C-NMR (125 MHz, pyridine-*d*₅): Table 2; HR-ESI-MS (positive-ion mode): *m/z* 1566.6893 [M+Na]⁺ (Calcd for C₇₀H₁₁₃NO₃₆Na: 1566.6935).

Rheedeioside C (3): Amorphous powder; $[\alpha]_D^{23}$ –20.5 (*c*=4.30, MeOH); IR (film) $\nu_{\max}$ 3395, 2941, 1730, 1647, 1560, 1077, 1049 cm^{–1}; ¹H-NMR (600 MHz, pyridine-*d*₅): Table 1; ¹³C-NMR (150 MHz, pyridine-*d*₅): Table 2; HR-ESI-MS (positive-ion mode): *m/z* 1608.7003 [M+Na]⁺ (Calcd for C₇₂H₁₁₅NO₃₇Na: 1608.7040).

Rheedeioside D (4): Amorphous powder; $[\alpha]_D^{25}$ –24.0 (*c*=2.10, MeOH); IR (film) $\nu_{\max}$ 3388, 2945, 1730, 1648, 1559, 1079, 1047 cm^{–1}; ¹H-NMR (600 MHz, pyridine-*d*₅): Tables 2 and 3; ¹³C-NMR (150 MHz, pyridine-*d*₅): Table 1; positive-ion HR-ESI-MS: *m/z* 1446.6471 [M+Na]⁺ (Calcd for C₆₆H₁₀₅NO₃₂Na: 1446.6512).

***cis*-Entadamide A β -D-Glucopyranoside (5):** Amorphous powder; $[\alpha]_D^{23}$ –12.0 (*c*=0.44, MeOH); IR (film) $\nu_{\max}$ 3335, 2945, 1748, 1637, 1575, 1074 cm^{–1}; ¹H-NMR (500 MHz, CD₃OD): Table 3; ¹³C-NMR (125 MHz, CD₃OD): Table 3; HR-ESI-MS (positive-ion mode): *m/z* 346.0927 [M+Na]⁺ (Calcd for C₁₂H₂₁NO₅Na: 346.0931).

Alkaline Hydrolysis of Rheedeioside A (1) A solution of rheedeioside A (**1**) (38.3 mg) in 50% aqueous 1,4-dioxane (1.0 ml) was treated with 10% aqueous KOH (1.0 ml) and stirred at 80 °C for 3 h. The reaction mixture was neutralized with Dowex HCR W2 (H⁺ form) and then the resin was removed by filtration. The residue of the reaction mixture was subjected to RPCC [Cosmosil (0.5 g), H₂O–MeOH (100:0→4:1→2:3→0:100, v/v)] and on evaporation of the 60% MeOH eluate, entagenic acid monodesmoside (**1a**, 25.2 mg) was obtained. Entagenic acid monodesmoside (**1a**): Amorphous powder; $[\alpha]_D^{23}$ –13.6 (*c*=0.13, MeOH); IR (film) $\nu_{\max}$ 3364, 2942, 1718, 1635, 1569, 1050 cm^{–1}; ¹H-NMR (600 MHz, pyridine-*d*₅): δ : 0.78, 0.90, 1.02, 1.09, 1.13, 1.14, 1.81, 2.11 (3H each, all s, H₃-25, 24, 29, 26, 23, 30, 27 and NHCOC(=O)CH₃), 3.22 (1H, dd, *J*=12, 4 Hz, H-3), 4.79 (1H, d, *J*=8 Hz, H-1'''), 5.03 (1H, d, *J*=8 Hz, H-1'), 5.20 (1H, d, *J*=7 Hz, H-1''), 5.43 (1H, d, *J*=8 Hz, H-1''), 5.67 (1H, brs, H-12); ¹³C-NMR (150 MHz, pyridine-*d*₅): Table 1; HR-ESI-MS (positive-ion mode): *m/z* 1140.5555 [M+Na]⁺ (Calcd for C₅₄H₈₇NO₂₃Na: 1140.5561). The residue of the 100% H₂O eluate was heated in 2 M HCl (1.0 ml) under reflux for 3 h. After cooling, the reaction mixture was poured into ice-water and neutralized with Amberlite IRA-400 (OH[–] form), and then the resin was removed by filtration. The filtrate was extracted with EtOAc and the aqueous layer was subjected to HPLC analysis [column: Shodex Asahipak NH 2P-50 4E, Φ =4.6 mm, *L*=25 cm; mobile phase: MeCN–H₂O (4:1, v/v); detection: optical rotation detector (JASCO 2090_{Plus}); and flow rate: 1.0 ml/min] to detect D-apiose, D-xylose and D-glucose, which were identified by comparison of their retention times with those of authentic samples, D-apiose (*t_R* 4.5 min, positive optical rotation), D-xylose (*t_R*: 7.5 min, positive optical rotation), and D-glucose (*t_R*: 10.3 min, positive optical rotation), respectively.

Acid Hydrolysis of Rheedeioside B Monodesmoside (1a) A solution of **1a** (12.5 mg) in 2 M HCl (1.0 ml) was heated under reflux for 3 h. After cooling, the reaction mixture was poured into ice-water and neutralized with Amberlite IRA-400 (OH[–] form), and then the resin was removed by filtration. The filtrate was extracted with EtOAc and the organic layer was evapo-

Table 3. NMR Spectroscopic Data for Compound **5** (C: 125 MHz and H: 500 MHz, CD₃OD), δ in ppm and *J* in Hz

	C	H
1	169.1	—
2	115.9	5.89 d, 10
3	148.8	6.93 d, 10
1'	40.4	3.40 2H, t, 7
2'	69.9	3.62 2H, t, 7
S–CH ₃	19.2	2.32 3H, s
1''	104.6	4.26 d, 8
2''	75.1	3.19 dd, 8, 8
3''	77.9	3.34 dd, 8, 8
4''	71.7	3.28 dd, 8, 8
5''	78.0	3.26 ddd, 8, 5, 2
6''	62.7	3.58 dd, 12, 5
		3.78 dd, 12, 2

rated under vacuum to give an aglycone (**1b**, 4.9 mg). The aglycone (**1b**) was crystallized from EtOH–H₂O and identified as entagenic acid (**8**). Entagenic acid (**1b**): Colorless crystals, mp 292–296 °C. $[\alpha]_D^{23}$ +28.6 (*c*=0.49, EtOH); ¹H- (400 MHz, pyridine-*d*₅) and ¹³C-NMR (100 MHz, pyridine-*d*₅); essentially the same as reported for entagenic acid (**8**)^{3,4}; HR-ESI-MS (positive-ion mode) *m/z*: 511.3392 [M+Na]⁺ (Calcd for C₃₀H₄₈O₅Na: 511.3399). The aqueous layer was subjected to HPLC analysis to detect D-glucosamine hydrochloride, D-xylose and D-glucose, which were identified by comparison of their retention times with those of authentic samples, D-glucosamine (*t_R*: 6.0 min, positive optical rotation).

Acetylation of Rheedeioside A (1) A solution of **1** (17.3 mg) in 2 ml of Ac₂O–pyridine (1:1) and DMAP (2.0 mg) was stirred at room temperature for 48 h. The reaction mixture was poured into ice-water and then extracted with EtOAc. The residue of the EtOAc extract was purified by silica gel CC [0.5 g, CHCl₃ and then CHCl₃:MeOH:H₂O] to give **1c** (19.0 mg). Nonadecacacetate (**1c**): Amorphous powder, $[\alpha]_D^{22}$ –37.0 (*c*=1.9, CHCl₃); ¹H-NMR (600 MHz, CDCl₃) δ : 0.62 (1H, br d, *J*=12 Hz, H-5), 0.68, 0.75, 0.84, 0.85, 0.91, 0.98, 1.33 (3H each, all s, H₃-24, 26, 25, 23, 29, 30 and 27, respectively), 1.85 (3H, s, CH₃CON–), 1.89 (3H, s), 1.96 (3H, s), 1.97 (6H, s), 1.98 (9H, s), 1.99 (3H, s), 2.01 (3H, s), 2.02 (3H, s), 2.04 (6H, s), 2.05 (6H, s), 2.081 (6H, s), 2.083 (6H, s), 2.09 (3H, s), 2.10 (3H, s) (20×CH₃COO–), 3.00 (1H, m, H-3), 4.43 (1H, d, *J*=6 Hz, H-1'), 4.45 (1H, d, *J*=6 Hz, H-1''), 4.58 (1H, d, *J*=8 Hz, H-1'), 4.60 (1H, d, *J*=6 Hz, H-1'''), 4.69 (1H, d, *J*=7 Hz, H-1''), 4.97 (1H, brs, H-1'''), 5.48 (1H, brt, *J*=4 Hz, H-12), 5.69 (1H, d, *J*=6 Hz, H-1'''); ¹³C-NMR (150 MHz, CDCl₃) δ : 15.5 (C-25), 16.5 (C-24), 17.3 (C-26), 18.6 (C-6), 20.2 (C-27), 20.4, 20.54, 20.57, 20.59 (×2), 20.62 (×2), 20.65, 20.69 (×2), 20.71 (×3), 20.84, 20.93 (×2), 20.96, 21.1, 21.3, 20.6 (20×CH₃COO–), 23.2 (CH₃CON–), 23.4 (C-11), 23.9 (C-30), 25.9 (C-2), 27.9 (C-23), 30.3 (C-20), 30.4 (C-22), 33.2 (C-29), 34.9 (C-7), 35.0 (C-21), 36.9 (C-10), 38.9 (C-1, 4), 40.9 (C-18), 41.3 (C-8), 46.1 (C-14), 46.3 (C-19), 46.9 (C-9), 47.0 (C-17), 55.0 (C-5), 69.6 (C-15), 76.2 (C-16), 90.1 (C-3), 126.4 (C-12), 141.0 (C-13), 172.8 (C-28), 103.4 (C-1'), 101.4 (C-1''), 101.7 (C-1'''), 101.3 (C-1'''), 93.1 (C-1'''), 100.5 (C-1), 107.1 (C-1'''), 169.0, 169.1, 169.3, 169.42, 169.43, 169.5, 169.6, 169.70, 169.74, 169.75, 169.88, 169.93, 170.1, 170.16 (×2), 170.19, 170.3, 170.4, 170.6, 170.7, 171.4 (21×CH₃COO–); HR-ESI-MS (positive-ion mode) *m/z*: 2406.9070 [M+Na]⁺ (Calcd for C₁₁₀H₁₅₃NO₅₆Na: 2406.9047).

Mild Alkaline Hydrolysis of Rheedeioside A (1) A solution of rheedeioside A (**1**) (17.3 mg) in 0.5% NaOMe (1.0 ml) was stirred at 20 °C for 18 h. The reaction mixture was neutralized with Dowex HCR W2 (H⁺ form) and then the resin was removed by filtration. The filtrate was extracted with EtOAc and then the aqueous layer was evaporated under vacuum to give a desacetylrheedeioside A (**1a**, 14.0 mg). The desacetylrheedeioside A (**1a**) was identified as rheedeioside B (**2**). Desacetylrheedeioside A (=rheedeioside B) (**1a**=**2**): Amorphous powder, $[\alpha]_D^{24}$ –25.1 (*c*=0.14, MeOH); HR-ESI-MS (positive-ion mode) *m/z*: 1566.6899 [M+Na]⁺ (Calcd for C₇₀H₁₁₃NO₃₆Na: 1566.6934).

Acid Hydrolysis of Rheedeiosides B, C and D (2–4) Solutions of **2**, **3** and **4** (1 mg each) were hydrolyzed in the same manner as described previously. HPLC analysis under the same conditions as above revealed the presence of D-glucosamine hydrochloride, D-xylose, D-apiose and D-glucose for **2**, D-glucosamine hydrochloride, D-xylose and D-glucose for **3**, and D-glucosamine hydrochloride, D-apiose, D-xylose and D-glucose for **4**. The sugars were identified by comparison of their retention times with those of authen-

tic samples, L-arabinose (t_R : 7.0 min, positive optical rotation).

Acknowledgements The authors are grateful for access to the superconducting NMR instrument (JEOL JNM-LA500 and ECA-600K) and the Thermo Fisher Scientific LTQ Orbitrap XL mass spectrometer at the Natural Science Center for Basic Research and Development, Hiroshima University. This work was supported in part by Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and the Japan Society for the Promotion of Science. Thanks are also due to the Takeda Science Foundation for the financial support.

References

- 1) Nzowa L. K., Barboni L., Teponno R. B., Ricciutelli M., Lupidi G., Quassinti L., Bramucci M., Tapondjou L. A., *Phytochemistry*, **71**, 254—261 (2010).
- 2) Tapondjou L. A., Miyamoto T., Mirjolet J.-F., Guilbaud N., Lacaille-Dubois M.-A., *J. Nat. Prod.*, **68**, 1185—1190 (2005).
- 3) Okada Y., Shibata S., Ikekawa T., Javellana M. J., Kamo O., *Phytochemistry*, **26**, 2789—2796 (1987).
- 4) Okada Y., Shibata S., Kamo O., Okuyama T., *Chem. Pharm. Bull.*, **36**, 5028—5030 (1988).
- 5) Dai J., Kardono L. B. S., Tsauri S., Padmawinata K., Pezzuto J. M., Kinghorn A. D., *Phytochemistry*, **30**, 3749—3752 (1991).
- 6) Kurashima K., Fujii M., Ida Y., Akita H., *Chem. Pharm. Bull.*, **52**, 270—275 (2004).
- 7) Mimura A., Sumioka H., Matsunami K., Otsuka H., *J. Nat. Med.*, **64**, 153—160 (2010).
- 8) Ishii T., Yanagisawa M., *Carbohydr. Res.*, **313**, 189—192 (1998).