#### Synthetic Methods

### Strain Release in C-H Bond Activation?\*\*

Ke Chen, Albert Eschenmoser,\* and Phil S. Baran\*

The field of C-H activation and the logic that underlies its use in complex-molecule synthesis are developing at a rapid pace.<sup>[1]</sup> This is due, in part, to the great potential that such transformations could have on the various "economies" of synthesis.<sup>[2]</sup> Comprehensive observations have pointed to the importance of steric and electronic factors governing the relative rates and selectivities of such reactions.<sup>[3]</sup> Yet, in order to plan complex-molecule total syntheses that utilize one or multiple C-H activation steps, a profound understanding of even the subtlest reactivity trends is needed. In particular, it has been observed on multiple occasions that equatorial C-H bonds react more rapidly than those oriented axially (Scheme 1).<sup>[4]</sup> This curious axial–equatorial rate ratio appears to be independent of the reagent system employed and in some cases can be exploited to achieve site-specific C-H activation.<sup>[4c,f,5]</sup> So far, explanations for these scattered observations have remained ambiguous. Here, a reactivity factor that apparently has been ignored thus far in this context is proposed, one that we suspect, besides steric hindrance to reagent approach and C-H bond nucleophilicity,<sup>[6]</sup> to be coresponsible for the more rapid activation of equatorial vs. axial C-H bonds in these tertiary settings.

In 1955, one of us proposed strain release (Scheme 2A) to explain the relative rates of reactions in which an equatorial hydrogen is also removed more rapidly than its axial counterpart, namely, in the oxidation of steroidal secondary alcohols with chromic acid (see Supporting Information for an English translation of this paper).<sup>[7]</sup> The rate enhancement in these reactions is attributed to a release of strain (1,3-diaxial interactions) in the transition state of going from an sp<sup>3</sup> to an sp<sup>2</sup> carbon. At the time, this work convincingly contradicted and corrected the theory<sup>[8]</sup> according to which the difference in oxidation rate of axial and equatorial alcohols was due to steric hindrance to proton abstraction by a base. In a later paper, the strain release hypothesis was corroborated in

[\*] Dr. K. Chen, Prof. P. S. Baran Department of Chemistry, The Scripps Research Institute 10550 North Torrey Pines Road, La Jolla, CA 92037 (USA) Fax: (+1) 858-784-7375 E-mail: pbaran@scripps.edu Prof. A. Eschenmoser Laboratory of Organic Chemistry, Swiss Federal Institute of Technology, Hönggerberg HCI H309 Wolfgang-Pauli-Strasse 10, 8093 Zürich (Switzerland) E-mail: eschenmoser@org.chem.ethz.ch

- [\*\*] We thank Tanja Gulder for the translation of ref. [7] and Jun Shi for technical assistance. Financial support for this work was provided by Amgen, Bristol-Myers Squibb, the Skaggs Institute, and the NIH (Grant CA184785).
  - Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.200904474.

C-H activation through hydroxylation with ozone:[4a] Relative reactivity ΟН O<sub>3</sub> then LAH 1 O<sub>3</sub> then LAH 5.6 C-H activation through hydroxylation with peracids:[4b] Relative reactivity p-nitroperoxy benzoic acid Me p-nitroperoxy Me benzoic acid 1.6 он Me 3° vs 2° selectivity C-H activation through hydroxylation with dioxirane:<sup>[4c]</sup> OН н TFDO, 10 min ∠\_Me ∼Me ∙Me 9 88% Me TFDO, 4 min only 3° ∩OH ∙Me Zme observed ≥ 95% C-H activation through carbene insertion:[4d] Relative reactivity н CHCl<sub>3</sub> NaOH CHCI TBAC Me Me 1 fRu tΒι Me Me CHCl<sub>3</sub> NaOH TBAC CHCI 6 tBu tBu C-H activation through Fe catalysis:[4e] Turnover н он FeL<sub>n</sub>, H<sub>2</sub>O<sub>2</sub> Me 1.4 Me Me Me FeLn, H2O2 Me ∠ Me 3.8 C-H activation through Fe catalysis:<sup>[4f]</sup> vs 2° selectivity н он FeLn, H<sub>2</sub>O<sub>2</sub> ca. 2.6 50% Me Me FeLn, H2O2 only 3°

Scheme 1. Observations from the literature. LAH = lithiumaluminum hydride; TFDO = methyl(trifluoromethyl)dioxirane; TBAC = tetrabutylammoniumcyanoborohydride.

69%

он

observed

collaboration with Westheimer and Rozek,<sup>[9a]</sup> as well as by work by Wilcox et al.<sup>[9b]</sup>

A key step in the total synthesis of eudesmane terpenes<sup>[5]</sup> caused us to revisit this principle, one that has so far mainly been used to explain differences in the rates of alcohol oxidation<sup>[7,9a,b]</sup> and solvolysis.<sup>[10]</sup> As shown in Scheme 2B, the power of the Curci (TFDO) oxidation<sup>[4c]</sup> was vividly demonstrated by the conversion of 1 to 2. Among five tertiary centers present in 1, H<sup>1</sup> was selectively activated, leading the alcohol 2 in 82% isolated yield on a gram scale. Purely electronic considerations might have led one to predict that the tertiary center of the isopropyl group (H<sup>5</sup> in Scheme 2B) would be oxidized first since its <sup>13</sup>C NMR shift is 0.9 ppm more upfield than the carbon attached to H<sup>1,[11]</sup> Steric considerations might also support the supposition that H<sup>5</sup> will react first. Such unusual efficiency and site-selectivity led



## Communications



**Scheme 2.** A) Strain release in alcohol oxidation. B) Site-selective C-H oxidation: a strain release phenomenon?

us to consider that their origin be related to strain release in going to the electrophilic transition state of the oxidation. In fact, numerous mechanistic studies of carbene,<sup>[12a-c]</sup> nitrene,<sup>[12d]</sup> dioxirane,<sup>[12e]</sup> and other<sup>[12f,g]</sup> intermolecular C–H oxidations have suggested a somewhat "flattened tetrahedrality" of the tertiary carbon center in the rate-determining transition states of such reactions (not an actual radical or cation formation but an electrophilic carbon that is bent as if to become trigonal).

Of special interest in this context are Houk's calculations on the mechanism of dioxirane-mediated oxidations.<sup>[13]</sup> They point to "a concerted transition state with O-H abstraction much more advanced than O-C bond formation", and to a "polarized nature of the transition state [that] cause[s] the concerted oxygen insertions into tertiary CH bonds to be highly favored" [see Eq. (1)]. These conclusions seem perfectly compatible with a participation of strain release as rateinfluencing co-factor; they qualitatively combine concertedness with the structural prerequisite for strain release to operate in the formation of the transition state.

$$\begin{array}{c} \mathsf{Q} \\ \mathsf{CF}_{3} \\ \mathsf{H}_{-H} \\ \mathsf{R}_{-H} \end{array} \xrightarrow{\delta^{*}} \left[ \begin{array}{c} \mathsf{R}_{-}^{*} \\ \mathsf{R}_{-}^{*}$$

The dihydrojunenol system with its steric repulsion between two 1,3-diaxial methyl groups offers a rare opportunity to determine whether strain release may be a contributing factor in oxidations by an electrophilic oxidant besides the factors of C–H bond nucleophilicity and steric hindrance to reagent approach.<sup>[3]</sup> It is for this very purpose that a specific eudesmane-based probe—demethyl dihydrojunenol carbamate (**3**)—was designed (Figure 1). In such a probe, any decrease in reactivity as compared to **1** would reflect the importance of strain release in going to the transition state, since the electronic reactivity factors (C–H bond nucleophilicity) in both settings are as nearly identical as they possibly can be,<sup>[14]</sup> as is the steric environment at the site of oxidation.



*Figure 1.* The design of an eudesmane-based probe to account for the enhanced reactivity of equatorial C-H bonds.

The synthesis of **3** was conducted in a similar fashion as the preparation of  $\mathbf{1}$ ,<sup>[5]</sup> as shown in Scheme 3. Thus, the



Scheme 3. Synthesis of eudesmane 3.

decalin framework of enone 4 was forged using methyl vinyl ketone, 3-methylbutyraldehyde, lithium dimethylcuprate, and 5-bromopentene in six steps with 44% overall yield. Stereoselective hydrogenation then delivered intermediate 5 in 80% vield as the major isomer. The establishment of three requisite steric centers of demethyl dihydrojunenol 6 was achieved in the subsequent Birch reduction in 73% yield. Lastly, carbamate formation smoothly delivered desired eudesmane probe 3 in nearly quantitative yield. The structure and stereochemical assignment were verified by X-Ray crystallography (see Figure 1). Notably, probe 3 exhibits an identical conformation in the solid state and nearly identical <sup>13</sup>C NMR chemical shifts relative to **1** (the <sup>13</sup>C NMR shift at the key tertiary carbon atom differ by 0.1 ppm).<sup>[11a,b,15]</sup> Not surprisingly, probe 3 reacted rapidly with TFDO in high yield to give the tertiary alcohol 7.

In order to establish the presence of any reactivity difference between **1** and **3** (Scheme 4), the following two experiments were conducted on a 1:1 mixture of **1** and **3** and halted before reaching full conversion (both experiments were run in triplicate and on scales ranging from 5–25 mg). Standard Curci conditions<sup>[4c]</sup> led to a 3.0–3.1:1 ratio of tertiary alcohols **2** and **7**, along with recovered starting materials. To confirm that this result is not reagent-specific, the mixture was absorbed to silica gel and exposed to a stream of ozone<sup>[16]</sup> at



**Scheme 4.** TFDO- and Ozone-mediated oxidation of a 1:1 mixture of **1** and **3** points to the strain-release phenomenon in tertiary C–H oxidation (sm: starting materials).

0 °C for 20 min delivering a mixture of tertiary alcohols 2 and 7 in a 3.9-4.1:1 ratio (in addition to recovered starting materials). Whereas the reaction of TFDO with both 1 and 3 was selective (only trace quantities of byproducts were observed), the reaction of 1 with ozone proceeded more efficiently (74% 2 based on recovered 1) than with 3 (numerous uncharacterized byproducts were observed and the product was formed in 30% based on recovered 3). Thus, 1 not only reacts faster with ozone than 3, it also reacts with greater selectivity.

The observed rate ratio of 3:1 to 4:1 in these experiments might seem modest ( $\Delta G^{\#} \approx 0.6 \text{ kcal mol}^{-1}$ ) for serving as evidence for the operation of a reactivity factor, particularly when compared with corresponding ratios in chromic acid oxidations of secondary alcohols. Yet there is a mechanisticically significant difference between the latter reaction and an oxidative hydroxylation of a tertiary C-H bond, in the product of which the carbon remains tetrahedral. Such rate ratios in oxidative C-H activations are expected to be much smaller than in chromic acid oxidations of secondary alcohols (see Scheme 2A), leading us to consider the observed ratios of Scheme 4 as significant. There is a parallel between these observations and those of classical solvolysis studies wherein similar rate differences are observed.<sup>[10]</sup> As alluded to above, these results suggest that strain release considerations can help to predict not only relative rate but also selectivity in complex settings.

In an attempt to delineate the scope of this conclusion, a series of experiments involving dioxirane, carbene, and nitrene C-H activation were conducted on 1,1-dimethylcyclohexane (8). If the strain release factor was operative also in methylene C-H activation, one would expect these conditions to favor activation at the C-3 carbon. As shown in Scheme 5, after correcting for statistics, TFDO and carbene activation show little to no site specificity (ca. 1:1) while nitrene activation shows a small effect (ca. 1.5:1 favoring C-3). The transition state is likely to be more tight in insertion



**Scheme 5.** Strain release is a minor factor in methylene C–H activation.  $H_2esp = \alpha, \alpha, \alpha', \alpha'$ -tetramethyl-1,3-benzenedipropionic acid.

reactions at methylene groups and, therefore, less sensitive to the strain release factor. Furthermore, the repulsion between CH<sub>3</sub> (axial) and H (axial) in the 3-position is roughly four times smaller (0.9 vs  $3.7 \text{ kcal mol}^{-1}$ ) than the corresponding repulsion between two axial methyl groups (8 vs 1) and so the effect on oxidation rates should be correpondingly smaller.<sup>[3e]</sup>

However, it is in more complex systems containing methylene groups where the strain release factor might be operative, such as the case of sclareolide (15) (Scheme 6).



*Scheme 6.* Strain release may be an important contributor in methylene activation in complex settings.

Although **15** contains 26 hydrogen atoms (2 tertiary and 12 methylene), it should be possible to predict which C–H bond will react first with an electrophilic oxidant. On electronic grounds, CH<sub>2</sub> positions at ring A may be considered most reactive since they are furthest away from the electron-withdrawing lactone ring C. Taking into account strain release, the equatorial  $\alpha$ -CH bond at position C-2 of ring A is both the least sterically hindered and, due to the presence of two 1,3-diaxial interactions of the axial hydrogen at position C-2, expected to be most prone to the effects of strain release in the transition state of oxidation. Indeed, when commercially available sclareolide (**15**) was submitted to Du Bois' nitrene insertion chemistry,<sup>[31]</sup> product **16** (verified by NMR spectroscopy, see Supporting Information for details) was obtained in nearly quantitative yield (based on sulfonamide).

The reactivity comparison of **1** and eudesmane probe **3** brings to light the previously unrecognized importance of a reactivity factor in C–H activation that is in all likelihood attributable to strain release, since both the steric and electronic characteristics of these two substrates are nearly identical at the reacting site. The corroboration of these results in other settings will of course require more experi-

# Communications

ments (e.g. using substrates such as **15**). For the time being, it is tempting to consider that the remarkably consistent rate differences in C–H activation of equatorial versus axial tertiary C–H bonds (see Scheme 1) may result from a cooperation of reactivity factors that involve 1) C–H bond nucleophilicity, 2) steric hindrance to reagent approach, and 3) strain release in transition state formation. Most importantly, the work presented here may aid in the planning and execution of total syntheses that rely on the simplifying power of C–H activation logic.

Received: August 11, 2009 Revised: October 29, 2009 Published online: November 24, 2009

**Keywords:** C–H functionalization · conformational analysis · natural products · strain release

- For reviews, see a) R. H. Crabtree, J. Chem. Soc. Dalton Trans. 2001, 2437–2450; b) J. A. Labinger, J. E. Bercaw, Nature 2002, 417, 507–514; c) A. R. Dick, M. S. Sanford, Tetrahedron 2006, 62, 2439–2463; d) K. Godula, D. Sames, Science 2006, 312, 67– 72; e) H. M. L. Davies, J. R. Manning, Nature 2008, 451, 417– 424; f) X. Chen, K. M. Engle, D.-H. Wang, J.-Q. Yu, Angew. Chem. 2009, 121, 5196–5217; Angew. Chem. Int. Ed. 2009, 48, 5094–5115, and references therein.
- [2] T. R. Newhouse, R. W. Hoffmann, P. S. Baran, *Chem. Soc. Rev.* 2009, 38, 3010–3021, and references cited therein.
- [3] See reference [1], and the following reports: a) D. Seyferth, Y. M. Cheng, J. Am. Chem. Soc. 1973, 95, 6763-6770; b) D. H. R. Barton, N. K. Basu, M. J. Day, R. H. Hesse, M. M. Perchet, A. N. Starratt, J. Chem. Soc. Perkin Trans. 1 1975, 2243-2251; c) P. Bovicelli, P. Lupattelli, E. Mincione, J. Org. Chem. 1992, 57, 5052-5054; d) H. Chen, S. Schlecht, T. C. Semple, J. F. Hartwig, Science 2000, 287, 1995-1997; e) M. E. González-Núnez, G. Castellano, C. Andreu, J. Royo, M. Báguena, R. Mello, G. Asensio, J. Am. Chem. Soc. 2001, 123, 7487-7491; f) H. M. L. Davies, E. G. Antoulinakis, J. Organomet. Chem. 2001, 617-3.20 618, 47-55; g) J. Yang, B. Gabriele, S. Belvedere, Y. Huang, R. Breslow, J. Org. Chem. 2002, 67, 5057-5067; h) C. G. Francisco, A. J. Herrera, E. Suárez, J. Org. Chem. 2003, 68, 1012-1017; i) M. M. Díaz-Requejo, T. R. Belderrain, M. C. Nicasio, P. J. Pérez, Dalton Trans. 2006, 5559-5566; j) O. Daugulis, V. G. Zaitsev, D. Shabashov, Q. Pham, A. Lazareva, Synlett 2006, 3382-3388; k) A. Company, L. Gómez, M. Güell, X. Ribas, J. M. Luis, L. Que, Jr., M. Costas, J. Am. Chem. Soc. 2007, 129, 15766-15767; 1) K. W. Fiori, J. Du Bois, J. Am. Chem. Soc. 2007, 129, 562-568; m) M. S. Chen, M. C. White, Science 2007, 318, 783-787; n) T. Kurokawa, M. Kim, J. Du Bois, Angew. Chem. 2009, 121, 2815-2817; Angew. Chem. Int. Ed. 2009, 48, 2777-2779, and references therein.
- [4] a) H. Varkony, S. Pass, Y. Mazur, J. Chem. Soc. Chem. Commun. 1974, 437–438; b) H.-J. Schneider, W. Müller, J. Org. Chem. 1985, 50, 4609–4615; c) R. Mello, M. Fiorentino, C. Fusco, R. Curci, J. Am. Chem. Soc. 1989, 111, 6749–6757; d) I. R. Likhotvorik, K. Yuan, D. W. Brown, P. A. Krasutsky, N. Smyth, M. Jones, Jr., Tetrahedron Lett. 1992, 33, 911–914; e) K. Chen, L. Que, Jr., J. Am. Chem. Soc. 2001, 123, 6327–6337; f) L. Gómez, I. Garcia-Bosch, A. Company, J. Benet-Buchholz, A.

Polo, X. Sala, X. Ribas, M. Costas, *Angew. Chem.* **2009**, *121*, 5830–5833; *Angew. Chem. Int. Ed.* **2009**, *48*, 5720–5723.

- [5] K. Chen, P. S. Baran, Nature 2009, 459, 824-828.
- [6] We use the term "C-H bond nucleophilicity" when referring to the "electronic reactivity factor". It seems to us that this term pragmatically denotes the effect of what hyperconjugative and inductive electronic effects might transmit from neighboring substituents to the reaction site. Thus, this term can refer to both activation as well as deactivation of the C-H bond by neighboring substituents towards electrophilic "C-H activation" insertion reactions.
- [7] J. Schreiber, A. Eschenmoser, *Helv. Chim. Acta* 1955, 38, 1529– 1536.
- [8] a) D. H. R. Barton, *Experientia* 1950, 6, 316; b) D. H. R. Barton, J. Chem. Soc. 1953, 1027.
- [9] a) J. Rocek, F. H. Westheimer, A. Eschenmoser, L. Moldoványl,
  J. Schreiber, *Helv. Chim. Acta* 1962, 45, 2554–2567; b) C. F.
  Wilcox, Jr., M. Sexton, M. F. Wilcox, *J. Org. Chem.* 1963, 28, 1079–1082.
- [10] a) H. C. Brown, R. S. Fletcher, J. Am. Chem. Soc. 1949, 71, 1845–1854; b) K. Okamoto, S. Saitô, H. Shingu, Bull. Chem. Soc. Jpn. 1969, 42, 3288–3297; c) J. F. King, M. J. Coppen, Can. J. Chem. 1971, 49, 3714–3723; d) J. F. King, T. W. S. Lee, Can. J. Chem. 1971, 49, 3724–3732, and references therein.
- [11] a) For a previous example of using (<sup>1</sup>H) nuclear magnetic resonance to predict C–H activation pattern, see: R. B. Boar, J. *Chem. Soc. Perkin Trans.* 1 1975, 1275–1277; b) for a correlation of fractional charge and carbon atom core energy to <sup>13</sup>C chemical shifts, see: K. Rasmussen, *Acta Chem. Scand. A* 1980, 34, 155– 156.
- [12] a) H. W. L. Davies, T. Hansen, M. R. Churchill, J. Am. Chem. Soc. 2000, 122, 3063 – 3070; b) U. H. Brinker, G. Lin, L. Xu, W. B. Smith, J.-L. Mieusset, J. Org. Chem. 2007, 72, 8434–8451; c) J. Hansen, J. Autschbach, H. W. L. Davies, J. Org. Chem. 2009, 74, 6555–6563; d) D. S. Breslow, E. I. Edwards, R. Leone, P. v. R. Schleyer, J. Am. Chem. Soc. 1968, 90, 7097–7102; e) R. Curci, L. D'Accolti, C. Fusco, Acc. Chem. Res. 2006, 39, 1–9; f) D. H. Giamalva, D. F. Church, W. A. Pryor, J. Org. Chem. 1988, 53, 3429–3432; g) A. Tenaglia, E. Terranova, B. Waegell, J. Org. Chem. 1992, 57, 5523–5528.
- [13] X. Du, K. N. Houk, J. Org. Chem. **1998**, 63, 6480-6483, and references therein; see also R. D. Bach, J. L. Andrés, M.-D. Su, J. J. W. McDouall, J. Am. Chem. Soc. **1993**, 115, 5768-5775.
- [14] Due to the rigid chair–chair conformation, any hyperconjugation (electronic)effect of the bridge-headed methyl in **1** or the proton in **3** should be negligible (see ref. [4d]). Strict consideration of Charton's preferred  $\sigma_i$  parameters point to nearly identical electronic reactivity factors (C–H bond nucleophilicity) in both settings. See M. Charton, *Prog. Phys. Org. Chem.* **1981**, *13*, 119; also, for an example of correlation of  $\sigma_i$  parameters and siteselectivity in TFDO C–H oxidation, see ref. [3e].
- [15] The chemical shift of the reaction center in probe 3 is exactly 0.1 ppm higher than in 1, which might indicate somewhat less C– H bond nucleophilicity. However, such a small difference should be regarded as negligible. For example, the <sup>13</sup>C shift of the tertiary carbons in *cis* and *trans* decalin differ by ca. 7 ppm, yet the rate ratio of oxidation with ozone is only ca. 5.6:1 (see Scheme 1 and ref. [4a]). See ref. [11b] for a correlation of <sup>13</sup>C NMR shift and fractional charge.
- [16] Z. Cohen, E. Keinan, Y. Mazur, T. H. Varkony, J. Org. Chem. 1975, 40, 2141–2142.

#### 9708 www.angewandte.org