

A C₂-Symmetric Chiral Bis-Sulfoxide Ligand in a Rhodium-Catalyzed Reaction: Asymmetric 1,4-Addition of Sodium Tetraarylborates to Chromenones [*J. Am. Chem. Soc.* 2010, 132, 4552–4553]. Jun Chen, Junmin Chen, Feng Lang, Xiangyang Zhang, Linfeng Cun, Jin Zhu, Jingen Deng, and Jian Liao*

Page 4552. The name for compound **2** should be given as (*R*)-phenyl *tert*-butylsulfoxide.

In Scheme 1 (and on page S2 in the Supporting Information), the absolute configuration of compound $\mathbf{3}$ was mistakenly drawn as S instead of R. The correct Scheme 1 is shown below:

Scheme 1. Synthesis of (R,R)-1,2-Bis(tert-butylsulfinyl)benzene [(R,R)-1]

JA103812E 10.1021/ja103812e

Published on Web 06/10/2010

Phytochrome as Molecular Machine: Revealing Chromophore Action during the Pfr — Pr Photoconversion by Magic-Angle Spinning NMR Spectroscopy [*J. Am. Chem. Soc.* 2010, 132, 4431–4437]. Thierry Rohmer, Christina Lang, Christian Bongards, Karthick Babu Sai Sankar Gupta, Johannes Neugebauer, Jon Hughes, Wolfgang Gärtner, and Jörg Matysik*

Page 4433. Changes in ¹⁵N chemical shifts were inadvertently not shown in Figure 3. The corrected figure is presented here.




Figure 3. Changes in 13 C and 15 N chemical shift of the u-[13 C, 15 N]-PCB chromophore in Cph1Δ2 during the Pfr → Lumi-F (A), Lumi-F → Meta-F (B), and Meta-F → Pr (C) transitions. For each transition, the starting state is taken as reference and the size of the circles and squares refers to the difference in 13 C and 15 N chemical shift in the next state.

JA104192U

10.1021/ja104192u Published on Web 06/14/2010