

Cu—N3	2.035 (2)	C5—C6	1.409 (4)
Cu—N4	2.042 (2)	C6—C7	1.434 (4)
O1—C1	1.193 (3)	C6—C14	1.396 (3)
Cu ¹ —N1	2.481 (2)	C7—C8	1.346 (4)
O2—C2	1.196 (4)	C8—C9	1.428 (4)
N1—C1	1.147 (3)	C9—C10	1.405 (4)
N2—C2	1.119 (4)	C9—C13	1.398 (3)
N3—C3	1.329 (3)	C10—C11	1.363 (4)
N3—C14	1.356 (3)	C11—C12	1.385 (4)
N4—C12	1.330 (3)	C13—C14	1.430 (3)
N1—Cu—N2	95.27 (9)	C7—C6—C14	118.6 (2)
N1—Cu—N3	91.70 (8)	C6—C7—C8	121.1 (2)
N1—Cu—N4	172.26 (8)	C7—C8—C9	121.6 (2)
N2—Cu—N3	167.98 (9)	C8—C9—C10	124.6 (2)
N2—Cu—N4	92.35 (9)	C8—C9—C13	118.3 (2)
N3—Cu—N4	80.56 (7)	C10—C9—C13	117.0 (2)
Cu—N1—C1	143.3 (2)	C9—C10—C11	119.2 (3)
Cu—N2—C2	166.3 (2)	C10—C11—C12	120.3 (3)
Cu—N3—C3	128.8 (2)	N4—C12—C11	122.4 (3)
Cu—N3—C14	113.3 (2)	N4—C13—C9	123.3 (2)
C3—N3—C14	117.8 (2)	N4—C13—C14	116.3 (2)
Cu—N4—C12	129.2 (2)	C9—C13—C14	120.4 (2)
Cu—N4—C13	112.8 (1)	N3—C14—C6	123.6 (2)
C12—N4—C13	117.7 (2)	N3—C14—C13	116.4 (2)
O1—C1—N1	176.7 (3)	C6—C14—C13	120.0 (2)
O2—C2—N2	178.3 (3)	Cu ¹ —N1—Cu	92.59 (8)
N3—C3—C4	122.5 (2)	Cu ¹ —N1—C1	118.9 (2)
C3—C4—C5	120.0 (2)	N1 ¹ —Cu—N1	87.41 (8)
C4—C5—C6	119.1 (2)	N1 ¹ —Cu—N2	99.49 (9)
C5—C6—C14	117.0 (2)	N1 ¹ —Cu—N3	90.58 (8)
C5—C6—C7	124.3 (2)	N1 ¹ —Cu—N4	92.73 (7)

Symmetry code: (i) $-x, -y, 2 - z$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989). Cell refinement: *CAD-4 Software*. Data reduction: *MolEN* (Fair, 1990). Program(s) used to solve structure: *MolEN*. Program(s) used to refine structure: *MolEN*. Molecular graphics: *ORTEP* (Johnson, 1965) in *SHELXTL/PC* (Sheldrick, 1990). Software used to prepare material for publication: *MolEN*.

We gratefully acknowledge financial support for this study from the EWU Foundation and the EWU North-west Institute for Advanced Studies.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: PT1036). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Cromer, D. T. & Mann, J. B. (1968). *Acta Cryst.* **A24**, 321–324.
 Enraf–Nonius (1989). *CAD-4 Software*. Version 5. Enraf–Nonius, Delft, The Netherlands.
 Fair, C. K. (1990). *MolEN. An Interactive Intelligent System for Crystal Structure Analysis*. Enraf–Nonius, Delft, The Netherlands.
 Johnson, C. K. (1965). *ORTEP*. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee, USA.
 Julve, M., Verdaguier, M., De Munno, G., Real, J. A. & Bruno, G. (1993). *Inorg. Chem.* **32**, 795–802.
 Lloret, G., Julve, M., Faus, J., Ruiz, R., Castro, I., Mollar, M. & Philoche-Levisalles, M. (1992). *Inorg. Chem.* **31**, 784–791.
 Mauro, A. E., Klein, S. I., Saldana, J. S., de Simone, C. A., Zukerman-Schpector, J. & Castellano, E. E. (1990). *Polyhedron*, **9**, 2937–2939.
 Parker, O. J. & Breneman, G. L. (1993). *Polyhedron*, **12**, 891–895.
 Sheldrick, G. M. (1990). *SHELXTL/PC*. Release 4.1. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

Acta Cryst. (1996). **C52**, 1091–1093

cis-(*R,S*)-Bis(benzyl methyl sulfoxide)dichloroplatinum(II)

JOHN EBBIGHAUSEN,^a NICHOLAS FARRELL,^b KARIN LÖVQVIST^c AND ÅKE OSKARSSON^c

^aDepartment of Chemistry, Cook Physical Building, Burlington, Vermont 05405-0125, USA, ^bDepartment of Chemistry, Virginia Commonwealth University, 1001 Main Street, Box 2006, Richmond, VA 23284, USA, and ^cInorganic Chemistry 1, Chemical Center, University of Lund, PO Box 124, S-221 00 Lund, Sweden. E-mail: ake.oskarsson@inorgk1.lu.se


(Received 10 July 1995; accepted 20 November 1995)

Abstract

The coordination around the Pt atom in the title compound, [PtCl₂(C₈H₁₀OS)₂], is pyramidally distorted from *cis* square-planar; two chloride ions form Pt—Cl bonds of 2.310 (3) and 2.307 (3) Å and two benzyl methyl sulfoxide groups form Pt—S bonds of 2.241 (2) and 2.233 (2) Å.

Comment

There should be three isomers of the complex *cis*-bis-(benzyl methyl sulfoxide)dichloroplatinum(II): the (*S,S*), (*R,R*) and *meso* forms. The crystal structure of the (*S,S*) form as a methanol solvate have been determined by X-ray methods (Antolini, Folli, Iarossi, Schenetti & Taddei, 1991). We have crystallized and determined the structure of the *meso* form, (I).



(I)

The title compound is composed of neutral *cis*-[PtCl₂(MeBzSO)₂] complexes (Fig. 1). The closest Pt···Pt distance is 7.274 (1) Å. Both the benzyl methyl sulfoxide molecules are bonded *via* their chiral S atoms. The structure analysis reveals that the two S atoms have different chiralities, *R* and *S*, giving the *meso* form of the platinum complex. The S atoms are in approximately tetrahedral environments with angles ranging from 100.1 (5) (C—S—C) to 117.1 (3)° (Pt—S—O). The S—O and S—C bond distances are close to those found in other sulfoxides (Almeida, Hubbard & Farrell, 1992; Antolini *et al.*, 1991; Melanson & Rochon, 1975, 1987; Melanson, Chevrotire & Rochon, 1985).




Fig. 1. Molecular structure and atomic numbering of *cis*-[PtCl₂(MeBzSO)₂]. Displacement ellipsoids are drawn at the 50% probability level. Only one of three locations of the disordered phenyl group is shown.

One of the benzyl groups was found to be disordered (see *Experimental* and Table 1). The bond distances and angles in the other phenyl ring have close to expected values (Table 2). The dihedral angles between the disordered benzyl groups and the coordination plane are 61 (2)°; for the other benzyl group the dihedral angle is 33 (1)°. The largest deviation from the least-squares plane through the pyramidally distorted Cl₂PtS₂ core is shown by Cl(1) [−0.054 (3) Å]. The non-rigidly refined benzyl group is planar within 0.02 (1) Å.

The Pt—S and Pt—Cl distances are normal compared with those found in similar compounds (see references cited above). The Cl—Pt—Cl angle seems to be the most rigid angle in compounds of the type *cis*-[PtCl₂(sulfoxide)₂], only varying between 87.2 (1) and 88.5 (1)°, while the other angles around the Pt atom are more affected by the orientation of the sulfoxide group and steric and packing effects of the different types of sulfoxides.

Experimental

Benzyl methyl sulfoxide (racemic mixture) reacts with the platinum salt K₂PtCl₄ in aqueous ethanol solution to give the expected product *cis*-[PtCl₂(MeBzSO)₂] as a pale yellow precipitate. The crude material is a mixture of diastereomers, as evidenced by two peaks in the ¹⁹⁵Pt NMR spectrum (CDCl₃) at −3509 and −3498 p.p.m. relative to Na₂PtCl₆ in D₂O as external standard. The ¹H NMR spectrum confirmed the presence of diastereomers with two separate peaks for methyl resonance and two doublets of doublets for the diastereotopic benzyl protons of the sulfoxide ligand. The peaks are shifted downfield upon coordination, as expected (Farrell, Kiley, Schmidt & Hacker 1990). Recrystallization from EtOH gave colourless crystals of one diastereomer:

$\delta(\text{Pt}) = -3498$; $\delta(\text{S—CH}_3) = 3.31$ (singlet) and $\delta(\text{S—CH}_2) = 4.95$ p.p.m. (doublets of doublet); $\nu(\text{SO}) = 1103 \text{ cm}^{-1}$ {*c.f.* values found in K[PtCl₃(MeBzSO)] (Almeida, Hubbard & Farrell, 1992); $\delta(\text{Pt}) = -3011$ (*d*₆-acetone); $\delta(\text{S—CH}_3) = 2.95$ (singlet) and $\delta(\text{S—CH}_2) = 4.87$ p.p.m. (doublets of doublet); $\nu(\text{SO}) = 1099 \text{ cm}^{-1}$ }.

Crystal data

[PtCl₂(C₈H₁₀OS)₂]

M_r = 574.439

Monoclinic

*P*2₁/*c*

a = 8.4502 (9) Å

b = 12.616 (2) Å

c = 18.219 (3) Å

β = 92.304 (9)°

V = 1940.7 (8) Å³

Z = 4

D_x = 1.959 Mg m^{−3}

Mo *K*α radiation

λ = 0.7107 Å

Cell parameters from 23 reflections

θ = 8.8–24.7°

μ = 7.794 mm^{−1}

T = 295 K

Prism

0.17 × 0.15 × 0.13 mm

Pale yellow

Data collection

CAD-4 diffractometer

ω –2 θ scans

Absorption correction:

numerical by integration

from crystal shape

T_{min} = 0.333, *T_{max}* =

0.436

6055 measured reflections

5895 independent reflections

3574 observed reflections

[*I* > 1.5 σ (*I*)]

R_{int} = 0.037

θ_{max} = 30°

h = −11 → 11

k = 0 → 17

l = 0 → 24

3 standard reflections

monitored every 60

reflections

intensity decay: 0.90%

Refinement

Refinement on *F*

R = 0.0529

wR = 0.0521

S = 1.776

3574 reflections

178 parameters

H atoms: see below

w = 1/ σ^2 (*F*)

(Δ/σ)_{max} = 0.0495

$\Delta\rho_{\text{max}}$ = 1.91 e Å^{−3}

$\Delta\rho_{\text{min}}$ = −2.39 e Å^{−3}

Extinction correction: none

Atomic scattering factors

from *International Tables*

for X-ray Crystallography

(1974, Vol. IV)

Table 1. Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	$U_{\text{eq}} = (1/3)\sum_i U_{ii}a_i^*a_j^*a_i \cdot a_j$			<i>U_{eq}/U_{iso}</i>
	<i>x</i>	<i>y</i>	<i>z</i>	
Pt	0.62149 (4)	0.18058 (3)	0.85723 (2)	0.0405 (1)
Cl1	0.4117 (3)	0.1413 (2)	0.7766 (2)	0.063 (1)
Cl2	0.6032 (4)	0.0098 (2)	0.9014 (2)	0.082 (3)
S1	0.8169 (3)	0.2217 (2)	0.9391 (1)	0.050 (1)
S2	0.6215 (3)	0.3450 (2)	0.8118 (1)	0.041 (1)
O1	0.7835 (8)	0.3104 (5)	0.9893 (4)	0.062 (4)
O2	0.7380 (8)	0.4181 (5)	0.8430 (3)	0.057 (4)
C1	0.996 (1)	0.2483 (9)	0.8891 (5)	0.055 (6)
C2	1.132 (1)	0.2816 (7)	0.9382 (5)	0.047 (5)
C3	1.249 (1)	0.2126 (7)	0.9612 (6)	0.054 (6)
C4	1.375 (1)	0.2475 (10)	1.0066 (6)	0.061 (6)
C5	1.385 (1)	0.3513 (10)	1.0289 (7)	0.066 (8)
C6	1.266 (1)	0.4215 (9)	1.0089 (6)	0.067 (8)
C7	1.140 (1)	0.3888 (8)	0.9623 (6)	0.057 (6)
C8	0.878 (1)	0.1110 (9)	0.9945 (7)	0.084 (9)
C10	0.426 (1)	0.4022 (8)	0.8189 (6)	0.062 (6)
C15	0.635 (1)	0.3417 (8)	0.7135 (5)	0.052 (6)

C20†	0.794 (2)	0.305 (2)	0.698 (2)	0.058 (6)
C21†	0.833 (3)	0.199 (2)	0.687 (2)	0.058 (6)
C22†	0.988 (3)	0.172 (2)	0.670 (1)	0.058 (6)
C23†	1.103 (2)	0.251 (3)	0.666 (1)	0.058 (6)
C24†	1.064 (2)	0.356 (3)	0.678 (1)	0.058 (6)
C25†	0.909 (2)	0.384 (2)	0.694 (2)	0.058 (6)
C30‡	0.813 (2)	0.333 (2)	0.696 (2)	0.050 (6)
C31‡	0.874 (3)	0.233 (2)	0.679 (1)	0.050 (6)
C32‡	1.034 (3)	0.222 (2)	0.665 (1)	0.050 (6)
C33‡	1.133 (2)	0.311 (2)	0.667 (1)	0.050 (6)
C34‡	1.072 (2)	0.410 (2)	0.683 (1)	0.050 (6)
C35‡	0.911 (2)	0.421 (2)	0.698 (1)	0.050 (6)
C40§	0.793 (2)	0.289 (2)	0.693 (2)	0.055 (6)
C41§	0.792 (2)	0.179 (2)	0.683 (2)	0.055 (6)
C42§	0.933 (3)	0.126 (2)	0.669 (1)	0.055 (6)
C43§	1.074 (2)	0.183 (2)	0.666 (1)	0.055 (6)
C44§	1.075 (2)	0.292 (2)	0.676 (2)	0.055 (6)
C45§	0.934 (3)	0.345 (2)	0.690 (2)	0.055 (6)

† Occupancy = 0.40 (3); U_{iso} . ‡ Occupancy = 0.32 (3); U_{iso} . § Occupancy = 0.30 (3); U_{iso} .

Table 2. *Selected geometric parameters* ($\text{\AA}$, °)

Pt—C11	2.310 (3)	S2—O2	1.448 (7)
Pt—C12	2.307 (3)	S2—C10	1.81 (1)
Pt—S1	2.241 (2)	S2—C15	1.800 (9)
Pt—S2	2.233 (2)	C1—C2	1.49 (1)
S1—O1	1.479 (7)	C15—C20	1.45 (2)
S1—C1	1.828 (10)	C15—C30	1.55 (2)
S1—C8	1.79 (1)	C15—C40	1.55 (2)
C11—Pt—C12	87.8 (1)	Pt—S2—C10	109.3 (3)
C11—Pt—S1	177.31 (10)	Pt—S2—C15	110.4 (3)
C11—Pt—S2	88.56 (9)	O2—S2—C10	109.0 (4)
C12—Pt—S1	92.4 (1)	O2—S2—C15	109.5 (4)
C12—Pt—S2	176.0 (1)	C10—S2—C15	100.1 (5)
S1—Pt—S2	91.19 (9)	S1—C1—C2	112.8 (7)
Pt—S1—O1	115.7 (3)	S2—C15—C20	106 (1)
Pt—S1—C1	108.2 (3)	S2—C15—C30	108 (1)
Pt—S1—C8	112.6 (4)	S2—C15—C40	109 (1)
O1—S1—C1	110.6 (5)	C20—C15—C30	14 (1)
O1—S1—C8	107.5 (5)	C20—C15—C40	7 (1)
C1—S1—C8	101.4 (5)	C30—C15—C40	21 (1)
Pt—S2—O2	117.1 (3)		

It was possible to resolve three positions of the disordered phenyl group (see Table 1) using a rigid phenyl group in the refinement with C—C 1.40, C—H 0.95 Å and C—C—C 120°. An occupancy factor for each group was refined without restrictions: 0.40 (3), 0.32 (3) and 0.30 (3) resulted, giving a total occupancy for the phenyl group of 1.02 (5), which is an acceptable value. The H atoms of the disordered phenyl group were included in the rigid group and therefore refined with the whole group. The H atoms on C8, C10 and C15 were not included. The H atoms of the ordered benzyl group were included in calculated positions and included in the structure-factor calculations. The largest residual maxima and minima in the $\Delta\rho$ map were 0.92 Å from S1 and 1.55 Å from Cl(1), respectively.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989). Cell refinement: *SET-4* (Enraf–Nonius, 1989). Data reduction: *TEXSAN PROCESS* (Molecular Structure Corporation, 1985). Program(s) used to solve structure: *MITHRIL* (direct methods) (Gilmore, 1983). Program(s) used to refine structure: *TEXSAN LS*. Software used to prepare material for publication: *TEXSAN FINISH*.

Financial support from Swedish Natural Science Research Council and the Crafoord Foundation is gratefully acknowledged.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: AB1300). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Almeida, S. G., Hubbard, J. L. & Farrell, N. (1992). *Inorg. Chim. Acta*, **193**, 149–157.
 Antolini, L., Folli, U., Iarossi, D., Schenetti, L. & Taddei, F. (1991). *J. Chem. Soc. Perkin Trans. 2*, pp. 955–961.
 Enraf–Nonius (1989). *CAD-4 Software*. Version 5.0. Enraf–Nonius, Delft, The Netherlands.
 Farrell, N., Kiley, D. M., Schmidt, W. & Hacker, M. P. (1990). *Inorg. Chem.* **29**, 397–403.
 Gilmore, C. J. (1983). *MITHRIL. Computer Program for the Automatic Solution of Crystal Structures from X-ray Data*. Department of Chemistry, University of Glasgow, Scotland.
 Melanson, R. & Rochon, F. D. (1975). *Can. J. Chem.* **53**, 2371–2374.
 Melanson, R. & Rochon, F. D. (1987). *Acta Cryst.* **C43**, 1869–1872.
 Melanson, R., Chevrotire, C. & Rochon, F. D. (1985). *Acta Cryst.* **C41**, 1428–1431.
 Molecular Structure Corporation (1985). *TEXSAN. TEXRAY Structure Analysis Package*. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.

Acta Cryst. (1996). **C52**, 1093–1095

(Isothiocyanato)[(1*RS*,4*RS*,8*SR*,11*SR*)-1,4,8,11-tetraazacyclotetradecane]copper(II) Thiocyanate, [Cu(NCS)(cyclam)](SCN)

TIAN-HUEY LU,^{a*} TAHIR H. TAHIROV,^a YUH-LIANG LIU,^b CHUNG-SUN CHUNG,^b CHIUNG-CHIA HUANG^c AND YAW-SHIUN HONG^c

^aDepartment of Physics, National Tsing Hua University, Hsinchu, Taiwan 300, ^bDepartment of Chemistry, National Tsing Hua University, Hsinchu, Taiwan 300, and ^cDepartment of Applied Chemistry, Chung-Cheng Institute of Technology, Taishi, Taiwan 335

(Received 13 July 1995; accepted 6 December 1995)

Abstract

The crystal structure of the title compound, [Cu(NCS)-(C₁₀H₂₄N₄)](SCN), has been determined by X-ray diffraction. A mirror plane passes through the metal atom and bisects the complex with the isothiocyanato ligand lying on the plane. The Cu^{II} ion is five-coordinate in a distorted square-pyramidal environment, with four amine N atoms in equatorial positions and the isothiocyanate N atom in an axial position; its coordination ge-