## Chelating Diphosphite Complexes of Nickel(0) and Platinum(0): Their Remarkable Stability and Hydrocyanation Activity

Michael J. Baker, a Karl N. Harrison, a A. Guy Orpen, a Paul G. Pringle and Gordon Shawb

<sup>a</sup> School of Chemistry, University of Bristol, Cantocks Close, Bristol BS8 1TS, UK <sup>b</sup> ICI Fibres, Research and Technology Dept., Wilton Centre, Middlesbrough, Cleveland TS6 8JE, UK

The synthesis, stability and hydrocyanation catalytic activity of nickel(0) complexes of the diphosphite **1** derived from 2,2'-biphenol are described; the X-ray crystal structures of the ligand **1** and its platinum(0) complex **2b** are reported.

Diphosphines are very widely used ligands because the chelate complexes which they form are very stable and they often show catalytic activity, a property which has been particularly valuable in enantioselective catalysis where the rigidity of the chelate is exploited.<sup>1</sup> In contrast, until recently diphosphites were unknown as ligands<sup>2</sup> despite the importance of monophosphite complexes *e.g.* in the nickel–phosphite catalysed hydrocyanation of butadiene.<sup>3</sup> We report here the synthesis of diphosphite 1, its metal(0) complexes **2a** and **b**, and the hydrocyanation catalytic activity of the nickel complex **2a**. Recently, Union Carbide have reported that rhodium complexes of ligands similar to **1** are very active hydroformylation catalysts.<sup>4</sup>

Ligand 1 is readily made from 2,2'-biphenol and PCl<sub>3</sub> in the presence of Et<sub>3</sub>N and then crystallised from CH<sub>2</sub>Cl<sub>2</sub>-pentane. The X-ray crystal structure<sup>†</sup> of 1 (see Fig. 1) shows that the ligand is intrinsically chiral because of the twists about the biphenyl C-C linkages [(O)C-C-C-C(O) torsion angles -41.4(4), -61.7(3) and  $-42.4(4)^{\circ}$ ]. The diastereoisomer that crystallized has the *RRR* (*SSS*) configuration. The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of 1 at ambient temperatures and at -80 °C is a singlet consistent with only one of the three possible diastereoisomers being present or, less likely, rapid interconversion of these isomers on the NMR timescale by rotation about the three biphenyl C-C linkages.

The zerovalent nickel and platinum complexes 2a and b are made by treatment of the appropriate  $[M(cod)_2]$  (M = Ni or Pt; cod = cycloocta-1,5-diene) with diphosphite 1. The platinum complex 2b was crystallized from tetrahydrofuran-



<sup>+</sup> Crystal data for 1: C<sub>36</sub>H<sub>24</sub>O<sub>6</sub>P<sub>2</sub>,  $M_r$  = 614.5, triclinic, space group  $P\bar{1}$ (No. 2), a = 9.210(2), b = 11.541(3), c = 14.183(4) Å,  $\alpha = 93.57(2)$ , β = 96.06(2),  $\gamma = 100.55(2)^\circ$ , U = 1468.3(6) Å<sup>3</sup>, Z = 2,  $D_c = 1.39$ g cm<sup>-3</sup>,  $\bar{\lambda} = 0.71069$  Å,  $\mu$ (Mo-K $\alpha$ ) = 1.89 cm<sup>-1</sup>, F(000) = 636, T = 293 K. Data were collected on a Nicolet P3m diffractometer for a unique hemisphere of reciprocal space with 3.5 < 20 < 50°. The structure was solved by direct methods and refined by least-squares to R 0.045 for 3635 unique, observed ( $I > 1.5\sigma(I)$ ), absorption corrected, intensity data.

pentane solution and the structure of its pentane solvate determined by X-ray diffraction.<sup>‡</sup> In the solid state structure of **2b** (which has exact  $C_2$  symmetry) the phosphorus atoms are inequivalent since both of the ligands have an SRR configuration (see Fig. 2) with (O)C-C-C-C(O) torsion angles 41.5(11), -106.5(8) and  $-42.3(13)^{\circ}$ . The configuration of the ligand in the solid complex 2b differs from 1 both in absolute configuration and in the extent of twisting about the central biphenyl C-C linkage [(O)C-C-C-C(O) torsion angles;  $-106.5(8)^{\circ}$  in **2b** and  $-61.7(3)^{\circ}$  in **1**]. The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of 2b at +30 °C is two singlets (5:1 ratio) with <sup>195</sup>Pt satellites and at -80 °C two triplets with satellites. The spectrum at low temperature is consistent with the solid state structure of 2b, *i.e.* an  $A_2X_2$  spin system arising from the inequivalent phosphorus environments; there are twelve possible diastereoisomers of 2b and it is therefore remarkable that apparently only one of these is present in solution at low



Fig. 1 Molecular struture of 1 showing atom labelling scheme. All hydrogens have been omitted for clarity. Important dimensions include; bond lengths (Å): P(1)-O(1) 1.631(2), P(1)-O(2) 1.635(2), P(1)-O(3) 1.631(2), P(2)-O(4) 1.631(2), P(2)-O(5) 1.613(2), P(2)-O(6) 1.630(2); bond angles (°): O(1)-P(1)-O(2) 99.5(1), O(1)-P(1)-O(3) 91.7(1), O(2)-P(1)-O(3) 101.9(1), O(4)-P(2)-O(5) 102.1(1), O(4)-P(2)-O(6) 92.7(1), O(5)-P(2)-O(6) 100.7(1); torsion angles (°): C(11)-C(12)-C(22)-C(21) -41.4(4), C(31)-C(32)-C(42)-C(41) -61.7(3), C(51)-C(52)-C(62)-C(61) -42.4(4).

‡ Crystal data for **2b**·2C<sub>5</sub>H<sub>12</sub>: C<sub>82</sub>H<sub>72</sub>O<sub>12</sub>P<sub>4</sub>Pt,  $M_r = 1568.5$ , monoclinic, space group *I2/a* (No. 15), a = 23.094(5), b = 14.044(2), c = 24.164(5) Å,  $\beta = 113.53(2)^\circ$ , U = 7370(2) Å<sup>3</sup>, Z = 4,  $D_c = 1.41$  g cm<sup>-3</sup>,  $\overline{\lambda} = 0.71069$  Å,  $\mu$ (Mo-K $\alpha$ ) = 20.7 cm<sup>-1</sup>, F(000) = 3024, T = 293 K. Data were collected on a Nicolet P3m diffractometer for a unique quadrant of reciprocal space with  $3.5 < 20 < 50^\circ$ . The structure was solved by heavy atom methods and refined by least-squares to *R* 0.039 for 4245 unique, observed ( $I > 1.5\sigma(I)$ ), absorption corrected, intensity data. For both structures atomic coordinates, bond lengths and angles, and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre. See Notice to Authors, Issue No. 1.



Fig. 2 Molecular structure of 2b showing atom labelling scheme. All hydrogens have been omitted for clarity. Important dimensions include; bond lengths (Å): Pt–P(1) 2.233(1), Pt–P(2) 2.263(2), P(1)–O(1) 1.609(4), P(1)–O(2) 1.634(4), P(1)–O(3) 1.609(5), P(2)–O(4) 1.608(5), P(2)–O(5) 1.611(4), P(2)–O(6) 1.618(5); bond angles (°): P(1)–Pt–P(2) 115.3(1), P(1)–Pt–P(1a) 114.3(1), P(1)–Pt–P(2a) 102.2(1), P(2)–Pt–P(2a) 107.8(1); torsion angles (°): C(11)–C(12)–C(22)–C(21) 41.5(11), C(31)–C(32)–C(42)–C(41) – 106.5(8), C(51)–C(52)–C(62)–C(61)–42.3(13).

temperatures. The phosphorus atoms are equilibrated on the NMR timescale at +30 °C presumably by rapid inversion of the P(biphenoxide) rings giving rise to the major <sup>31</sup>P NMR singlet. The <sup>195</sup>Pt{<sup>1</sup>H} NMR spectrum at +30 °C shows two quintets (5:1 ratio) at similar chemical shift and we therefore conclude that a second diastereoisomer of **2b** is present at ambient temperatures. Similar solution NMR behaviour is observed for the nickel analogue **2a**.

## J. CHEM. SOC., CHEM. COMMUN., 1991

Table 1 Catalytic hydrocyanation of butadiene experiments

Catalyst	Yield of product/g	% Nitriles <sup>b</sup>	% 3PNc	Turnover
2a	2.38	61	52	1430
3	2.15	79	69	35
$3 + 6L^{d}$	3.19	75	63	310

<sup>*a*</sup> Conditions used: butadiene (4 cm<sup>3</sup>, 46 mmol) and Me<sub>2</sub>CO·HCN (3.1 cm<sup>3</sup>, 34 mmol) with catalyst (0.088 mmol) in 10 cm<sup>3</sup> metal bomb, stirred for 18 h at 140 °C. Each run was duplicated to ensure consistency; % yields determined by GC. The total number of turnovers for each catalyst was determined by sequentially reducing the amount of catalyst used until the yield of nitriles was seen to drop significantly. <sup>*b*</sup> The major non-hydrocyanation product is 4-vinyl-1-cyclohexene. <sup>*c*</sup> Proportion of the nitrile yield that is 3-pentenenitrile rather than 2-methyl-3-butenenitrile. <sup>*d*</sup> Complex **3** in the presence of 6 equiv. of tris(*p*-tolyl)phosphite.

The Du Pont catalyst  $[Ni{P(OC_6H_4Me-p)_3}_4]$  3 is outstandingly effective for the hydrocyanation of butadiene<sup>5</sup> but the intrinsic instability of the catalyst necessitates the presence of large excesses of phosphite ligand. By contrast the nickel chelate 2a is a very robust compound, e.g. toluene solutions of 2a when exposed to air for 14 days show no indication of decomposition (by <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy) whereas exposure of similar solutions of 3 to air lead to complete decomposition within 6 hours. The nickel chelate 2a is a catalyst for the hydrocyanation of butadiene. From the data given in Table 1, it can be seen that 2a, though less selective than 3, achieves at least 4 times the number of catalytic turnovers of the commercial catalyst system (3 + 6 equivalents)of ligand<sup>5</sup>) and 40 times the number of catalytic turnovers of catalyst 3 in the absence of added ligand. It is clear that the high stability of chelating diphosphite complexes gives them great potential as catalysts.

We should like to thank the SERC and ICI for supporting this work and Johnson Matthey for their generous loan of platinum salts.

Received, 30th November 1990; Com. 0/05409G

## References

- 1 See, *Homogeneous Catalysis with Metal-Phosphine Complexes*, ed. L. H. Pignolet, Plenum Press, New York, 1983.
- 2 J. R. Bleeke, A. J. Donaldson and W. J. Peng, Organometallics, 1988, 7, 33.
- 3 C. A. Tolman, R. J. McKinney, W. C. Seidel, J. D. Druliner and W. R. Stevens, *Adv. Catal.*, 1985, **33**, 1, and references therein.
- 4 US Pat 4 769 498, 1988, to Union Carbide.
- 5 US Pat 3 766 237, 1973, to Du Pont.