# **CHEMISTRY** A European Journal



# **Accepted Article** Title: Residue and site-specific labeling of CCR5 RNA pseudoknot reveals internal motions that correlate with microRNA binding

Authors: Bin Chen, Andrew P Longhini, Felix Nußbaumer, Christoph Kreutz, Jonathan Dinman, and T Kwaku Dayie

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Chem. Eur. J. 10.1002/chem.201705948

Link to VoR: http://dx.doi.org/10.1002/chem.201705948

Supported by ACES



# Residue and site-specific labeling of *CCR*5 RNA pseudoknot reveals internal motions that correlate with microRNA binding

Bin Chen<sup>[a,b]</sup>, Andrew P. Longhini<sup>[b]</sup>, Felix Nußbaumer<sup>[c]</sup>, Christoph Kreutz<sup>[c]</sup>, Jonathan Dinman<sup>[a]</sup>\*, & T. Kwaku Davie<sup>[b]</sup>\*

Abstract: Conformational dynamics of RNA molecules play a critical role in governing their biological functions. Measurements of RNAs' dynamic behaviour shed important light on sites that interact with their binding partners or cellular stimulators. However, such measurements using solution NMR are difficult for large RNAs (>70 nt) due to severe spectral overlap, homonuclear <sup>13</sup>C scalar couplings, and linebroadening. In this work, a strategic combination of solid-phase synthesis, sitespecific isotopic labeled phosphoramidites and enzymatic ligation is introduced. This approach allowed us to positionspecifically insert isotopic probes into a 96 nt CCR5 RNA. Accurate measurements of functional dynamics using the Carr-Purcell-Meiboom-Gill (CPMG) relaxation dispersion (RD) experiments enabled extraction of the exchange rates and populations of this RNA. NMR chemical shift perturbation analysis of the RNA/microRNA-1224 complex indicated that A90-C1' of the pseudoknot exhibits similar changes in chemical shift observed in the excited state. This work demonstrates the general applicability of a NMR-labeling strategy to probe functional RNA structural dynamics.

The C-C chemokine receptor type 5 (CCR-5) is a coreceptor involved in membrane fusion of human immunodeficiency type 1 virus (HIV-1) particle and plays a critical role in facilitating viral entry into human cells.<sup>[1]</sup> Significant efforts have been made to develop therapeutic agents or strategies either targeting the receptor itself <sup>[2]</sup> or manipulating its gene <sup>[3]</sup> for treatment of HIV-1 infection. Recently, a highly-conserved pseudoknot motif was identified in the CCR-5 encoding mRNA that stimulates programmed -1 ribosomal frameshifting (-1 PRF) and that acts as an mRNA destabilizing element by directing translating ribosomes to a -1 frame termination codon.<sup>[4]</sup> Interaction between this element and a microRNA was proposed to

[a]	Dr. B. Chen, Prof. Dr. J. D. Dinman
	Department of Cell Biology and Molecular Genetics
	University of Maryland
	4062 Campus Dr., College Park, MD 20742 (USA)
	E-mail: dinman@umd.edu
[b]	Dr. A. P. Longhini, Prof. Dr. T. K. Dayie
	Center for Biomolecular Structure & Organization

- Center for Biomolecular Structure & Organization Department of Chemistry & Biochemistry University of Maryland 8314 Paint Branch Dr, College Park, MD 20782 (USA) Email: Dayie@umd.edu
- Mr. F. Nußbaumer, Prof. Dr. C. Kreutz Institute of Organic Chemistry and Center for Molecular Biosciences
   Innsbruck (CMBI), University of Innsbruck Innrain 80/82, 6020 Innsbruck (Austria)
   E-mail:christoph.kreutz@uibk.ac.at

Supporting information for this article is given via a link at the end of the document

stimulate -1 PRF, thus fine-tuning gene expression by controlling mRNA abundance, suggesting that this element may be novel for controlling HIV-1 viral replication.<sup>[4]</sup> Unlike other non-coding functional RNA elements currently being pursued as targets for various diseases,<sup>[5]</sup> the pseudoknot motif is within the coding region of *CCR5* mRNA.<sup>[4]</sup> Therefore, any information about the *CCR5* pseudoknot structure and interaction with miRNAs will provide insight into gene regulation connected with -1 PRF mechanism and foundation for RNA/miRNA-based drug design and development.

While structures are desirable, not all RNA molecules are amenable to high resolution structural characterization by X-ray crystallization and nuclear magnetic resonance (NMR) spectroscopy, mostly because RNAs are highly flexible. Their inherent plasticity renders them capable of sampling a vast ensemble of conformations.<sup>[6]</sup> RNA structures also typically undergo conformational changes in response to ligand binding, cellular signals or environmental fluctuations; indeed, interconversions among different conformational states are crucial for RNAs to fulfil their diverse biological functions.<sup>[7]</sup> Therefore, to better understand how *CCR*5 functions in -1 PRF and gene regulation, we need to not only determine its high resolution structures in free and in complex with miRNA-1224, but also understand its internal motions and dynamic behavior.

NMR, as a proven and powerful approach, has been applied extensively to characterize RNA dynamics over a wide range of timescales.<sup>[8]</sup> However, these applications have mainly been limited to small RNA systems (<50 nucleotides, nt) due to severe spectral overlap and line broadening, in addition to conformational heterogeneity in larger RNA species.<sup>[8a,9]</sup> To bypass these problems, "divide-and-conquer" strategies have been employed for structural characterization of sub-domains derived from larger functional RNA molecules. However, these approaches have led to questions regarding whether the isolated sub-domains adopt the same structures, behave with the same dynamic properties, and carry out the same functions as those encoded by the full-length RNA molecules.<sup>[10]</sup> A hybrid solid-liquid phase transcription method allowing region-specific isotope labeling for RNAs larger than 50 nt was previously reported.<sup>[11]</sup> This, however, was found to be highly sequence restrictive and generated very low yields for large RNAs.[11] Recently, coupling of site-specific isotope-labeling techniques with enzymatic ligation has expanded the size limitation of RNA NMR.<sup>[12]</sup> Of particular promise are approaches combining sitespecific isotope-labeling, solid-phase synthesis, and enzymatic ligation in investigating RNA structure and dynamics. Indeed, this chemo-enzymatic approach significantly simplifies the NMR spectra, allows unambiguous resonance assignment and identification of key residues involved in molecular recognition,

and enables accurate measurements of dynamic features at atomic resolution that are typically difficult to obtain for large RNA molecules.  $^{\rm [8b,e]}$ 

We report herein the characterization of site-specific motions of a 96nt CCR5 RNA -1 PRF pseudoknot element that are correlated to interaction with miR-1224 based on the strategy described above using NMR spectroscopy. We first prepared a key precursor (8,1'-13C) modified ATP by coupling (8-13C)adenine with (1'-13C)-ribose using chemo-enzymatic synthesis with an excellent yield above 90%,<sup>[13]</sup> followed bv dephosphorylation with recombinant shrimp alkaline phosphatase to afford (8,1'-<sup>13</sup>C)-adenosine (Supplementary Figure 1) (compound 1 in Scheme 1). The exocyclic amino  $N^6$ group of  $(8,1'-^{13}C)$ -adenosine (1) was then benzoylated to yield 2. The selective protection of the 2'-hydroxyl group with tertbutyldimethylsilyl chloride via transient blocking of the 5'- and 3'using di-tert-butylsilandiylditriflate gave hydroxyl groups intermediate 3. Selective removal of the 3',5'-di-tertbutylsilanediyl group and subsequent tritylation gave N6-Bz-5'-O-DMT-(8,1'-<sup>13</sup>C) adenosine (4). Lastly, the (8,1'-<sup>13</sup>C)-adenosine phosphoramidite building block 5, which is not commercially available, was obtained for the first time by treatment with 2'cyanoethyl N,N-diisopropylchlorophosphoramidite (Scheme 1). This building block is crucial for insertion of isotopic probes into CCR5 RNA residue-specifically to facilitate subsequent NMR studies.



**Scheme 1.** Synthesis of N<sup>6</sup>-Benzoyl-8,1'-<sup>13</sup>C-2'-O-TBDMS-adenosine phosphoramidite **5**. Reaction conditions: **(a)** Benzoylchloride, trimethylsilyl chloride, pyridine, 2.5h, rt, then water, 28% NH<sub>3</sub>, 39%. **(b)** Di-*tert*-butylsilyl bis(trifluoromethanesulfonate) in DMF, 0°C, 1h, then *tert*-butyldimethylsilyl chloride, 3 h, 60°C, 40%. **(c)** HF.Py in CH<sub>2</sub>Cl<sub>2</sub>, 0°C, 1h, then DMT-Cl in pyridine, rt, 3h, 91%. **(d)** CEP-Cl, D*i*PEA in THF, rt, 3h, 80%. Orange dot = <sup>13</sup>C.

Based on previous mutational studies that suggested Stem 2 of CCR5 pseudoknot motif interacts with miR-1224,<sup>[4]</sup> the RNA was split into two fragments such that a 24 nt donor fragment containing eight adenosine residues spans the proposed interacting sequence (Figure 1a). The donor fragments bearing (8,1'-13C)-modified adenosine at each of five positions were synthesized using RNA solid-phase synthesis (Supplementary Table 1 & Supplementary Figure 2). Unlabeled 2'-O-TBDMSprotected nucleoside phosphoramidite building blocks were first utilized to optimize the conditions for synthesizing the 24 nt CCR5 donor fragments. A prolonged coupling time (6 min) was employed to guarantee yields greater than 98%. Next, the synthesized 2'-O-TBDMS-(8,1'-<sup>13</sup>C)-adenosine phosphoramidite was introduced into the donor fragments at different locations of the RNA in combination with three other unlabeled 2'-O-TBDMSprotected nucleoside phosphoramidite building blocks. After successful incorporation of the (8,1'-13C)-adenosine label, we

deprotected these donor fragments using methylamine/ammonium hydroxide followed by treatment with 3M triethylamine trihydrofluoride (*TEA*, *3HF*) in anhydrous DMSO.<sup>[14]</sup> This desilylation agent was very efficient even in the presence of low moisture. After purification to homogeneity by denaturing polyacrylamide gel electrophoresis (PAGE), we obtained 0.3 µmol of the donor fragments typically from a 1 µmol synthesis scale column (**Supplementary Table 1**).

Next, we ligated an unlabeled T7 RNA polymerase synthesized 72 nt acceptor fragment to a solid phase synthesized residue/site-specific labeled 24-nt donor fragment (Figure 1). The unlabeled acceptor fragment had a protective 5'triphosphate group and a reactive 3'-OH group by virtue of using T7 RNA polymerase for the in vitro transcription. The synthesized residue/site-specific labeled 24-nt donor fragment was phosphorylated by T4 polynucleotide kinase to obtain the desired monophosphorylated group at the 5'-terminus. A donor fragment with different labeling patterns could also be prepared by in vitro transcription with T7 RNA polymerase and dephosphorylated using RNA 5'-polyphosphatase prior to ligation (Figure 1b). In general (Detailed in Supplementary Materials), excess amounts of the acceptor fragment and the DNA splint were used in the ligation reaction to achieve maximum conversion of the donor fragment into the ligated residue/site-specific isotope labeled product with a typical ligation yield (based on the donor fragment) of >60% (Figure 1b).



Figure 1 Ligation of the CCR5 pseudoknot element. (a) Scheme for preparation of segmentally and selectively <sup>13</sup>C-labeled CCR5 RNA. The unlabeled 72 nt acceptor fragment was synthesized by in vitro transcription. The donor was prepared by either in vitro transcription followed by enzymatic dephosphorylation using RNA 5' polyphosphatase or solid phase synthesis followed by phosphorylation with T4 polynucleotide kinase to obtain the desired monophosphorylated group at the 5'-terminus. A 40mer DNA splint was used to facilitate RNA ligation. (b) A 12% denaturing PAGE showing the ligation results. L1: de-phosphorylated donor fragment obtained from in vitro transcription; L2: phosphorylated donor fragment from solid phase synthesis; L3: DNA splint; L4: acceptor fragment; L5: ligation reaction at 0 h using the donor from L1; L6: after 3 h reaction of L5; L7: ligation reaction at 0 h using the donor from L2; L8: after 3 h reaction of L7; L9: CCR5 mRNA pseudoknot prepared by in vitro transcription. Similar to prior observation, [8f] addition of the DNA splint to the reaction system resulted in changes of the migration pattern of all components in the system. The numbers by the right side of the image (1, 2, 3, and 4) correspond to full length CCR5, 72nt acceptor, DNA splint, and 24nt donor fragment, respectively. Typically, excess amounts of the acceptor fragment and the DNA splint were applied and the ligation yield was >60% based on the conversion of the donor fragment which, in most cases, was prepared with different isotope labels.

The residue/site-specific insertion of <sup>13</sup>C probes into *CCR*5 RNA was confirmed by NMR spectroscopy. 2D <sup>1</sup>H-<sup>13</sup>C heteronuclear multiple quantum coherence (HMQC) spectra obtained for the

### COMMUNICATION

ligated and residue/site-specific labeled CCR5 RNA in the absence of miR-1224 exhibited a single cross peak corresponding to either ribose C1' or base C8 of the residue A95 (Figure 2a & 2b). Unambiguous assignments of A95 C1'/H1' was readily achieved by overlaying the ribose C1' HMQC spectrum with that of the CCR5 RNA containing all seventeen adenosine residues prepared from in vitro transcription with atom-specifically labeled (8,1'-<sup>13</sup>C)-ATP and unlabeled CTP/GTP/UTP (Figure 2c). Continued incorporation of <sup>13</sup>C probes at different positions of the RNA resulted in identification of ribose C1'/H1' atoms of A76, A79, A85, and A90 (Figure 2d & 2e) and base C8/H8 atoms of these residues (Supplementary Fig. 3a & 3b).



Figure 2 Heteronuclear multiple quantum coherence (HMQC) spectra of the CCR5 pseudoknot element. (a) Ribose C1' and (b) base C8 HMQC spectra of ligated CCR5 RNA bearing site-specific (8,1')-13C-labeled probes at residue A95. The donor fragment was prepared by solid phase synthesis on a DNA synthesizer first using unlabeled RNA amidites followed by incorporation of a selective (8, 1')-13C-labeled adenosine phosphoramidite at different positions. Upon ligation to the unlabeled acceptor fragment, the resulting ligated full length CCR5 pseudoknot element afforded a single peak at either ribose C1' or base C8 region. The projected 1D <sup>1</sup>H spectrum in (a) showed a single peak across a broader H1' region, demonstrating the success in isotope labeling of a specific residue of the full length 96nt RNA. (c) Overlay of ribose HMQC spectra obtained from two CCR5 RNA pseudoknot samples. The spectrum in blue was recorded for a full length CCR5 RNA with all 17 adenosine residues labeled selectively with (8,1')-13C atoms; while the other in red was from the ligated CCR5 A95 RNA. (d) Ligation between the unlabeled acceptor and four different residue-specific labeled donor fragments resulted in resonance assignments of four additional residues: A90, A85, A79, and A76 at both ribose C1' and base C8 regions. The spectrum in blue is the same as that in (c). This method allowed for straightforward resonance assignments and provided staring points for assignment of the neighboring residues based on 2D/3D NOESY experiments. (e) Secondary structure of CCR5 pseudoknot RNA based on previous SHAPE experiments.<sup>[4]</sup>

To distinguish the three remaining adenosine residues in the donor fragment, we used conventional NMR techniques and prepared a *CCR*5 RNA sample by ligating the unlabeled acceptor fragment and a donor fragment labeled only with eight site-specific (8,1'- $^{13}$ C)-adenosines. A 2D edited-edited NOESY spectrum obtained for the RNA sample showed inter-residue cross-peaks between A75H1' and A76H8, and between A95H1' and A96H8, allowing straightforward and unambiguous assignments of residues A75 and A96 (**Supplementary Fig. 3**).

The last adenosine A88 within the donor fragment region was exclusively determined (**Supplementary Figure 3**). The eight adenosine ribose anomeric atoms in the donor fragment of the ligated *CCR*5 RNA were well-resolved in the 2D HMQC spectrum (**Supplementary Figure 3b**).

In sum, this labeling strategy provides a number of advantages. First, use of isotopic probes residue/site-specifically helps to resolve substantial ambiguities in resonance assignments of RNAs of this size. Second, these labels are effective and reliable for quickly assigning large (>60 nt) RNAs. This represents an important advance for RNA structural studies by NMR spectroscopy, currently stuck at a median size of 25nt in the PDB. Third, these labels remove complicated scalar coupling effects or dipolar interactions with adjacent nuclei, thereby providing an ideal platform for performing, especially, CPMG RD experiments to investigate *CCR*5 conformational exchanges on micro- to millisecond timescales. <sup>[13b,15]</sup>; other RD experiments such as  $R_1$  and CEST will also benefit from this labeling pattern<sup>[8b,f,g]</sup>.



Figure 3 Chemical exchange in CCR5 revealed by <sup>13</sup>C CPMG RD experiments. (a) A 2D <sup>1</sup>H-<sup>13</sup>C HMQC spectrum of a ligated CCR5 RNA sample labeled only with site-specific (8,1'-13C)-adenosines within the donor fragment. This was recorded in the absence of miR-1224, at 298 K, at 600 MHz proton Larmor frequency, with  $T_{relax}$  = 0 ms. The expected eight anomeric H-C cross-peaks are well-resolved. (b) <sup>13</sup>C CPMG RD profiles of the ligated CCR5 RNA at 150 MHz carbon Larmor frequency. Red dots represent experimental data and black ones represent repeated experiments. Solid lines are best RD curve fittings of the CPMG profiles. Residues A75, A76, A88, A90, and A95 show non-flat dispersion profiles while A85 and A96 show flat <sup>13</sup>C dispersion profiles. Note that A79's resonance was too broad to fit. (c) CPMG RD profiles of A90 ribose anomeric carbon of the ligated CCR5 RNA at 150 (red dots) and 200 MHz (blue dots) carbon Larmor frequency. Black circles represent repeated experiments and solid lines are the best fits of the CPMG profiles. (d) NMR titration of the ligated CCR5\_A90 RNA with miR-1224. The 2D HMQC spectrum in magenta recorded in the absence of miR-1224 exhibits a single cross peak corresponding to ribose C1' of the residue A90. Upon addition of miR-1224, the ribose A90-C1' chemical shift changed dramatically from 90.3 ppm to 88.3 ppm (in cyan). The large chemical shift change (up to 2.0 ppm) indicates that the interaction with miR-1224 alters the local chemical environment of the A90 residues. In the presence of an equimolar amount of miR-1224, the CCR5 RNA is not completely converted to its bound form, with ~9% remaining in the free conformation. This is consistent

#### WILEY-VCH

with the EMSA result (Supplementary Figure 6). However, adding 20% extra amount of miR-1224 converts it to the completely bound form (data not shown).

A pseudo-3D <sup>1</sup>H-<sup>13</sup>C HMQC version of CPMG RD experiment was used to minimize relaxation loses and the effect of magnetic field inhomogeneity.<sup>[16]</sup> The eight well-dispersed <sup>13</sup>C probes of the ligated CCR5 RNA in the 2D <sup>1</sup>H-13C correlation map with  $T_{relax} = 0$  (Figure 3a) allowed for accurate measurements of the functional dynamics and subsequent extraction of information about the exchange processes in which the pseudoknot participates. As a result, the 1' anomeric carbons of adenosines A75, A76, A88, A90, and A95 of the ligated CCR5 RNA at 150 MHz carbon Larmor frequency showed non-flat dispersion profiles while that of A85 and A96 exhibited flat dispersion profiles (Figure 3b). Unfortunately the signal intensity of A79-C1' rapidly decreases during the CPMG relaxation delay which prevents further characterization and analysis. Additionally A76 and A96 appear to have shoulder peaks suggestive of multiple conformations. More detailed analyses are needed to elucidate the precise nature of these conformations.

CPMG RD experiments were also conducted at 200 MHz carbon Larmor frequency (magnet field strength of 18.8 T) (Figure 3c & Supplementary Figure 4). Subsequent global fitting resulted in extracting kinetic and thermodynamic parameters of the conformational transition.[13b,17] Initially, we expected to obtain the same dynamic parameters because the seven probes report on the same biological exchange process. Surprisingly, fits to a two-state exchange model resulted in residue-specific variations in dynamic behaviors: two probes (A85 and A96) showed no conformational exchange, whereas the other five displayed different thermodynamic parameters (Table 1). A76 samples a minor-populated state of  $P_{\rm B} \approx 30\%$ , while residues A75, A88, A90 and A95 sample an invisible-populated state of  $P_{\rm B}$  <10% (Table 1). The exchange rate constants  $(k_{ex})$  of A75, A88, and A90 were very close to one another within experimental error, whereas the rate parameters of A76 and A95 were roughly one magnitude higher (Table 1). Longitudinal relaxation rates were also recorded for these sites. Larger longitudinal relaxation rate of residues A85 and A96 indicated the presence of rapid local motion and high flexibility, indicating complicated motions occuring on various time scales (Supplementary Table 2 and Supplementary Figure 5). Thus these probes experience local independent complex motions or conformational exchanges.

A 2D <sup>1</sup>H-<sup>13</sup>C HMQC spectrum of the ligated and residue/sitespecific labeled *CCR5\_*A90 RNA in the absence of miR-1224 exhibited a cross peak corresponding to the ribose C1'/H1' (**Figure 3d**). Upon addition of miR-1224, the ribose H1' chemical shift was barely perturbed from 5.86 ppm to 5.83 ppm. In contrast, the ribose C1' chemical shift was strongly perturbed from 90.3 ppm to 88.3 ppm (**Figure 3d**). A similar result was obtained for the A90-C8/H8 chemical shift changes (data not shown here). The large chemical shift adjustment (up to 2.0 ppm) unambiguously indicated that miR-1224 binding alters the local chemical environment of the A90-C1' atom. It is notable that the chemical shift ( $\delta_{\rm C}$  90.3 ppm) of A90-C1' in the free form *CCR*5 pseudoknot suggests that the A90 residue either basepairs or is structurally constrained to a fixed conformation. The presence of miR-1224 triggered a highly specific interaction, flipping A90 out to participate in an intermolecular interaction, possibly base stacking with the miR-1224 bases but not base pairing as indicated by its up field chemical shift value ( $\delta_c$  88.3 ppm). The A90-C1' chemical shift of the minor-populated invisible state ( $|\Delta\omega|$ : 0.8 ± 0.1 ppm) likely samples local base-flipping motions to facilitate intermolecular recognition. The base C8 and ribose C1' of *CCR5\_*G89 also showed large chemical shift changes (data not shown), suggesting that the predicted intermolecular interaction with miR-1224. More detailed analysis using these and other RD measurements will be needed to clearly delienate the sign of chemical shift changes.

Our NMR data indicate that the secondary structure of the *CCR*5 RNA pseudoknot is slightly different from that derived from SHAPE experiments.<sup>[4]</sup> We are continuing with resonance assignment using this strategy to obtain a more accurate secondary structure of the *CCR*5 RNA pseudoknot using NMR spectroscopy.

In summary, in this study we have expanded the labeling protocol from previously reported (6,1'-<sup>13</sup>C) cytidine labels to (8,1'-<sup>13</sup>C) base and ribose building blocks for preparing residue/site-specifically isotopic labeled 96nt *CCR*5 RNA to study conformational dynamics.<sup>[8f,15a]</sup> The strategic combination of solid-phase synthesis of RNA with labeled phosphoramidites, enzymatic ligation, and NMR spectroscopy allows investigations into the structures and dynamics of large RNA systems, than have been possible until now, with biologically important functions that are otherwise impeded by severe spectral overlap.

#### **Experimental Section**

See details in Supplementary materials.

#### Acknowledgements

The authors would like to thank Dr. Paul Paukstelis for access to the DNA synthesizer and help with setup of DNA/RNA synthesis. This work was funded by grants to JDD from the NIH (R01 GM117177), and to TKD from NSF for NMR instrumentation (DBI1040158) and NIH (U54GM103297).

**Keywords**: CCR5 mRNA pseudoknot • MicroRNA • NMR spectroscopy • RNA solid-phase synthesis • Dynamics

a) G. Alkhatib, C. Combadiere, C. C. Broder, Y. Feng, P. E. Kennedy, P. M. Murphy, E. A. Berger, *Science*, 1996, 272, 1955-1958; b) R. W. Doms and J. P. Moore, *J. Cell Biol.* 2000, 151, 9-14; c) M. Samson, F. Libert, B. J. Doranz, *et al. Nature*, 1996, 382, 722–725; d) P. A. Zimmerman, A. Buckler-White, G. Alkhatib, *et al. Mol. Med.*, 1997, 3, 23–36; e) S. A. Gallo, C. M. Finnegan, M. Viard, *et al. Biochim. Biophys. Acta - Biomembr.*, 2003, 1614, 36–50; f) D. C. Chan, and P. S. Kim, *Cell*, 1998, 93, 681–684.

- COMMUNICATION
- a) M. Baba, O. Nishimura, N. Kanzaki, *et al. Proc. Natl. Acad. Sci. USA*, 1999, 96, 5698–5703; b) T. Dragic, A. Trkola, D. A. Thompson, *et al. Proc. Natl. Acad. Sci. USA*,2000, 97, 5639–5644; c) C. Watson, S. Jenkinson, W. Kazmierski, T. Kenakin, *Mol. Pharmacol.* 2005, 67, 1268–1282; d) K. Maeda, H. Nakata, Y. Koh, *et al. J. Virol.* 2004, 78, 8654–8662; e) A. Trkola, T. J. Ketas, K. A. Nagashima, *et al. J. Virol.* 2001, 75, 579–588; f) J. M. Strizki, C. Treblay, S. Xu, *et al. Antimicrob. Agents Chemother.* 2005, 49, 4911–4919; g) F. Tsamis, S. Gavrilov, F. Kajumo, *et al. J. Virol.* 2003, 77, 5201–5208.
- [3] P. Tebas, D. Stein, W. W. Tang, et al. N. Engl. J. Med., 2014, 370, 901–10.
- [4] A. T. Belew, A. Meskauskas, S. Musalgaonkar, et al. Nature, 2014, 512, 265–269.
- [5] a) M. Matsui D. R. Corey, *Nat. Rev. Drug Discov.* 2016, 16, 167–179;
  b) H. Ling, M. Fabbri, G. A. Calin, *Nat. Rev. Drug Discov.* 2013, 12, 847–865.
- [6] H. M. Al-Hashimi, N. G. Walter, Curr. Opin. Struct. Biol. 2008, 18, 321– 329.
- [7] E. A. Dethoff, J. Chugh, A. M. Mustoe, H. M. Al-Hashimi, *Nature*, 2012, 482, 322–330.
- [8] a) B. Furtig, C. Richter, J. Wohnert, H. Schwalbe, *ChemBioChem*, 2003, 4, 936–962; b) A. L. Hansen, H. M. Al-Hashimi, *J. Am. Chem. Soc.* 2007, 129, 16072–16082; c) Q. Q. Zhang, X. Sun, E. D. Watt, H. M. Al-Hashimi, *Science*, 2006, 311, 653–656; d) J. Rinnenthal, J. Buck, J. Ferner, A. Wacker, B. Furtig, H. Schwalbe, *Acc. Chem. Res.*, 2011, 44, 1292–1301; e) J. R. Bothe, E. N. Nikolova, C. D. Eichhorn, J. Chugh, A. L. Hansen, H. M. Al-Hashimi, *Nat. Methods*, 2011, 8, 919–931; f) B. Chen, R.M. LeBlanc, T. K. Dayie, *Angew. Chemie Int. Ed.* 2016, 55,

2724–2727; g) B. Zhao, A. L. Hansen, Q. Zhang, J. Am. Chem. Soc. 2014, 136, 20–23.

- [9] L. G. Scott, and M. Hennig in *Methods Mol. Biol.* vol. 452, Humana Press, Totowa, NJ, 2008, pp. 29-61.
- a) S. Spriggs, L. Garyu, R. Connor, M. F. Summers, *Biochemistry*, 2008, 47, 13064–13073; b) P. J. Lukavsky, *Virus Res.* 2009, 139, 166– 171; c) L. R. Ganser, H. M. Al-Hashimi, *Proc. Natl. Acad. Sci. USA*. 2016, 113, 13263–13265.
- [11] Y. Liu, E. Holmstrom, J. Zhang, et al. Nature, 2015, 522, 368–372.
- [12] a) C. S. Thakur, J. N. Sama, M. E. Jackson, B. Chen, T. K. Dayie, J. Biomol. NMR, 2010, 48, 179–192; b) L. J. Alvarado, R. M. LeBlanc, A. P. Longhini, et al. ChemBioChem, 2014, 15, 1573–1577; c) I. Kawahara, K. Haruta, Y. Ashihara, et al. Nucleic Acids Res. 2012, 40, e7; d) I. Kim, P. J. Lukavsky, J. D. Puglisi, J. Am. Chem. Soc. 2002, 124, 9338–9339; e) V. D'Souza, A. Dey, D. Habib, M. F. Summers, J. Mol. Biol. 2004, 337, 427–442; f) K. Lu, X. Heng, L. Garyu, et al. Science, 2011, 334, 242–245.
- [13] a) A. P. Longhini, R. M. LeBlanc, T. K. Dayie, *Methods*, 2016, 103, 11– 17; b) A. P. Longhini, R. M. LeBlanc, O. Becette, *et al. Nucleic Acids Res.* 2015, 44, e52.
- [14] E. Westmanu R. Stromberg, Nucleic Acids Res. 1994, 22, 2430–2431.
- [15] a) C. H. Wunderlich, R. Spitzer, T. Santner, K. Fauster, M. Tollinger, C. Kreutz, J. Am. Chem. Soc. 2012, 134, 7558–7569; b) D. M. Korzhnev, X. Salvatella, M. Vendruscolo, et al. Nature, 2004, 430, 586–590.
- [16] S. Grzesiek, A. Bax, *J Biomol NMR*, 1995, 6, 335–339.
- [17] C. H. Wunderlich, R. Spitzer, T. Santner, K. Fauster, M. Tollinger, C. Kreutz, J. Am. Chem. Soc. 2012, 134, 7558–7569.

## COMMUNICATION

### WILEY-VCH

#### Entry for the Table of Contents (Please choose one layout)

#### Layout 1:

Table 1 Kinetic and thermodynamic parameters extracted from CPMG RD experiments<sup>[a]</sup>

Residues (ribose C1')	<i>k</i> <sub>ex</sub> [S <sup>-1</sup> ]	<b>Р</b> в [%]	Δω [ppm]	X <sup>2</sup>	
A75	179 ± 138	8 ± 5	0.8 ± 0.1	1.08	—
A76	1014 ± 188	30 ± 12	$0.3 \pm 0.2$	1.12	
A88	356 ± 97	3 ± 1	0.6 ± 0.1	0.86	
A90	165 ± 121	9 ± 5	0.8 ± 0.1	1.46	
A95	943 ± 197	6 ± 2	$0.5 \pm 0.2$	0.96	

[a]: The CPMG RD experiments were carried out on the ligated *CCR*5 RNA sample labeled with site-specific  $(8,1^{-1^3}C)$ -adenosines within the donor fragment in the absence of miR-1224 at 298 K. The CPMG data of the ribose C1' of all five adenosine residues were fitted together to slow exchange two-state models and the results were derived from 1000 Monte Carlo runs (Detailed in Supplementary Materials). Note that strictly P<sub>B</sub> and  $\Delta w$  are not separable in the absence of other data.

### COMMUNICATION

(8,1'-13C)-adenosine phosphoramidite

A highly conserved pseudoknot Author(s), Corresponding Author(s)\* element within the mRNA encoding human C-C chemokine receptor type Page No. – Page No. 5 regulates its own gene's expression through a translational recoding event. Title A residue/site-specific labeling strategy was applied to insert isotopic probes into the pseudoknot. This ((Insert TOC Graphic here)) approach enables unambiguous assignments of key residues involved in intermolecular interaction with miRNA-1224 and results in accurate measurements of their dynamics at atomic resolution. Layout 2: COMMUNICATION Bin Chen<sup>[a]</sup>, Andrew P. Longhini<sup>[b]</sup>, Felix Nußbaumer<sup>[c]</sup>, Christoph Kreutz<sup>[c]</sup>, Jonathan Dinman<sup>[a]</sup>\*, & T. Kwaku Dayie<sup>[b]</sup>\* 87.5 CCR5 A95-C1'/H1 88.5 89.0 89.5 Residue and site-specific labeling of 90.0 90.5 Solid phase synthesis CCR5 RNA pseudoknot reveals 5.8 <sup>1</sup>H ppr 5.6 & enzymatic ligation

internal motions that correlate with

microRNA binding