ACS Medicinal Chemistry Letters

Technology Note

Subscriber access provided by CORNELL UNIVERSITY LIBRARY

Discovery of fluorescence polarization probe for the ELISAbased antagonist screening of #1-adrenergic receptors

Zhao Ma, Zhenzhen Liu, Tianyu Jiang, Tianchao Zhang, Huateng Zhang, Lupei Du, and Minyong Li ACS Med. Chem. Lett., Just Accepted Manuscript • DOI: 10.1021/acsmedchemlett.6b00048 • Publication Date (Web): 26 Aug 2016 Downloaded from http://pubs.acs.org on August 27, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Medicinal Chemistry Letters is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Discovery of fluorescence polarization probe for the ELISA-based antagonist screening of α_1 -adrenergic receptors

Zhao Ma, Zhenzhen Liu, Tianyu Jiang, Tianchao Zhang, Huateng Zhang, Lupei Du and Minyong ${\rm Li}^*$

Department of Medicinal Chemistry, Key Laboratory of Chemical Biology (MOE), School of Pharmacy, Shandong University, Jinan, Shandong 250012, China.

*KEYWORDS.α*₁*-adrenergic receptors, fluorescence polarization, ligandscreening, fluorescent probe, ELISA, GPCR.*

(This article is dedicated to Professor Lin Xia on the occasion of her 80th birthday)

ABSTRACT: High-throughput screening (HTS) of ligand library to find new active molecules for GPCRs is still a major interest, as well as an actual challenge. Fluorescence polarization (FP) assay portrays an essential role in HTS, however, in many cases, it was restricted by the absence of FP probes, the narrow measurement window, and low signal-to-noise (S/N) ratio. Herein, based on the modification of our previous probe 1 (QFL), we discovered an FP probe 3 (QGGFL) for α_1 -adrenergic receptors (α_1 -ARs), which has satisfactory fluorescence intensity, specific binding ability to receptors and suitable fluorescence properties to that were compatible with the filters in the FP system. Meanwhile, an "ELISA-like" strategy was designed for FP-based HTS assay, in which proteins were adhered into a solid phase, to improve the measurement window and S/N ratio. With fluorescent antagonist QGGFL and the ELISA strategy, we succeeded in establishing the first competitive binding FP assay for α_1 -AR antagonists as the alternative of the radioligand binding assay.

G protein-coupled receptors (GPCRs) represent the largest family of transmembrane proteins in the human genome that account for about 40% of therapeutic agents in the prescription pharmaceutical industry.1,2 Unbiased identification of compounds interacting with GPCRs is strongly dependent on highthroughput screening (HTS) of chemical libraries. Therefore, developing analytical methods to find new lead compounds targeting GPCRs is a great interest, as well as an actual challenge.³ Currently, HTS approaches for GPCR ligands mainly involve in the ligand-receptor binding assay using radioactive or fluorescent tracers.⁴ Radioligand binding is a wellestablished tool for screening at GPCRs, but with drawbacks of safety concerns and costly consumables.⁵ In parallel to the rapid development of fluorescent techniques, fluorescence assay is becoming a multifaceted tool in life science.⁶ Among these techniques, the nonradioactive, low-cost and selfreferencing FP assay using fluorescent ligands propose another credible alternative.^{7,8} Polarization value is defined as the ratio of fluorescence intensities parallel (I_I) and perpendicular (I_{\perp}) with respect to the plane-polarized excitation light.^{9, 10} In general, fast rotation of freely moving fluorophores will scramble polarization, while constrained fluorophores that bind to bulk receptors, will emit highly polarized fluorescence.



Figure 1. Chemical structure, U-shaped preferential conformation and structure modification strategy of our previous fluorescent ligand 1 (QFL) of the α_1 -AR.

Fluorescent ligands of GPCRs sprang up in recent decades, but their applications were restricted primarily to receptor visualization and localization.^{11,} ¹²Although fluorescent ligand-based assay has opened a brand new standpoint in the fields of GPCRs, it still requires further development.¹³ Due to the absence of FP probes, the narrow measurement window, and low signal-to-noise (S/N) ratio, FP assays have not been extensively applied.¹³ The key issue in such a matter is that the polarization is not totally or completely inexistent when fluorescent ligands bind to GPCRs because there is not quite a difference on rotation rate between the free fluorescent ligand and ligand confined by the receptor.

 α_i -Adrenergic receptors(α_i -ARs) are members of GPCR superfamily that mediate many physiological processes such as smooth muscle contraction, myocardial inotropy and chronotropy, and hepatic glu-

cose metabolism.¹⁴ As therapeutic targets, the antagonists of α_1 -AR play critical roles in the treatment of hypertension, benign prostatic hyperplasia (BPH) and lower urinary tract symptoms (LUTS).Moreover, someq1-adrenoceptor antagonists could induce prostate cancer cell apoptosis.^{15, 16} However, the screening of α_1 -AR antagonists is heavily dependent on radioligand binding assay at present.17,18 Although several fluorescent probes have been synthesized to study α . ARs in cells, a few of them are qualified for the highthroughput screening of antagonists, and no FPbased assay has been discovered or reported.¹⁸⁻²² In this paper, through the structure modification of our previous fluorescent antagonist QFL (1) of α_1 -AR²¹, we designed and synthesized an excellent FP antagonist probe QGGFL (3) with maintained binding affinity and increased fluorescence quantum yield (~10-fold), which could serve as an efficient toolkit to achieve an FP-based HTS assay (Figure 1). To settle the matters of narrow measurement window and low S/N ratio in FP assay, an "ELISA-like" strategy was designed to amplify FP effect. $^{\scriptscriptstyle 23,\,24}$ This strategy is to adhere $\alpha_{i}\text{-}AR$ proteins to the solid phase and then to measure the FP effect of fluorescent ligands binding to these proteins. When bound to the confined solid-phase proteins, the rotation of fluorescent ligands was constrained greatly, which would produce a considerable polarization effect.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59 60



Scheme 1. Synthesis of fluorescent ligands 2 and 3.

To achieve the FP-based ligand binding assay, the first thing we need to do is to afford an excellent FP probe with bright fluorescence and high binding affinity to receptors. In particular, the fluorescence intensity must be strong enough in order to increase S/N ratio and measure up to the microplate reader. Among masses of fluorophores, fluorescein performs brightly as the signal emitter in FP assay by virtue of its high quantum yield.^{25, 26} However, in our previous research on α_1 -AR, the reported fluorescent quinazoline-fluorescein conjugate QFL (1. Figure 1), possessing nanomolar binding affinity to α_1 -ARs, showed very low quantum yield (Table 1).²¹As a result, this compound failed to be employed in quantitive screening

of α_1 -AR antagonists and could not become an FP probe.²¹ With the help of Chemdraw 3D software, we found QFL's essential conformation with a crowded U-shape (Figure 1). In such a conformation, the quinazoline group comes close to the fluorescein core, which may exert an influence the conjugate plane of fluorophores, and thereby the quantum yield decreases. While the quinazoline moiety represents an interesting pharmacophoric scaffold of antagonist to develop FP agents that bind to the α -AR receptor tightly, to circumvent the issue about fluorescence intensity, two novel fluorescent ligands, QGFL and QGGFL (2 and 3, Figure 1) were designed by introducing the linker, y-aminobutyric acid (GABA), to conjugate pharmacophore and fluorophore. Although GABA isthe inhibitory neurotransmitter targeted at GABA receptors, the new compounds including GABA linker here will show no agonism activity to GABA receptors, because the primary amino group at 4-position in GABA is necessary for its agonism activity²⁷. As the linker, GABA could increase the flexibility of molecules and reduce the steric hindrance between quinazoline and fluorescein groups. The target compounds were conveniently synthesized as depicted in Scheme 1, in which piperazinesubstituted chlorine of 6 to afford compound 7, and then the amide condensation and N-BOC deprotection reactions were involved to introduce one or two GABA(s) to the key intermediates 9 and 10. Finally, the key intermediate reacted with FITC to provide 2 (QGFL) and 3 (QGGFL), respectively.

The affinities of two target compounds 2 (QGFL) and 3 (OGGFL) were assessed routinely by competitive binding assays on three α_1 -AR subtypes stably expressed in CHO cells, using radiolabeled $[^{3}H]$ prazosin, an established α_{1} -AR competitive antagonist. As expected, each compound revealed up to nanomolar affinities for all α_1 -AR subtypes (Table 1). This result also indicated that the new compounds could serve as antagonists of α_1 -AR. It should be noted that our previous research indicated clearly that the large moiety was permitted to conjugate to the N4 position of 2-piperazine group in the quinazoline pharmacophore without losing much binding affinity, but bulk fluorescein was not under consideration for selection.²¹ These results revealed that introducing a linker between fluorescein and quinazoline groups was beneficial for affinity improvement. Moreover, compound 3 (QGGFL) with the longer linker showed higher affinity than compound 2(QGFL).

A summary of spectral properties for all compounds is shown in Table 1as well. Compared to the reported compound 1 (QFL), the newly synthesized fluorescent ligands 2 (QGFL) and 3 (QGGFL) displayed about 15 nm red-shift in wavelength. Significantly, the major improvement in optical properties among them was fluorescence quantum yield and the quantum yield was positively associated with the length of the linker. In brief, compound 3 (QGGFL) 1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59 60 with two GABAs as the linker displayed about 10-fold quantum yield (27.2%) than compound 1 (QFL) without linker (2.6%), and compound 2(QGFL) with one GABA linker showed 6-fold yield (17.1%).

Table 1. Characterization of fluorescent ligands in spectroscopy and binding affinity to α_1 -ARs.

Compd	λ_{ex}	$\lambda_{em}/$	$\Phi/$	K_i^a/nM		
	nm	nm	%	α_{iA}	α_{1B}	α_{iD}
Phentolamine				1.3	7.7	13.8
QFL ²¹	485	517	2.6	4.7	7.3	20.7
2	500	530	17.1	33.5	8.1	31.3
3	500	530	27.2	4.0	2.7	23.8
Atropine				NA	NA	NA
3 1 1	1.0				_	

^a K_i was calculated from IC₅₀ using the Cheng-Prusoff equation (Equation S₃).

An essential requirement of a fluorescent ligand of GPCR is to visualize the target protein. As an improved fluorescent ligand, compound 3 (QGGFL) shows the reasonable receptor affinity and appropriate quantum yield, especially the latter. Next, we established the transfected HEK293A cells transiently with α_{IB} -AR and determined the ability of QGGFL to bind to such intact cells expressing α_1 -AR in culture. In confocal imaging study, the α_{1B} -AR transfected HEK293A cells could be "dyed" by strong green fluorescence when incubated with 300nM of compound 3 (QGGFL) for 20 min, and the strong fluorescence would weaken while a co-incubation of QGGFL (300 nM) and an excess of prazosin (3 µM) was carried Meanwhile, the untransfected out(Figure S₃). HEK293A cells were also employed as a negative control, in which very weak fluorescence was observed. This negative result indicated that non-specific binding is absent between QGGFL and cell components. The untransfected HEK293A cells also express some β-ARs natively, so these imaging results also indicated that QGGFL showed no interactions with the aminergic β -ARs. As is known, the α_{1B} -AR subtype mainly distributes on cytomembrane.28 In our confocal imaging, most cells with highlighted membrane by QGGFL came into view, indicative of a cell surface localization for α_{1B} -AR, which was highly consistent with the conclusion from Piascik and declared that the probe **3** (QGGFL) could recognize α,adrenoceptors specifically.²⁸ In addition, to check the specificity or selectivity of QGGFL, we measured its activities to three kinases (BMX, KIT and ALK) because quinazoline may cause the inhibitory activity against kinases.²⁹ As shown in Figure S5, we found all the GI_{50} values were greater than 10 μ M. Moreover, QGGFL also showed low affinities to other adrenergic receptors (α_2 - and β_1 -ARs). As a result, we can conclude that this fluorescent antagonist QGGFL with quinazoline core has few off-target activities.

At present up to, we have gotten an excellent fluorescent ligand 3 (QGGFL) with improved fluorescence intensity, the specific binding ability to α_1 -AR and suitable fluorescent properties that were compatible with the filters in our FP system. In order to identify whether the probe QGGFL may serve as an FP probe or if it could be qualified for the FP-based ligand screening of α_1 -AR, polarization at 520 nm of OGGFL with various concentrations of α ,-AR proteins was measured firstly using the 96-well or 384-well no-binding flat bottom black plate (Corning, USA) in classical FP assay. However, as shown in Figure S4Aa, the polarization values seemed no correlation with the amount of proteins. This set of irregular data incubated that the measurement window was too narrow for FP assay. To amplify the polarization signal, the ELISA strategy was surrogated (Figure S₄B), in which the enzyme or antigen was attached to the solid-phase surface.^{23, 24} By adhering to stationary phase, the constrained proteins could slow the spin of fluorescent ligands down, and thus, the fluorescent ligand would produce the obvious polarization effect. To achieve this strategy, 384-well flat bottom black plate with high protein-binding force (Fluotrac 600, Greiner) was employed in 1 h incubation at room temperature. By contrast with the irregular data in Figure S4Aa, three nice saturation curves in Figure S4Ba showed that addition of 16 nM QGGFL into increasing concentrations of three α_1 -ARproteins gave a saturable increase in FP. We also measured the Z' factor in both the traditional assay and the ELISA-like assay (Figure S4c), which is a statistical benchmark to judge whether the response in a particular assay is large enough to warrant further attention.^{30, 31}

The Z' factors calculated for our ELISA-based FP assay are 0.84, 0.77 and 0.70 for α_{1A} -, α_{1B} -, α_{1D} -AR proteins, respectively (Figure S4Bc), and in the traditional FP assay, the Z' factors are -0.49, -0.64 and -0.56, respectively (Figure S4Ac). A Z' factor greater than 0.5 means the assay is considered acceptable for a good HTS, while that less than o confirms there is too much overlap between the positive and negative controls for the assay to be used. In addition, when treated with excess membrane proteins (0.4 mg/mL), the FP value displays a dose-dependent enhancement, which yields K_d value of 0.56 ± 0.14, 1.21 ± 0.49 and 0.64 ± 0.30 nM for α_{1A} -, α_{1B} - and α_{1D} -AR, respectively (Figure 2). The results above indicates that compound₃ (QGGFL) is gualified to become a FP probe in the evolutionary ELISA-based FP assay. The specific interaction between QGGFL and α_1 -AR proteins reached equilibrium in 60 min and was saturable. Therefore, the FITC-derivative compound 3 (QGGFL) is suitable for the measurement of binding affinity of ligands.

We further investigated whether this "ELISA-like" FP strategy could be used in HTS forα₁-AR antago-

nists. Competitive binding experiments were carried out in the presence of 16 nM QGGFL and α_1 -AR membrane proteins (625 ng of α_{1A} -AR and α_{1B} -AR, 1.25 µg of α_{1D} -AR, respectively, Table S1). As known competitive antagonists, prazosin, doxazosin, and phentolamine could reduce FP in a concentration-dependent manner, which gave three anticipated sigmoidal response curves (Figure 2) and convincing IC_{50} and K_i values to three α_1 -AR subtypes with the similar values acquired from radioligand binding assay (Table S2). The difference of K_i values in the two assays may be caused by the different systems: the unbound ligands in radioligand assay were washed away while the FP values were recorded in the homogeneous reaction system. FP was almost enacted on the muscarinic receptor antagonist atropine, which could serve as the negative group for atropine do not bind to α -AR. Comparing with the radioligand competitive binding assay, the safer fluorescent ligands were employed and fewer membrane proteins (>10 fold) were consumed in the FP binding assay (Table S1). According to these results, we also concluded that the new fluorescent antagonist QGGFL could serve as a competitive antagonist.



Figure 2. Saturation of increasing concentrations of compound 3 (QGGFL) binding to α_{1A} -AR (A), α_{1B} -AR (B) and α_{1D} -AR (C) proteins (Final con. o.4 mg/mL). Competition curves of prazosin, doxazosin, phentolamine and atropine to three α_i -AR subtype, including α_{1A} -AR (D), α_{1B} -AR (E) and α_{1D} -AR (F) using the established "ELISA-like" FP assay.

In conclusion, developing analytical methods to identify compounds interacting with GPCRs or to find new drugs targeting GPCRs is drawing more and more attentions from scientists and researchers. Although FP assay can portray important roles in the HTS of the chemical library for GPCRs, it is restricted in many cases due to the lack of FP probes, the narrow measurement window, and low S/N ratio. To overcome these limitations in α_i -ARs, we firstly discovered and synthesized an ideal FP probe QGGFL, compound **3**, for α_i -ARs, which has improved quantum yield, high receptor affinity and the ability of receptor imaging specifically in living cells. Moreover, an "ELISA-like" strategy was designed, in which proteins are confined to the solid phase, in order to amplify the FP signal and increase the measurement window and S/N ratio. Based on the FP probe QGGFL and the ELISA strategy, we established a safe and affordable FP-based screening method for α_i -ARs for the first time, which could provide the convincing data of binding affinity to three α_i -AR subtypes of known antagonists, such as prazosin, doxazosin, and phentolamine, with the classical radioligand binding assay. To the best of our knowledge, this research is the first report that the ELISA strategy is applied in a small-molecule probe-based FP assay. It is our expectation that this strategy would find broad applications in the FP-based HTS for other target proteins to help addressing their signal issue.

ASSOCIATED CONTENT

Supporting Information

Full experimental procedures, analytical and spectral characterization data of all compounds, and results of cell imaging and FP assay. This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Author

*Minyong Li, Tel./fax: +86-531-8838-2076; E-mail: mli@sdu.edu.cn

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

ACKNOWLEDGMENT

The present work was supported by grants from the Qilu Young Professorship of Shandong University, the International Postdoctoral Exchange Fellowship Program, the Fok Ying Tong Education Foundation (No. 122036), the Program for Changjiang Scholars and Innovative Research Team in University (No. IRT13028), the Major Project of Science and Technology of Shandong Province (No. 2015ZDJS04001), the Shandong Key Research & Development Project (No. 2015GSF118166) and the Fundamental Research Funds of Shandong University (No. 2014JC008). Our cell imaging work was performed at the Microscopy Characterization Facility, Shandong University.

ABBREVIATIONS

GPCRs, G protein-coupled receptors; HTS, highthroughput screening; FP, fluorescence polarization; S/N, signal-to-noise; α_i -ARs, α_i -adrenergic receptors; BPH, benign prostatic hyperplasia; LUTS, lower urinary tract symptoms; GABA, γ -aminobutyric acid; ELISA, enzyme linked immunosorbent assay, EDCI, 1-Ethyl-3-(3dimethylaminopropyl)carbodiimide hydrochloride; HOBT, 1-hydroxybenzotriazole; FITC, fluorescein isothiocyanate; TFA, trifluoroacetic acid; BMX, bone marrow X

1

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56 57

58 59 60 kinase; ALK, anaplastic lymphoma kinase; KIT, v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homologue.

REFERENCES

- 1 Ke, N.; Nguyen, K.; Irelan, J.; Abassi, Y. A. Multidimensional GPCR profiling and screening using impedance-based label-free and real-time assay. In *G Protein-Coupled Receptor Screening Assays*; Springer: New York, 2015; pp. 215-226.
- 2 Sengupta, S.; Ye, K.; Scott, A. D.; Niu, B.; Bailey, M. H.; McLellan, M. D.; Wendl, M. C.; Wyczalkowski, M. A.; Ding, L. Sequence and structure-guided approach to identify functional mutations in G-protein coupled receptors. *Cancer Res.* 2015, 75, 61-61.
- 3 Oueslati, N.; Hounsou, C.; Belhocine, A.; Rodriguez, T.; Dupuis, E.; Zwier, J. M.; Trinquet, E.; Pin, J.-P.; Durroux, T. Time-resolved FRET strategy to screen GPCR ligand library. In *G Protein-Coupled Receptor Screening Assays*; Springer: New York, 2015; pp. 23-36.
- 4 Smith, D. S.; Eremin, S. A. Fluorescence polarization immunoassays and related methods for simple, highthroughput screening of small molecules. *Anal. Bioanal. Chem.* 2008, 391, 1499-1507.
- 5 Allen, M.; Reeves, J.; Mellor, G. High throughput fluorescence polarization: a homogeneous alternative to radioligand binding for cell surface receptors. *J. Biomol. Screen.* 2000, *5*, 63-69.
- 6 Leopoldo, M.; Lacivita, E.; Berardi, F.; Perrone, R. Developments in fluorescent probes for receptor research. *Drug discovery today* **2009**, *14*, 706-712.
- 7 Banks, P.; Gosselin, M.; Prystay, L. Fluorescence polarization assays for high throughput screening of G protein-coupled receptors. *J. Biomol. Screen.* **2000**, *5*, 159-167.
- 8 Jones, J. W.; Greene, T. A.; Grygon, C. A.; Doranz, B. J.; Brown, M. P. Cell-free assay of G-protein-coupled receptors using fluorescence polarization. *J. Biomol. Screen.* **2008**, *13*, 424-429.
- 9 Jia, T.; Fu, C.; Huang, C.; Yang, H.; Jia, N. A highly sensitive naphthalimide-based fluorescence polarization probe for detecting cancer cells. *ACS Appl. Mater. Interfaces* **2015**, *7*, 10013-10021.
- 10 Owicki, J. C. Fluorescence polarization and anisotropy in high throughput screening: perspectives and primer. *J. Biomol. Screen.* **2000**, *5*, 297-306.
- 11 Ma, Z.; Du, L.; Li, M. Toward fluorescent probes for Gprotein-coupled receptors (GPCRs). J. Med. Chem. 2014, 57, 8187-8203.
- 12 Vernall, A. J.; Hill, S. J.; Kellam, B. The evolving smallmolecule fluorescent-conjugate toolbox for class A GPCRs. *Br. J. Pharmacol.* 2014, *171*, 1073-1084.
- 13 Cottet, M.; Faklaris, O.; Zwier, J. M.; Trinquet, E.; Pin, J.-P.; Durroux, T. Original fluorescent ligand-based assays open new perspectives in G-protein coupled receptor drug screening. *Pharmaceuticals* **2011**, *4*, 202-214.
- 14 Piascik, M. T.; Perez, D. M. α₁-Adrenergic receptors: new insights and directions. *J. Pharmacol. Exp. Ther.* **2001**, *298*, 403-410.
- 15 Kyprianou, N.; Benning, C. M. Suppression of human prostate cancer cell growth by α_1 -adrenoceptor antagonists doxazosin and terazosin via induction of apoptosis. *Cancer Res.* **2000**, *60*, 4550-4555.
- 16 Partin, J.; Anglin, I.; Kyprianou, N. Quinazoline-based α₁adrenoceptor antagonists induce prostate cancer cell apoptosis via TGF-β signalling and IκBα induction. *Br. J. Cancer* **2003**, *88*, 1615-1621.

- 17 Nagatomo, T.; Tsuchihashi, H.; Sasaki, S.; Nakagawa, Y.; Nakahara, H.; Imai, S. Displacement by α_1 -adrenergic agonists and antagonists of ³H-prazosin bound to the α_1 -adrenoceptors of the dog aorta and the rat brain. *Jpn. J. Pharmacol.* **1985**, *37*, 181-187.
- 18 Mackenzie, J. F.; Daly, C. J.; Pediani, J. D.; McGrath, J. C. Quantitative imaging in live human cells reveals intracellular α_1 -adrenoceptor ligand-binding sites. *J. Pharmacol. Exp. Ther.* **2000**, *294*, 434-443.
- 19 Ma, J.; Hou, Z.; Song, Y.; Wang, L.; Guo, E. Visual and quantitative screening of α_1 -adrenoceptor antagonists in living cells using quantum dots. *ACS Comb. Sci.* **2014**, *16*, 155-159.
- 20 Morishima, S.; Suzuki, F.; Nishimune, A.; Yoshiki, H.; Akino, H.; Yokoyama, O.; Muramatsu, I. Visualization and tissue distribution of α_{1L} -adrenoceptor in human prostate by the fluorescently labeled ligand Alexa-488-silodosin. *J. Urol.* **2010**, *183*, 812-819.
- 21 Zhang, W.; Ma, Z.; Li, W.; Li, G.; Chen, L.; Liu, Z.; Du, L.; Li, M. Discovery of quinazoline-based fluorescent probes to α_1 -adrenergic receptors. *ACS Med. Chem. Lett.* **2015**, *6*, 502-506.
- 22 Ma, Z.; Lin, Y.; Cheng, Y.; Wu, W.; Cai, R.; Chen, S.; Shi, B.; Han, B.; Shi, X.; Zhou, Y.; Du, L.; Li, M. Discovery of the first environment-sensitive near-infrared (NIR) fluorogenic ligand for α₁-adrenergic receptors imaging in vivo. *J. Med. Chem.* **2016**, *59*, 2151-2162.
- 23 Elder, P.; Lewis, J. An enzyme-linked immunosorbent assay (ELISA) for plasma testosterone. *J. Steroid Biochem.* **1985**, *22*, 635-638.
- 24 de Camargo, Z. P.; Guesdon, J.; Drouhet, P. E.; Improvisi, L. Enzyme-linked immunosorbent assay (ELISA) in the paracoccidioidomycosis. *Mycopathologia* **1984**, *88*, 31-37.
- 25 Marholz, L. J.; Wang, W.; Zheng, Y.; Wang, X. A fluorescence polarization biophysical assay for the naegleria DNA hydroxylase Tet1. *ACS Med. Chem. Lett.* **2016**, *7*, 167-171.
- 26 Johnson, I.; Spence, M. T. *The Molecular Probes Handbook, A Guide to Fluorescent Probes and Labeling Technologies*; Life Technologies: Eugene, OR, 2010.
- 27 Dewey, S. L.; Brodie, J. D.; Ashby Jr, C. R. By administering a drug that increases central nervous system GABA (4aminobutyric acid) levels; for example gabapentin, valproic acid, progabide, gamma-hydroxybutyric acid, fengabine, cetylGABA, topiramate, tiagabine, or acamprosate. US Patents: US, 2005.
- 28 Chalothorn, D.; McCune, D. F.; Edelmann, S. E.; García-Cazarín, M. L.; Tsujimoto, G.; Piascik, M. T. Differences in the cellular localization and agonist-mediated internalization properties of the α_1 -adrenoceptor subtypes. *Mol. Pharmacol.* **2002**, *61*, 1008-1016.
- 29 Ravez, S.; Castillo-Aguilera, O.; Depreux, P.; Goossens, L. Quinazoline derivatives as anticancer drugs: a patent review (2011 - present). *Expert Opin. Ther. Pat.* 2015, 25, 789-804.
- 30 Zhang, J.-H.; Chung, T. D.; Oldenburg, K. R. A simple statistical parameter for use in evaluation and validation of high throughput screening assays. *J. Biomol. Screen.* **1999**, *4*, 67-73.
- 31 Lei, Y.; Hu, T.; Wu, X.; Wu, Y.; Bao, Q.; Zhang, L.; Xiang, H.; Sun, H.; You, Q.; Zhang, X. Affinity-based fluorescence polarization assay for high-throughput screening of polyl hydroxylase 2 inhibitors. *ACS Med. Chem. Lett.* **2015**, *6*, 1236-1240.

Lay Summary

As an important HTS method, fluorescence polarization assay is restricted in many cases due to the lack of FP probes, the narrow measurement window, and low S/N ratio. For the case of α_1 -ARs, we firstly designed an ideal FP probe QGGFL with improved quantum yield, high affinity and high specificity to α_1 -ARs. Then an "ELISA-like" strategy was adopted to amplify the FP signal and increase the measurement window and S/N ratio. Finally, based on the FP probe and the ELISA strategy, we established a safe and affordable FP-based screening method for α_1 -ARantagonists for the first time.

