A New Stereoselective Synthesis of Cyclopropanes Containing Quaternary Stereocentres *via* Organocatalytic Michael Addition to Vinyl Selenones

Francesca Marini,^{a,*} Silvia Sternativo,^a Francesca Del Verme,^a Lorenzo Testaferri,^a and Marcello Tiecco^a

 ^a Dipartimento di Chimica e Tecnologia del Farmaco, Sezione di Chimica Organica, Università degli Studi di Perugia, Via del Liceo 1, 06123 Perugia, Italy
 Fax: (+39) 075-585-5116; e-mail: marini@unipg.it

Received: April 3, 2009; Revised: June 12, 2009; Published online: August 5, 2009

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/adsc.200900222.

Abstract: A novel organocatalytic method for the stereoselective synthesis of highly substituted cyclopropanes is reported. The Michael adducts, generated through the addition of α -substituted cyanoacetates to easily accessible vinyl selenones catalyzed by a bifunctional ureidic catalyst, smoothly cyclize by intramolecular alkylation induced by a de-ethoxycarbonylation process. The *one-pot* sequence generates cyclopropanes bearing adjacent tertiary and quaternary stereocentres as single Z-isomers in moderate to high yields and good enantiomeric excesses.

Keywords: asymmetric organocatalysis; cyclopropanes; Michael addition; quaternary stereocenters; selenones

The cyclopropane ring is the basic structural motif of a wide range of natural products and biologically active compounds, such as conformationally restricted amino acids or peptides, enzyme inhibitors and therapeutic agents with antipsychotic, antifungal, antibacterial, antitumoral or antiviral activities.^[1] Furthermore, the three-membered ring, due to its strong angular strain and great ring-opening ability, is a recognized intermediate in the construction of complex molecular skeletons.^[2] On these grounds, the development of simple and efficient methods for the stereocontrolled preparation of variously substituted cyclopropanes remains an interesting task for synthetic chemists. In the last years great attention has been focused on the identification of new asymmetric methodologies based on readily available achiral starting materials and chiral catalysts. Thus, asymmetric Simmons-Smith reactions or cyclopropanations of olefins by metal-catalyzed decomposition of diazo compounds have been successfully explored.^[3] Very recently significant advances have also been made in the development of organocatalytic cyclopropanations based on Michael initiated ring closure reactions (MIRC).^[3,4] In these processes, the addition of an ylide or an enolate containing a good leaving group to an electrophilic alkene generates an anionic intermediate, which easily undergoes a ring closure reaction by intramolecular alkylation. During our studies on the asymmetric Michael addition of a-substituted cyanoacetates to vinyl selenones catalyzed by ureidic or thioureidic bifunctional catalysts, we observed that the diastereomeric mixture of Michael adducts 4aA and 5aA affords the cyclopropane 6aA as a single isomer by simple treatment with KCN in DMF (Scheme 1).^[5]

We explained this process as a Krapcho-type deethoxycarbonylation, followed by an intramolecular nucleophilic substitution of the selenonyl moiety by the enolate intermediate. The excellent leaving group properties of this species have been already exploited for other synthetically useful inter- or intramolecular substitutions.^[5,6] This first result prompted us to study the sequence in detail and develop a novel *one-pot*



Scheme 1. Formation of the cyclopropane 6aA.

© 2009 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim





Figure 1. Ureidic or thioureidic catalysts 1a-f.

procedure for the enantioselective preparation of substituted cyclopropanes containing adjacent all carbon quaternary and tertiary stereocentres starting from readily available vinyl selenones.^[7] Some catalytic approaches for the stereoselective construction of such cyclopropanes are available in the literature^[3,4f,k,8] including methods for the preparation of nitrile-substituted cyclopropanes,^[9] but only few of them employ organocatalysts.

Preliminary experiments were carried out in order to examine the addition step which is responsible of the enantioselectivity of the sequence. The commercial ethyl α -phenylcyanoacetate **2a** and the (E)-vinyl selenone $3A^{[6]}$ were used as the model substrates in the presence of the commercial **1f** or easily accessible 1a-e chiral urea or thiourea derivatives reported in Figure 1, which have recently emerged as efficient catalysts for a broad range of highly enantioselective Michael additions.^[10] These bifunctional catalysts, bearing an hydrogen-bond donor group besides a basic site on a chiral scaffold, accelerate the 1,4-additions and improve yields and stereoselectivities by simultaneous activation of both the Michael acceptor and the carbon nucleophile. The results of the experiments confirmed the efficiency of the ureidic catalyst 1a, which gave in toluene the best results in terms of yield, diastereo- and enantioselectivity (Table 1).

The stereoinduction is slightly affected by the ester size with a decrement of the enantioselectivity as the size of the ester increases (Table 1, entries 1, 5, 6). When the temperature was raised at 50 °C the reaction time was shortened at 48 h, albeit with a loss of selectivity (Table 1, entry 4). It is interesting to point out that an attempt to effect the addition of **2a** on the phenyl (*E*)-2-phenylethenyl sulfone, structurally related to **3A**, failed indicating a different behaviour between the sulphur^[11,12] and the selenium^[5] containing compounds.

After the optimization of the addition step, the ring formation was examined. The crude mixture of **4aA**

		$\begin{array}{ccc} NC & CO_2R & SeO_2Ph \\ Ph & Ph \\ 2a - c & 3A \end{array}$			1a - f 20% mol NC * CO ₂ R solvent, 4 Å MS Ph SeO ₂ Ph 4 and 5				
Entry	Catalyst		R	Solvent	Time [h]	Yield [%] ^[b]	4:5 ^[c]	<i>ee</i> [%] ^[d]	
1	1 a	2a	Et	toluene	90	94	85:15	80	
2	1 a	2a	Et	CH_2Cl_2	144	78	78:22	56	
3	1a	2a	Et	THF	144	73	74:26	54	
4	1a	2a	Et	toluene	48 ^[e]	99	82:18	64	
5	1a	2b	Bn	toluene	90	87	79:21	64	
6	1 a	2c	<i>i</i> -Pr	toluene	90	97	77:23	66	
7	1b	2a	Et	toluene	72	_	_	_	
8	1c	2a	Et	toluene	120	81	78:22	68 ^[f]	
9	1d	2a	Et	toluene	144	21	68:32	58 ^[f]	
10	1e	2a	Et	toluene	144	< 20	68:32	56 ^[f]	
11	1f	2a	Et	toluene	144	40	42:58	50 ^[f]	

Table 1. Selected results for the Michael addition of **2a–c** to vinyl selenone **3A**.^[a]

^[a] Catalysts **1a-f** (0.02 mmol, 20 mol%) and **3A** (0.1 mmol) were dissolved in the indicated solvent (0.4 mL), then 4 Å MS (20 mg) and **2a-c** (0.15 mmol, 1.5 equiv.) were added at room temperature.

^[b] Yields refer to the mixture of diastereoisomers **4** and **5**.

^[c] Determined by ¹H NMR analysis after passing through a plug of silica gel.

^[d] The *ee* values, determined by chiral HPLC analysis, refer to the adducts **4**. The minor diastereomers **5** have *ee* lower than 20%.

^[e] The reaction was carried out at 50 °C.

^[f] The opposite enantiomer was obtained.

asc.wiley-vch.de

© 2009 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

 Table 2. Optimization of the cyclization step.



Entry	Reaction Conditions ^[a]	Yield [%] ^[b]	ee [%] ^[c]
1	LiCl, HMPA, 80°C, 2 h	65	66
2	LiCl, toluene/HMPA 1:1, 100°C, 24 h	31 ^[d]	66
3	LiCl, DMSO, 160 °C, 2 h	[37] ^[e]	_
4	LiCl, toluene, 12-crown-4, 80°C, 24 h	[48] ^[e]	_
5	NaBr, DMSO, 100 °C, 5 h	[51] ^[e]	_
6	aq. NaOH, MeOH, r.t., 24 h	[f]	_
7	$EtONa/EtOH, 0$ °C \rightarrow r.t., 2 h	54	76
8	EtONa/EtOH, 0°C→r.t., 2 h	41	64 ^{g]}

[a] 1a (0.02 mmol, 20 mol%) and 3A (0.1 mmol) were dissolved in toluene (0.4 mL), then 4 Å MS (20 mg) and 2a (0.15 mmol, 1.5 equiv.) were added at room temperature. After 90 h the toluene (if necessary) was removed under reduced pressure and the indicated reagents and solvent were added.

^[b] Determined on isolated **6aA**.

- ^[c] The *ee* values were determined by chiral HPLC analysis.
 ^[d] Products of nucleophilic substitution of the selenonyl group by chloride ion were also formed.
- ^[e] The alkene **7** was recovered as the main product with the yield reported in parentheses.
- ^[f] A complex mixture of products was formed.
- ^[g] The opposite enantiomer was obtained by using catalyst **1c**.

and **5aA**, obtained by simple removal of the toluene under vacuum, was submitted to treatment with LiCl or NaBr in a polar aprotic solvent (Krapcho-type protocols)^[13] or with an aqueous solution of NaOH in MeOH^[14] (Table 2). An alternative treatment with NaOEt in EtOH was also attempted. Only in the presence of LiCl in HMPA (method A) and NaOEt in EtOH (method B) was the cyclopropane **6aA** formed in good yields.

The stereochemical assignment of $6aA^{[15]}$ allows us to assign the 3S absolute configuration at the tertiary stereocenter of the major adduct 4aA. Reasonably, the LiCl or the NaOEt, respectively, attack the ethyl or the carbonyl group of the ester moiety forming a nitrile enolate which promotes the 3-exo-tet ring closure reaction. The process is completely diastereoselective, giving the sole Z-isomer with good enantiomeric excesses (Table 2, entries 1 and 7). In the other conditions more complex reaction mixtures, containing products generated by intermolecular nucleophilic substitution of the selenonyl group or considerable amounts of the elimination product 7, were obtained. The cyclization with method B generates the cyclo-



Scheme 2. Cyclization of adducts 4dA and 5dA.

propane **6aA** with an enantiomeric excess which is almost identical to that of the major adduct **4aA** and seems not affected by the presence of the poorly enantioenriched minor adduct **5aA**. This observation suggests that under these conditions only **4aA** is quantitatively converted into the cyclic compound **6aA**.

Additional experiments performed on the chromatographically separated adducts, generated from cyanoacetate 2d and the vinyl selenone 3A, confirmed this hypothesis. In fact, the cyclization of the major adduct 4dA proceeds smoothly, while the conversion of the minor isomer 5dA generates 6dA in a very poor yield (28% yield) together with unidentified byproducts (Scheme 2). As demonstrated by the ¹H NMR analyses of the crude reaction mixtures in both cases the *E*-cyclopropane was not present.

The one-pot process, in which the addition, deethoxycarbonylation and cyclization steps are carried out sequentially and without isolation of the intermediates, was applied to a range of β -aryl-substituted vinyl selenones containing electron-withdrawing or electron-donating groups at different positions of the phenyl ring. Different α -aryl cyanoacetates were also used as the Michael donor. Methods A and B were both used for the cyclization. The cyclopropanes 6 were generated in moderate to good yields, complete diastereoselectivity and good enantioselectivity (Table 3 entries 1-8). The results obtained with EtONa in EtOH are particularly interesting. In fact, this strategy provide higher enantiomeric excesses and comparable yields than those with the Krapcho protocol which involves the hazardous HMPA. The β alkyl-substituted vinyl selenone 3G was also examined as the Michael acceptor in the presence of the α arylcyanoacetates 2a and 2f. The addition step proceeded smoothly and at a faster rate than in the presence of the corresponding aryl compounds. Only the treatment with LiCl in HMPA gave the Z-2-alkyl-substituted cyclopropanes 6aG and 6fG^[16] in high vields and moderate selectivity (Table 3, entries 9-11).

Further experiments demonstrated that the presence of an α -aryl group on the starting cyanoacetate **2** is essential for the de-ethoxycarbonylation of the adducts **4**. In fact, the reactions between the α -allylcyaTable 3. Formation of the cyclopropanes 6.

			NC CC	D ₂ Et +SeO ₂ R 3	[⊃] h <u>a) 1a (20 mol%), toluene, r.t.</u> <u>b)</u> method A or B		NC, Ar R		
Entry		Ar	R			Metho Yield [%] ^[c]		Method $B^{[b]}$ Yield [%] ^[c] ee [%] ^[c]	
1	2a	Ph	3A	Ph	(-)-6aA	65	66	54	76
2	2a	Ph	3B	p-MeO-C ₆ H ₄	(–)-6aB	64	66	40	74
3	2a	Ph	3C	p-Me-C ₆ H ₄	(–)-6aC	51	68	58	76
4	2a	Ph	3D	p-Cl-C ₆ H ₄	(–)-6aD	40	72	40	72
5	2a	Ph	3E	m-F-C ₆ H ₄	(–)-6aE	55	74	65	74
6	2a	Ph	3F	o -Me- C_6H_4	(–)-6aF	56	48	45	68
7	2d	p-Br-C ₆ H ₄	3A	Ph	(–)-6dA	52	61	50	70
8	2e	p-F-C ₆ H ₄	3D	p-Cl-C ₆ H ₄	(–)-6eD	42	74	66	66
9	2a	Ph	3G	C_6H_{13}	(–)-6aG	81	48	18	54
10	2a	Ph	3G	$C_{6}H_{13}$	(–)-6aG	90 ^[e]	54	_	_
11	2f	p-MeO-C ₆ H ₄	3G	$C_{6}H_{13}$	(-)-6fG	91	52	-	-

^[a] Method A: LiCl (2.5 equiv.) in HMPA.

^[b] Method B: EtONa (2.5 equiv.) in EtOH.

^[c] Determined on isolated compounds.

^[d] Determined by chiral HPLC analysis.

^[e] Reaction carried out at -20 °C for 144 h.

noacetate and **3A** under the experimental conditions described in Table 3 did not afford any cyclopropanation product.

The stereochemical assignment of the cyclopropanes **6aA** and **6fG**^[15,16] suggests that in the presence of **1a** the Michael addition occurs preferentially on the *Re* face of the β -substituted unsaturated selenones. Thus, the facial selectivity is opposite to that observed in similar reactions with Michael acceptors containing nitro or carbonyl groups and urea and thiourea-substituted 9-*epi*-dihydroquinine derivatives.^[17]

The presence of the non-planar selenonyl group should modify the spatial arrangement of the ternary complex formed by the catalyst, the Michael acceptor and the nucleophile. In conclusion, we have described the first organocatalytic synthesis of enantioenriched cyclopropanes starting from β-substituted vinyl selenones. The key step of this procedure is the Michael addition of α -aryl-substituted cyanoacetates to β -substituted vinyl selenones,^[18] followed by a de-ethoxycarbonylation and a ring closure reaction. In most cases, the reactions can be carried out with comparable or even better results using EtONa in EtOH which avoids the use of HMPA with advantages for health and environment. The practical one-pot sequence, which occurs with complete diastereoselectivity and good enantioselectivity, is based on the peculiar properties of the selenonyl moiety which acts both as an electron-withdrawing group during the addition step and as a leaving group during the cyclization. Our procedure nicely complements the recent rhodium-catalyzed cyclopropanation of electron-rich olefins by 2-diazo-2-phenylacetonitrile, which selectively affords enantioenriched *E*-nitrile-substituted cyclopropanes.^[9a] Moreover, it expands the use of privileged organocatalysts in the field of selenium chemistry and opens attractive prospects for up to now scarcely explored organocatalytic selenium-based transformations.

Experimental Section

General Procedure for the *One-Pot* Synthesis of Enantioenriched Z-Cyclopropanes 6

In an ordinary vial equipped with a Teflon-coated stir bar, catalyst **1a** (0.04 mmol, 20 mol%) and the selenones **3A–F**^[19] or **3G**^[20] (0.2 mmol) were dissolved in undistilled toluene (0.8 mL) with 20 mg of 4 Å MS. The α -substituted cyanoace-tate **2a**, **d–f** (0.3 mmol, 1.5 equiv.) were added at room temperature (15–18 °C) and the resulting mixtures were stirred in air for 48–144 h. The solvent was removed under reduced pressure and the crude reaction mixtures were submitted to cyclization following method A or B.

Method A: the crude reaction mixtures were dissolved in HMPA (1.5 mL). Then LiCl (0.5 mmol) was added and the mixtures were stirred at 80 °C for 2–4 h. The reactions were quenched with water (20 mL) and extracted with diethyl ether $(3 \times 10 \text{ mL})$. The organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under vacuum. The residues were purified by flash chromatography to yield the cyclopropanes **6**.

Method B: the crude residues were dissolved in EtOH (1 mL) and a solution of EtONa (0.5 mmol) in EtOH

(1 mL) was added at 0 °C. The resulting solutions were allowed to warm to room temperature and stirred for 1–4 h. The reaction mixtures were quenched with water (20 mL) and extracted with diethyl ether (3×10 mL). The organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under vacuum. The residues were purified by flash chromatography to yield the cyclopropanes **6**.

Acknowledgements

Financial support from MIUR, National Projects PRIN 2007, University of Perugia and Consorzio CINMPIS, Bari is gratefully acknowledged.

References

- a) Small ring compounds in organic synthesis VI, in: Top. Curr. Chem. (Ed.: A. de Meijere), Springer, Berlin, 2000, vol. 207; b) J. Salaun, Curr. Med. Chem. 1995, 2, 511; c) W. A. Donaldson, Tetrahedron 2001, 57, 8589; d) R. Faust, Angew. Chem. 2001, 113, 2312; Angew. Chem. Int. Ed. 2001, 40, 2251; e) A. Reichelt, S. F. Martin, Acc. Chem. Res. 2006, 39, 433.
- [2] For reviews on cyclopropanes as intermediates in organic synthesis see: a) H.-U. Reissig, R. Zimmer, *Chem. Rev.* 2003, 103, 1151; b) H. N. C. Wong, M. Y. Hon, C. W. Tse, Y. C. Yip, J. Tanko, T. Hudlicky, *Chem. Rev.* 1989, 89, 165; c) M. Rubin, M. Rubina, V. Gevorgyan, *Chem. Rev.* 2007, 107, 3117.
- [3] For recent reviews see: a) H. Pellissier Tetrahedron 2008, 64, 7041; b) H. Lebel, J.-F. Marcoux, C. Molinaro, A. B. Charette, Chem. Rev. 2003, 103, 977; c) R. C. Hartley, S. T. Caldwell, J. Chem. Soc. Perkin Trans. 1 2000, 477.
- [4] For some recent examples of asymmetric MIRC reactions see: a) C. C. C. Johansson, N. Bremeyer, S. V. Ley, D. R. Owen, S. C. Smith, M. J. Gaunt, Angew. Chem. 2006, 118, 6170; Angew. Chem. Int. Ed. 2006, 45, 6024; b) R. K. Kunz, D. W. C. MacMillan, J. Am. Chem. Soc. 2005, 127, 3240; c) S. H. McCooey, T. McCabe, S. J. Connon, J. Org. Chem. 2006, 71, 7494; d) H. M. Hansen, D. A. Longbottom, S. V. Ley, Chem. Commun. 2006, 4838; e) H. Xie, L. Zu, H. Li, J. Wang, W. J. Wang, J. Am. Chem. Soc. 2007, 129, 10886; f) R. Rios, H. Sunden, J. Vesely, G.-L. Zhao, P. Dziedzic, A. Cordova, Adv. Synth. Catal. 2007, 349, 1028; g) A. Hartikka, A. T. Slosarczyk, P. I. Arvidsson, Tetrahedron: Asymmetry 2007, 18, 1403; h) J. Vesely, G.-L. Zhao, A. Bartoszewicz, A. Cordova, Tetrahedron Lett. 2008, 49, 4209; i) V. Wascholowski, H. M. Hansen, D. A. Longbottom, S. V. Ley, Synthesis 2008, 1269; j) I. Ibrahem, G.-L. Zhao, R. Rios, J. Vesely, H. Sunden, P. Dziedzic, A. Cordova, Chem. Eur. J. 2008, 14, 7867; k) J. Lv, J. Zhang, Z. Lin, Y. Wang, Chem. Eur. J. 2009, 15, 972; 1) Y. Xuan, S. Nie, L. Dong, J. Zhang, M. Yan, Org. Lett. 2009, 11, 1583.
- [5] F. Marini, S. Sternativo, F. Del Verme, L. Testaferri, M. Tiecco, Adv. Synth. Catal. 2009, 351, 103.

- [6] a) M. Tiecco, A. Carlone, S. Sternativo, F. Marini, G. Bartoli, P. Melchiorre, Angew. Chem. 2007, 119, 7006; Angew. Chem. Int. Ed. 2007, 46, 6882; b) M. Tiecco, L. Testaferri, A. Temperini, R. Terlizzi, L. Bagnoli, F. Marini, C. Santi, Org. Biomol. Chem, 2007, 5, 3510; c) M. Tiecco, L. Testaferri, L. Bagnoli, G. Scarponi, A. Temperini, F. Marini, C. Santi, Tetrahedron: Asymmetry 2007, 18, 2758; d) M. Tiecco, L. Testaferri, A. Temperini, L. Bagnoli, F. Marini, C. Santi, Chem. Eur. J. 2004, 10, 1752; e) V. R. Ward, M. A. Cooper, A. D. Ward, J. Chem. Soc. Perkin Trans. 1 2001, 944; f) M. A. Cooper, A. D. Ward, Aust. J. Chem. 1997, 50, 181; g) A. Toshimitsu, C. Hirosawa, S. Tanimoto, S. Uemura, Tetrahedron Lett. 1992, 33, 4017; h) A. Krief, W. Dumont, J.-N. Denis, J. Chem. Soc. Chem. Commun. 1985, 571; i) A. Krief, W. Dumont, A. F. De Mahieu, Tetrahedron Lett. 1988, 29, 3269; j) M. Tiecco, D. Chianelli, L. Testaferri, M. Tingoli, D. Bartoli, Tetrahedron 1986, 42, 4889; k) M. Tiecco, D. Chianelli, L. Testaferri, M. Tingoli, D. Bartoli, Tetrahedron 1986, 42, 4897; l) I. Kuwajima R. Ando, T. Sugawara, Tetrahedron Lett. 1983, 24, 4429; m) R. Ando, T. Sugawara, M. Shimizu, I. Kuwajima, Bull. Chem. Soc. Jpn. 1984, 57, 2897; n) selenonium groups can also be easily displaced by nucleophiles. This property has been employed to develop a one-pot cyclopropanation reaction starting from vinylselenonium salts: o) Y. Watanabe; Y. Ueno, T. Toru, Bull. Chem. Soc. Jpn. 1993, 66, 2042.
- [7] Vinyl selenones are prepared by oxidation of the corresponding selenides (see Supporting Information). For an update review on the synthesis of vinyl selenides see: G. Perin, E. J. Lenardao, R. G. Jacob, R. B. Panatieri, *Chem. Rev.* 2009, 109, 1277.
- [8] a) J. R. Denton, H. M. L. Davies, Org. Lett. 2009, 11, 787; b) H. M. L. Davies, G. H. Lee, Org. Lett. 2004, 6, 2117.
- [9] a) J. R. Denton, K. Cheng, H. M. L. Davies *Chem. Commun.* 2008, 1238; b) S. Arai, K. Nakayama, T. Ishida, T. Shioiri, *Tetrahedron Lett.* 1999, 40, 4215.
- [10] a) A. G. Doyle, E. N. Jacobsen, Chem. Rev. 2007, 107, 5713; b) M. S. Taylor, E. N. Jacobsen, Angew. Chem. 2006, 118, 1550; Angew. Chem. Int. Ed. 2006, 45, 1520; c) S. J. Connon, Chem. Eur. J. 2006, 12, 5418; d) S. J. Connon, Chem. Commun. 2008, 2499; e) S. J. Connon Synlett 2009, 354.
- [11] During the preparation of this manuscript, the asymmetric conjugate addition of carbon nucleophiles to β -substituted vinyl sulfones catalyzed by C6'-OH *Cinchona* alkaloids has appeared in the literature. The authors reported that the success of the addition is strictly dependent by the presence of the strong electron-withdrawing group CF₃SO₂ and that the phenyl (*E*)-2-phenylethenyl sulfone is not reactive. See: H. Li, J. Song, L. Deng *Tetrahedron* **2009**, *65*, 3139. We tested one of the Deng's C6'-OH *Cinchona* alkaloid catalysts on **3A** obtaining the adducts **4a/5a** in low yield and poor diastereo- and enantioselectivity.
- [12] For Cinchona alkaloids-catalyzed asymmetric conjugate addition of carbon nucleophiles to β-unsubstituted vinyl sulfones see: a) T.-Y. Liu, J. Long, B.-J. Li, L. Jiang, R. Li, Y. Wu, L.-S. Ding, Y.-C. Chen, Org.

Biomol. Chem. **2006**, *4*, 2097; b) H. Li, J. Song, X. Liu, L. Deng, *J. Am. Chem. Soc.* **2005**, *127*, 8948.

- [13] a) A. P. Krapcho, Synthesis 1982, 805; b) A. P. Krapcho, Synthesis 1982, 893.
- [14] C. Cheng, H. Lu, F. Lee, Eur. J. Med. Chem. 1991, 26, 125.
- [15] The relative configuration of 6aA was assigned by comparison of the ¹H NMR spectrum with that reported in the literature (ref.^[15a]). The absolute configuration was assigned as 1*S*,2*S* by its (-) optical rotation typical for the (2*S*)-isomers of 2-phenyl substituted cyclopropane-carboxylic derivatives. See: a) W. von E. Doering, L. R. Robertson, E. E. Ewing, *J. Org. Chem.* 1983, 48, 4280; b) A. B. Chmurny, D. J. Cram, *J. Am. Chem. Soc.* 1973, 95, 4237.
- [16] The relative and absolute configuration (1*S*,2*R*) of compound **6fG** was assigned by comparison of its ¹H NMR spectrum and specific optical rotation, respectively, with those reported in the literature. See: T. Kusumoto, A. Nakayama, K. Sato, T. Hiyama, S. Takehara, T. Shoji, M. Osawa, T. Kuriyama, K. Nakamura, T. Fujisawa, *Tetrahedron Lett.* **1991**, *32*, 939.
- [17] a) S. H. McCooey, S. J. Connon, Angew. Chem. 2005, 117, 6525; Angew. Chem. Int. Ed. 2005, 44, 6367; b) J.

Wang, H. Li, L. Zu, W. Jiang, H. Xie, W. Duan, W. Wang, *J. Am. Chem. Soc.* **2006**, *128*, 12652; c) C. Gu, L. Liu, Y. Sui, J.-L. Zhao, D. Wang, Y.-J. Chen, *Tetrahedron: Asymmetry* **2007**, 18, 455.

- [18] Simple cyanoacetates or other unsubstituted activated methylenic compounds add to β -substituted vinyl selenones generating directly 2-substituted cyclopropanecarboxylic esters through an alternative cascade process which involves a Michael addition, a proton exchange and a cyclization by displacement of the PhSeO₂ group.^[6j,l,m] The organocatalytic version of this process merits investigations. These are currently under way in our laboratory.
- [19] According to Tiecco's procedure (ref.^[6]), the (*E*)-vinyl selenones **3A–F** were prepared from the corresponding commercial or easily available vinyl bromides. See:
 a) H.-W. You, . K.-J. Lee *Synlett* **2001** 105; b) C. Kuang, H. Senboku, M. Tokuda, *Tetrahedron* **2005**, *61*, 637.
- [20] The vinyl selenide precursor of the (E)-vinyl selenone **3G** was prepared from commercial 1-octene and PhSeBr following a literature procedure: S. Raucher, M. R. Hansen, M. A. Colter, J. Org. Chem. 1978, 43, 4885.