

Intramolecular Stetter cyclisation of Morita–Baylis–Hillman adducts: a versatile approach towards bicycloenediones†

Pierre Wasnaire, Thierry de Merode and István E. Markó*

Received (in Cambridge, UK) 10th July 2007, Accepted 22nd August 2007

First published as an Advance Article on the web 31st August 2007

DOI: 10.1039/b710492h

Bicyclic enediones of various sizes can be efficiently assembled by intramolecular Stetter cyclisation of readily available Morita–Baylis–Hillman adducts.

Bicyclic structures are ubiquitous subunits in a wide variety of natural products such as coronatine¹ and guaiol² (Fig. 1). The occurrence of these substructures has stimulated the development of numerous and elegant procedures for their efficient construction.³ As part of an ongoing research programme directed towards the rapid and flexible synthesis of some of these naturally occurring compounds, we have recently described a method for the assembly of substituted hydrindanones and decalones based upon the reductive cyclisation of Morita–Baylis–Hillman adducts.⁴

In this communication, we wish to report on the discovery and development of a simple methodology for the rapid construction of various bicyclic enediones that hinges upon two key steps: a Morita–Baylis–Hillman coupling,⁵ followed by an intramolecular Stetter reaction.⁶ As depicted in Fig. 2, cleavage of the C-4/C-5 bond of bicyclic enedione **3**, accompanied by the concomitant insertion of an oxygenated function at C-8, reveals the Morita–Baylis–Hillman adduct **4**, itself readily available from enone **5** and aldehyde **6**.

Our approach thus began with the union of cyclic enones **5a–d** with 4-pentenal **6a** (Scheme 1). After some experimentation, it was found that a slight modification of the procedure reported by Yamada and Ikegami,⁷ using *n*-Bu₃P and *rac*-binol, promoted the coupling of 2-cyclopentenone **5a** and 2-cyclohexenone **5b** in excellent yields. In the case of 2-cycloheptenone **5c** and 2-cyclooctenone **5d**, the Nozaki Et₂AlI-mediated protocol was required⁸ and the desired adducts **7c–d** were obtained in satisfactory yields. Alcohols **7a–d** were then subjected to

acetylation, under Yamamoto's conditions,⁹ providing the β-acetoxy enones **8a–d** in good to essentially quantitative yields. Finally, chemoselective ozonolysis of the terminal alkene of **8a–d**, in the presence of Sudan red as an indicator,¹⁰ produced the key cyclisation precursors **9a–d** in 75–95% yield (Scheme 1).

With the desired precursors in hand, we next turned our attention to the crucial cyclisation step: the intramolecular ring closure of an aldehyde onto a cyclic enone. To perform this reaction, the aldehyde had to be transformed initially into an acylanion equivalent. Among several possibilities, the Stetter reaction which exploits this umpolung reactivity¹¹ and which has recently been the subject of renewed interest attracted our attention.¹² Although the Stetter reaction can be carried out sometimes with good yields and selectivities, its synthetic scope still remains to be explored.¹³

Our studies began with an examination of the intramolecular Stetter cyclisation of the cyclohexenone-aldehyde derivative **9b** (Table 1). Under standard Stetter conditions (0.1 equiv. of thiazolium salt **10a** and 0.6 equiv. of Et₃N in EtOH at reflux), the desired bicyclic enedione **11b** could be obtained in 50% yield (Entry 1). Several reaction parameters were next varied in order to optimise the yields; some selected examples are shown in Table 1. When 1 equiv. of commercially available thiazolium salt **10a** and 1.2 equiv. of Et₃N were employed, in EtOH at reflux, the enedione **11b** could be isolated in a satisfactory 66% yield (Entry 5).

Having delineated suitable conditions for the intramolecular Stetter reaction, we next applied them to the ring closure of β-acetoxyenones **9a–d** (Table 2). When cyclopentenone **9a** was submitted to these conditions, the corresponding bicyclic enedione **11a** was obtained in 50% yield (Entry 1).¹⁴ Increasing the ring size of the substrates led to improved yields. Cycloheptenone and cyclooctenone derivatives **9c–d** afforded the desired enediones **11c–d** in an excellent 80% yield. (Entries 3 and 4).

At this stage, the viability of our approach had been demonstrated and it became evident that Morita–Baylis–Hillman adducts were suitable intermediates to access various bicyclic enediones. In order to extend our methodology, it was decided to synthesise a more challenging adduct: the perhydroazulene core **15**. To achieve this objective, cyclopentenone **5a** was coupled to




Fig. 1 Selected natural products containing a bicyclic core.

Université catholique de Louvain, Département de Chimie, 1 Place Louis Pasteur, B-1348, Louvain-la-Neuve, Belgium.
E-mail: marko@chim.ucl.ac.be; Fax: +32 10/472788;
Tel: +32 10/478773

† Electronic supplementary information (ESI) available: Experimental details. See DOI: 10.1039/b710492h





Fig. 2 Proposed retrosynthetic analysis of bicyclic enediones.



Scheme 1 Assembly of the cyclisation precursors. *Reagents and conditions:* (i) 3 equiv. **5a** or **5b**, 0.6 equiv. n -Bu₃P, 0.3 equiv. binol, THF, rt; (ii) **5c** or **5d**, 1.2 equiv. **6a**, 1.4 equiv. Et₂AlI, CH₂Cl₂, 0 °C; (iii) 1.5 equiv. Ac₂O, Sc(OTf)₃ cat., CH₃CN, rt; (iv) O₃, Sudan red 7B, CH₂Cl₂ then Ph₃P, −78 °C to rt.

Table 1 Optimisation of the intramolecular Stetter reaction

<p>9b catalyst 10 base solvent, T° 11b</p> <p>10a: R⁴ = CH₂Ph, X = Cl 10b: R⁴ = C₂H₅, X = Br</p>					
Entry	Catalyst (equiv.)	Base (equiv.)	Solvent	T (°C)	Yield ^a (%)
1	10a (0.1)	Et ₃ N (0.6)	EtOH	78	50
2	10a (0.1)	Et ₃ N (0.6)	CH ₂ Cl ₂	40	20
3	10a (0.1)	Et ₃ N (0.6)	dioxane	80	32
4	10a (0.2)	Et ₃ N (0.7)	<i>t</i> -BuOH	80	55
5	10a (1.0)	Et ₃ N (1.2)	EtOH	78	66
6	10b (1.0)	Et ₃ N (1.2)	EtOH	78	47
7	10b (1.0)	DBU (1.2)	EtOH	78	24

^a Yields refer to pure, isolated products.

5-hexenal **6b**, under Ikegami's conditions, affording hydroxyenone **12** which was obtained in 81% yield. Acetylation of the hydroxyl group, followed by selective ozonolysis of the terminal alkene, generated the key intermediate **14** in 64% overall yield (Scheme 2). When submitted to our Stetter conditions, substrate **14** produced the corresponding perhydroazulenedione derivative **15** in a satisfactory 46% yield. To the best of our knowledge, this is the first example of the construction of a 7-membered ring using the Stetter reaction.


Though enediones are potent functionalities, their reactivity has been little examined.¹⁵ As a relevant preliminary study, we elected to reduce the conjugated C–C double bond using McMurry's procedure,¹⁶ *i.e.* TiCl₃ in acidic aqueous solution. In the presence of 4 equiv. of aqueous TiCl₃ in acetone, enedione **11a** afforded smoothly the corresponding *cis*-hydrindanone **16a**, as a single

Table 2 Intramolecular Stetter reaction of Morita–Baylis–Hillman adducts

Entry ^a	Substrate	Product	Yield ^b (%)
1	<p>9a</p>	<p>11a</p>	50
2	<p>9b</p>	<p>11b</p>	66
3	<p>9c</p>	<p>11c</p>	80
4	<p>9d</p>	<p>11d</p>	80

^a All reactions were performed in the presence of 1 equiv. **10a** and 1.2 equiv. of Et₃N, in ethanol for 2.5 h at reflux. ^b Yields refer to pure, isolated products.

diastereoisomer, in 85% yield (Table 3).¹⁷ Applying the same conditions to the prochiral enedione **11b** led to the *trans*-decalone **16b** in essentially quantitative yield. Similarly bicyclo[6.4.0]enedione **11d** was reduced stereoselectively to the *trans*-diketone **16d** in 86% yield.



Scheme 2 Preparation of a perhydroazulene. *Reagents and conditions:* (i) 3 equiv. **5a**, 0.6 equiv. n -Bu₃P, 0.3 equiv. binol, THF, rt; (ii) 1.5 equiv. Ac₂O, Sc(OTf)₃ cat., CH₃CN, rt; (iii) O₃, Sudan red 7B, CH₂Cl₂ then Ph₃P, −78 °C to rt; (iv) 1 equiv. **10a**, 1.2 equiv. Et₃N, EtOH, reflux, 2.5 h.

Table 3 Reduction of enediones

Entry	Substrate	Product	Yield ^a (%)
1			85
2			97
3			86

^a Yields refer to pure, isolated products.

In summary, we have developed a short and connective synthesis of bicyclic enediones. This new methodology involves two key steps: a Morita–Baylis–Hillman coupling and an intramolecular Stetter reaction. Various bicyclic structures can be readily assembled *via* this versatile protocol. Moreover, the Stetter reaction has been shown to be a powerful synthetic tool that led, not only to 6-*endo*, but also to 7-*endo-trig* cyclisations. Current efforts are now aimed at delineating the full scope of this methodology, examining the reactivity of enediones and using these substrates in the total synthesis of relevant natural products. The results of these investigations will be reported in due course.

Financial support for this work by the Université catholique de Louvain, the Fonds pour la Recherche dans l'Industrie et l'Agriculture (FRIA, studentships to P.W.) and Merck Sharp and Dohme (Merck Academic Development Award to I.E.M.) is gratefully acknowledged.

Notes and references

- 1 A. Ichihara, K. Shiraishi, H. Sato, S. Sakamura, K. Nishiyama, R. Sakai, A. Furusaki and T. Matsumoto, *J. Am. Chem. Soc.*, 1979, **99**, 636.
- 2 H. Minato, *Tetrahedron Lett.*, 1961, **2**, 280.
- 3 For general reviews, see: V. Singh, S. R. Iyer and S. Pal, *Tetrahedron*, 2005, **61**, 1917; P. Jankowski, S. Marczak and J. Wicha, *Tetrahedron*, 1998, **54**, 12071.
- 4 P. Wasnaire, M. Wiaux, R. Touillaux and I. E. Markó, *Tetrahedron Lett.*, 2006, **47**, 985.
- 5 K. Morita, Z. Suzuki and H. Hirose, *Bull. Chem. Soc. Jpn.*, 1968, **41**, 2815; A. B. Baylis and M. E. D. Hillman, *German Patent 2155113*, 1972. For a general review, see: D. Basavaiah, A. J. Rao and T. Satyanarayana, *Chem. Rev.*, 2003, **103**, 811. For selected examples, see reference 4 and those cited therein.
- 6 For some leading references, see: Q. Liu and T. Rovis, *Org. Process Res. Dev.*, 2007, **11**, 598; W. Schrader, P. P. Handayani, C. Burstein and F. Glorius, *Chem. Commun.*, 2007, 716; T. E. Reynolds, C. A. Stern and K. A. Scheidt, *Org. Lett.*, 2007, **9**, 2581; O. L. Epstein and T. Rovis, *Tetrahedron*, 2006, **62**, 11477; J. R. de Alaniz and T. Rovis, *J. Am. Chem. Soc.*, 2005, **127**, 6284; M. Christmann, *Angew. Chem., Int. Ed.*, 2005, **44**, 2632; A. E. Mattson, A. R. Bharadwaj and K. A. Scheidt, *J. Am. Chem. Soc.*, 2004, **126**, 2314; D. Enders and T. Balensiefer, *Acc. Chem. Res.*, 2004, **37**, 534; D. Enders and O. Niemeier, *Synlett*, 2004, 2111. For early references, see: H. Stetter and M. Schreckenberger, *Angew. Chem., Int. Ed. Engl.*, 1973, **12**, 81; H. Stetter, *Angew. Chem., Int. Ed. Engl.*, 1976, **15**, 639. For general reviews, see: H. Stetter and H. Kuhlmann, *Org. React.*, 1991, **40**, 407.
- 7 Y. M. A. Yamada and S. Ikegami, *Tetrahedron Lett.*, 2000, **41**, 2165.
- 8 A. Itoh, S. Ozawa, K. Oshima and H. Nozaki, *Bull. Chem. Soc. Jpn.*, 1981, **54**, 274; W. Pei, H. Wei and G. Li, *Chem. Commun.*, 2002, 2412.
- 9 K. Ishihara, M. Kubota, H. Kurihara and H. Yamamoto, *J. Org. Chem.*, 1996, **61**, 4560.
- 10 T. Veysoglu, L. A. Mitscher and J. K. Swayze, *Synthesis*, 1980, 807.
- 11 D. Seebach, *Angew. Chem., Int. Ed. Engl.*, 1979, **18**, 239.
- 12 For general reviews, see: M. Christmann, *Angew. Chem., Int. Ed.*, 2005, **44**, 2632; D. Enders and T. Balensiefer, *Acc. Chem. Res.*, 2004, **37**, 534; J. S. Johnson, *Angew. Chem., Int. Ed.*, 2004, **44**, 2632. For selected recent examples, see: J. L. Moore, M. S. Kerr and T. Rovis, *Tetrahedron*, 2006, **62**, 11478 and references cited therein; Z. Zhou, F. Ji, M. Cao and G. Yang, *Adv. Synth. Catal.*, 2006, **348**, 1826; T. Nakamura, O. Hara, T. Tamura, K. Makino and Y. Hamada, *Synlett*, 2005, 155; D. Enders, K. Breuer and J. H. Teles, *Helv. Chim. Acta*, 1996, **79**, 1899.
- 13 For applications to total synthesis, see: P. E. Harrington and M. A. Tius, *J. Am. Chem. Soc.*, 2001, **123**, 8509; B. M. Trost, C. D. Shuey, F. DiNinno and S. S. McElvain, *J. Am. Chem. Soc.*, 1979, **101**, 1284.
- 14 For **11a**, increasing the reaction time resulted in decreased yield.
- 15 See for examples: B. A. Pearlman, A. G. Padilla, J. T. Hach, J. L. Havens and M. D. Pillai, *Org. Lett.*, 2006, **8**, 2111; S. El Sheikh, N. Kausch, J. Lex, J.-M. Neudoerfl and H.-G. Schmaltz, *Synlett*, 2006, 1527; H. Asahara, E. Mochizuki and T. Oschima, *Tetrahedron Lett.*, 2006, **47**, 7881; K. Yamamoto, M. F. Hentemann, J. G. Allen and S. J. Danishefsky, *Chem.-Eur. J.*, 2003, **9**, 3242; J. Yu and E. J. Corey, *J. Am. Chem. Soc.*, 2003, **125**, 3232 and references cited therein; C. Liu and D. J. Burnell, *J. Org. Chem.*, 1997, **62**, 3683; W. Kroutil, M. E. Lastera-Sánchez, S. J. Maddrell, P. Mayon, P. Morgan, S. M. Roberts, S. R. Thornton, C. J. Todd and M. Tüter, *J. Chem. Soc., Perkin Trans. 1*, 1996, 2837.
- 16 L. C. Blaszcak and J. E. McMurry, *J. Org. Chem.*, 1974, **39**, 258.
- 17 A 24 h reaction time is necessary to complete the conversion of **11a** into the desired diketone **16a** due to the formation of a transient and unexpected 5-hydroxyenone intermediate.