This article was downloaded by: [Duke University Libraries] On: 28 September 2014, At: 12:43 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

http://www.tandfonline.com/loi/uopp20

A SIMPLE PROCEDURE FOR THE PREPARATION OF 6-CHLORO-3-ACETOXY-1-ACETYLINDOLE

Alexander Balbuzano-Deus^a, Juan C. Rodríguez-Domínguez^a, Anaís Fernández-Villalobo^a, Miguel López-López^a & Gilbert Kirsch^b ^a Departamento de Química, Centro de Química Farmacéutica, Calle 200 y 21 Atabey, Playa, CP, 11600, Ciudad de la Habana, CUBA ^b Laboratoire d'Ingénierie Moléculaire et Biochimie Pharmacologique 1 Boulevard Arago, 57070, Metz, FRANCE Published online: 11 Feb 2009.

To cite this article: Alexander Balbuzano-Deus , Juan C. Rodríguez-Domínguez , Anaís Fernández-Villalobo , Miguel López-López & Gilbert Kirsch (2006) A SIMPLE PROCEDURE FOR THE PREPARATION OF 6-CHLORO-3-ACETOXY-1-ACETYLINDOLE, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 38:1, 87-88

To link to this article: http://dx.doi.org/10.1080/00304940609355983

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>

OPPI BRIEFS

A SIMPLE PROCEDURE FOR THE PREPARATION OF 6-CHLORO-3-ACETOXY-1-ACETYLINDOLE

| Submitted by | Alexander Balbuzano-Deus [†] , Juan C. Rodríguez-Domínguez [†] , |
|--------------|--|
| (10/24/05) | Anaís Fernández-Villalobo [†] , Miguel López-López [†] and Gilbert Kirsch ^{*††} |

† Departamento de Química, Centro de Química Farmacéutica, Calle 200 y 21 Atabey, Playa, CP: 11600, Ciudad de la Habana, CUBA.
†† Laboratoire d'Ingénierie Moléculaire et Biochimie Pharmacologique 1 Boulevard Arago, 57070, Metz, FRANCE e-Mail: kirsch@sciences.univ-metz.fr.

The 6-chloro-3-acetoxy-1-acetylindole (3) moiety is present either as aglycon or as a part of different chromogenic compounds which are very useful for the quality control of water and food.^{1,2} The literature procedure³ for the preparation of the title compound starts from the commercially available 5-chloro-2-methylaniline and involves five steps. A long reaction time is required for the aminoalkylation and the overall yield of the product is only 12%. In order to simplify the process, we have developed a two-step procedure using Ullman's method,⁴ thus shortening the reaction time and improving the overall yield to about 38% Each step of the procedure was carried out with reproducible results and in excellent purity of the intermediate and of the final product.



EXPERIMENTAL SECTION

Thin layer chromatography (TLC) were performed on pre-coated plates of silica gel GF-254 (Merck) with CHCl₃:AcOEt:AcOH (8:6:1) as the solvent system. The tlc plates were visualized by means of a CAMAG UV-Vis lamp with a wavelenght of 254 nm. The ¹H and ¹³C NMR spectra were recorded on a Bruker AC 250F spectrometer in deuterated dimethyl- sulfoxide as solvent and TMS as an internal standard. Mps were determined on a Gallen-kamp capillary apparatus with a system of measurement and temperature control and are uncorrected.

^{© 2006} by Organic Preparations and Procedures Inc.

N-(2-Carboxy-5-chlorophenyl)glycine (2).- To a suspension of 10 g (52 mmol) of 2,4dichlorobenzoic acid, 9.95 g (133 mmol) of glycine and 0.40 g (6.3 mmol) of copper powder in 70 mL of dimethylformamide, was slowly added 21.7 g (157 mmol) of potassium carbonate. The mixture was heated at reflux with a good stirring for 6 h. When all the 2,4-dichlorobenzoic acid had been consumed, the mixture was poured over 70 mL of cold 6M hydrochloric acid and stirred for 30 min. The precipitate obtained was collected and washed with cold water until neutral pH. The solid was dried at 50°C to constant weight affording 10.1 g, (84%) of product as colorless crystals, mp. 225°C (dec.), *lit.*⁵ 228°C; mp⁵ of monosodium salt 278-280°C.

¹H NMR: δ 7.88 (d, 1H, H6, J_{6,4} = 8.5 Hz), 6.62 (dd, 1H, H4 J_{4,6} = 8.5 Hz, J_{4,3} = 1.9 Hz), 6.53 (d, 1H, H3 J_{1,3} = 1.9 Hz), 3.97 (s, 2H, CH₂).

¹³C NMR: δ (171.6 (CO), 169.6 (CO), 150.3 (C), 140.3 (C), 133.3 (CH), 115.3 (CH), 110.5 (CH), 109.2 (C).

6-Chloro-3-acetoxy-1-acetylindole (3).- A mixture of 10.5 g (45.7 mmol) of *N*-(2-carboxy-5-chlorophenyl)glycine, 70 mL of acetic anhydride and 14.4 g (175 mmol) of dry sodium acetate were heated at reflux. When the gas evolution had ceased, the mixture while still hot was poured into a beaker and left to cool overnight at 0°C. The precipitate was collected and poured into 70 mL of ice water and stirred for 1 h, collected again and recrystallized from ethyl acetate to yield 4.8 g (45%) of the product as a yellow solid, mp 111-113°C, *lit.*⁶ 112-113°C.

¹H NMR: $\delta 8.34$ (d, 1H, H7, J_{7,5} = 1.9 Hz); 7.90 (s, 1H, H2), 7.52 (d, 1H, H4, J_{4,5} = 8.5 Hz), 7.34 (dd, 1H, H5, J_{5,7} = 1.9 Hz, J_{5,4} = 8.5 Hz), 2.61 (s, 3H, CH₃, AcN); 2.28 (s, 3H, CH₃, AcO).

¹³C NMR: δ 169.36 (C=O, Ac), 167.95 (C=O, AcO), 133.02 (C9), 132.57 (C6), 130.08 (C5), 123.56 (C8), 122.27 (C4), 119.06 (C7), 116.33 (C2), 115.59 (CH), 23.36 (C, CH₃, AcN), 20.25 (C, CH₃, AcO).

REFERENCES

- R. D. Gonzalez, L. M. Tamagnini, P. D. Olmos and G. B. de Sousa, *Food Microbiology*, 20, 601 (2003).
- 2. P. W. Sadler and R. L. Warren, J. Am. Chem. Soc., 78, 1251 (1956).
- J. N. Roth and W. J. Ferguson, US Patent 5,210,022 (1993); Chem. Abstr., 119, 90788 (1993).
- R. F. Pellón, A. Estévez-Braun, M. L. Docampo, A. Martin and A. G. Ravelo, Synlett, 1606 (2005).
- 5. M. Heller and A. Hessel, J. prakt. Chem., [2], 120, 73 (1929).
- 6. S. J. Holt and P. W. Sadler, Proc. Roy. Soc. [B], 148, 481 (1958).