

Published on Web 09/12/2006

Gold(I) Catalyzed Isomerization of 5-en-2-yn-1-yl Acetates: An Efficient Access to Acetoxy Bicyclo[3.1.0]hexenes and 2-Cycloalken-1-ones

Andrea Buzas and Fabien Gagosz*

Laboratoire de Synthèse Organique, UMR 7652 CNRS/Ecole Polytechnique, 91128 Palaiseau, France Received June 15, 2006; E-mail: gagosz@dcso.polytechnique.fr

Gold(I) complexes have emerged as efficient and mild catalysts for the activation of alkynes toward addition by a variety of nucleophiles.¹ The potential of such catalysts has been highlighted by numerous studies related to the conversion of various enynes into cycloisomerized products.²

In this respect and by analogy with the isomerization of 1,5enynes to bicyclo[3.1.0]hexenes (eq 1),^{2c} we surmised that 5-en-2-yn-1-yl acetates such as **1** might be valuable precursors for the synthesis of acetoxy bicyclo[3.1.0]hexene (e.g., **3**) after a gold catalyzed sequence of allene (e.g., **2**) formation³ and cycloisomerization (eq 2). This approach would be advantageous since the presence of the acetoxy functionality at the ring junction might allow the synthesis of more elaborate structures such as 2-cyclohexen-1-ones.



Following our recent success in using the air stable crystalline $Ph_3PAuNTf_2$ ⁴ catalyst for the formation of C–C or C–O bonds, we chose this catalytic system to validate this approach (Table 1).

Treatment of propargylic acetate 4a with 1% of Ph₃PAuNTf₂ in CH₂Cl₂ at room temperature furnished the desired bicyclo[3.1.0]hexene 6a in 49% yield along with 41% of the postulated intermediate allene 5a (entry 1). Heating the reaction mixture at 40 °C did not improve the yield (entry 2). At 60 °C, the reaction was faster and afforded **6a** in good yield (entry 3). Changing the electronic properties of the phosphine had a significant effect on the course of the transformation (entries 4 and 5). While the electron deficient (p-CF₃Ph)₃PAuNTf₂ catalyst^{4c} gave **6a** in a modest 32% yield, the bulkier and more electron rich Ad₂n-BuPAuNTf₂^{4c} furnished 6a in 78% yield. The yield could even be improved to 94% by using 1% of Ad2n-BuPAuNTf2 and heating the reaction mixture at 60 °C for 0.5 h (entry 6). We next turned our attention on the use of biphenylphosphine based catalysts (entries 7 and 8).5 Remarkably, the use of 7 and 8 provided 6a in excellent yields (98% and 96%) in CH₂Cl₂ at room temperature.⁶ Furthermore, reacting allene $5a^7$ with 1% of 7 in CH₂Cl₂ at room temperature cleanly furnished 6a in 98% yield, thus proving the intermediacy of allene 5a in this transformation. It is worth notice that the use of Au(III), Pt(II), or Ag(I) catalysts did not furnish the desired bicyclo[3.1.0]hexene 6a.8 In light of these results, catalyst 7 was retained in the study of the scope of this transformation.

The reaction proved to be quite general and various substituted 5-en-2-yn-1-yl acetates $4\mathbf{b}-\mathbf{j}$ reacted using 1% of **7** as the catalyst to furnish the corresponding bicyclo[3.1.0]hexenes $6\mathbf{b}-\mathbf{j}$ in generally good yields (38%–99%) (Table 2). Substitution at the alkene was first examined. The reaction was exceptionally fast with methallyl derivative $4\mathbf{b}$ furnishing bicyclo[3.1.0]hexene $6\mathbf{b}$ in 97% yield (entry 1). Crotyl derivative $4\mathbf{c}$ reacted more slowly and gave $6\mathbf{c}$ as the sole product in 93% yield despite the fact that a cis/trans

Table 1. Optimization of the Catalytic System

I	OAc Ph	cataly	st 1% Ph			
4a Č			5a	6a ^{''}		
en	ntry catalys	st	conditions	time	yield (4a / 5a / 6a) ^a	
1	Ph_3AuNTf_2		CH ₂ Cl ₂ , rt	6h	0% / 41% / 49%	
2	Ph ₃ AuNTf ₂		CH ₂ Cl ₂ , reflux	6h	0% / 37% / 48%	
3	Ph ₃ AuNTf ₂		1,2-DCE, 60℃	2h	0% / 4% / 82%	
4	(pCF ₃ Ph) ₃ P/	AuNTf ₂	CH ₂ Cl ₂ , rt	6h	0% / 53% / 32%	
5	Ad ₂ n-BuPAu	INTf ₂	CH ₂ Cl ₂ , rt	6h	0% / 21% / 78%	
6	Ad ₂ n-BuPAuNTf ₂		1,2-DCE, 60℃	0.5h	0% / 0% / 94%	
7	7		CH ₂ Cl ₂ , rt	1h	0% / 0% / 98%	
8	8		CH ₂ Cl ₂ , rt	5h	0% / 0% / 96%	
	$\overset{\underline{R}}{\sim}\overset{\underline{P}}{}_{P}Au-NTf_{2}$	7 _{R'} 8	R= <i>t-</i> Bu, R'= H R=Cy, R'= <i>i-</i> Pr		^a isolated yields	

Table 2. Gold(I) CatalyzedFormation of Bicyclo[3.1.0]hexenes

en	try	substrate		product		time y	vield ^b
1	C Ph	Ac	4b	OAc Ph	6b	5 min	97%
2	C Ph	Ac	4c (E:Z=85:1		6c	5 h	93%
3	ر Ph	DAc	4d (dr=1:1)	Ph-	6d	30 min (exo:ende	98% o =1:2.8)
4	PhO	OAc	4e Ph		6e	2 h	99%
5	CI		4f ⁽		6f	2 h	с 38%
6	Pł	OAc	4g	OAc	6g	20 min	99%
7	/		4h		6h	2.5 h	72%
8	ە ب	AC CO ₂ E	Et 4i	OAc CO2Et CO2Et	6i	15 min	96%
9	C ₅ I	OAc H ₁₁	4j 99% ee	C ₅ H ₁₁	6j	15 min	92% 90% ee

^{*a*} Reaction conditions: 0.25 M substrate in CH₂Cl₂, 1% of **7**, rt. ^{*b*} Isolated yields. ^{*c*} A total of 44% of the corresponding allene were also isolated.

mixture of substrate was used (entry 2). Introduction of a methyl group at the allylic position was also tolerated, producing bicyclo-[3.1.0]hexene **6d** as a 1:2.8 mixture of diastereoisomers (entry 3).^{9,11}



^a Isolated vields

Scheme 1. Proposed Mechanism



The aromatic ring could be substituted, but with a decrease in efficiency when the 2,3-dichlorophenyl derivative 4f was used as the substrate (entries 4 and 5). The rearrangement proceeded as well when the propargylic position of the enyne was substituted with an alkyl chain (entries 6-9). Various substituted bicyclo[3.1.0]hexenes 6g-j possessing two adjacent quaternary centers at the ring junction were thus obtained in yields ranging from 72% to 99%. Finally, the cycloisomerization of enantioenriched substrate 4j was attempted. We were pleased to observe the rapid formation of bicyclo[3.1.0]hexene 6j which was isolated in 92% yield. Interestingly, the stereochemical information of the substrate was nearly completely transferred to the final product.¹⁰

The functionalized acetoxy bicyclo[3.1.0]hexene products lend themselves to a number of useful transformations. For example they can be efficiently converted into 2-cycloalken-1-ones by simple treatment with K₂CO₃ in methanol thus highlighting the general utility of this transformation (Table 3). The cleavage of the cyclopropane ring seems to be directed by the substitution pattern of the cylopropyl ring. Bicyclo[3.1.0]hexenes 6a, 6c, and 6d afforded the corresponding 2-cyclohexen-1-ones 7a, 7c, and 7d, while bicyclo[3.1.0]hexene **6b** bearing a methyl group at the ring junction gave 2-cyclopenten-1-one 7b.

To account for these observations, a mechanistic manifold for the formation of the bicyclo[3.1.0]hexenes is proposed in Scheme 1. Gold(I) activation of the triple bond in alkyne 8 promotes the formation of allene 9 through a [3,3]-sigmatropic rearrangement. A further gold(I) activation of the allene induces the nucleophilic attack of the pendant alkene resulting in the formation of the cationic vinyl-gold species 10.11 Subsequent formation of the cyclopropyl ring assisted by electron donation from gold(I) affords gold(I) carbene 11. A final 1,2-hydride shift regenerates the gold(I) catalyst and produces bicyclo[3.1.0]hexene 12.

The proposed mechanism suggests that intermediate 10 could be trapped by a nucleophile. In agreement with this hypothesis, cyclohexene 13 was formed in 81% yield as a 1:2 mixture of diastereoisomers when the rearrangement of 4b was performed in MeOH (eq 3). Interestingly, the nucleophilic addition of the alkene onto the gold(I) activated allene seems to be faster than the addition

of methanol onto the same activated allene. Subsequent basic treatment of 13 afforded 2-cyclohexen-1-one 14 in 99% yield. Furthermore, treatment of enantioenriched propargylic acetate 4j furnished 15 in 72% yield as a 1:1.5 mixture of epimers under the same experimental conditions (eq 4). A one-pot cycloisomerization deprotection sequence was also attempted and led to the formation of the corresponding 2-cyclohexen-1-one 16 with a complete retention of the chiral information. Note that the cyclohexenones obtained by that route (e.g., 16) are regioisomeric with those generated by opening the cyclopropane as in Table 3.

4b
$$\xrightarrow{1\% 7}_{MeOH, rt}$$
 $\xrightarrow{OAc}_{Ph} \xrightarrow{K_2CO_3}_{MeOH, rt}$ $\xrightarrow{O}_{Ph} \xrightarrow{(eq 3)}_{14 99\%}$ (eq 3)
 OAc $\xrightarrow{13 81\% (1/2)}_{C_5H_{11}}$ $\xrightarrow{O}_{MeOH, rt}$ 4j $\xrightarrow{1-1\% 7, MeOH}_{2-K_2CO_3}$ $\xrightarrow{O}_{Ph} \xrightarrow{(eq 3)}_{14 99\%}$ (eq 4)
15 72% (1/1.5) 16 79% 99% ee

In summary, we have developed an efficient gold(I) catalyzed cycloisomerization of 5-en-2-yn-1-yl acetates that provides an efficient access to acetoxy bicyclo[3.1.0]hexenes which can be further transformed into 2-cycloalken-1-ones. Cyclohexenones are key building blocks in numerous total syntheses. Further studies related to this new gold(I)-catalyzed process as well as its application to the synthesis of natural products are underway.

Acknowledgment. The authors wish to thank Prof. S. Z. Zard for helpful discussions.

Supporting Information Available: Experimental procedures and spectral data for new compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

References

- (1) For recent reviews, see: (a) Hashmi, A. S. K. Angew. Chem., Int. Ed. 2005, 44, 6990-6993. (b) Hoffman-Röder, A.; Krause, N. Org. Biomol. *Chem.* **2005**, *3*, 387–391. (c) Echavarren, A. M.; Nevado, C. *Chem. Soc. Rev.* **2004**, *33*, 431–436. (d) Arcadi, A. Di Giuseppe S. *Curr. Org. Chem.* **2004**, *8*, 795–812. (e) Hashmi, A. S. K. *Gold Bull.* **2003**, *36*, 3–9.
- (2) For selected examples of Au⁺ cycloisomerization of 1,n enynes, see (1,3 For selected examples of Au⁺ cycloisomerization of 1,n enynes, see (1,5-enyne): (a) Zhang, L.; Wang, S. J. Am. Chem. Soc. 2006, 128, 1442–1443, 1,4-enyne: (b) Shi, X.; Gorin, D. J.; Toste, F. D. J. Am. Chem. Soc. 2005, 127, 5802–5803, 1,5-enyne: (c) Luzung, M. R.; Markham, J. P.; Toste, F. D. J. Am. Chem. Soc. 2004, 126, 10858–10859. (d) Sherry, B. D.; Toste, F. D. J. Am. Chem. Soc. 2004, 126, 15978–10979. (e) Gagosz, F. Org. Lett. 2005, 7, 4129–4132. (f) Zhang, L.; Kozmin, S. A. J. Am. Chem. Soc. 2005, 127, 6962–16963. (g) Zhang, L.; Kozmin, S. A. J. Am. Chem. Soc. 2004, 126, 11807–1 (5, enyme). J. Am. Chem. Soc. 2004, 126, 11806-11807. 1,6-enyne: (h) Nieto-Oberhuber, C.; Munoz, M. P.; Lopez, S.; Jiménez-Nùnez, E.; Nevado, ; Herrero-Gomez, E.; Raducan, M.; Echavarren, A. M. Chem.-Eur. J. 2006, 12, 1677-1693.
- (3) For examples of Au⁺ catalyzed 3,3-rearrangements of propargylic esters, see: (a) ref 2a. (b) Zhang, L. J. Am. Chem. Soc. 2005, 127, 16804-16805. (c) Marion, N.; Diez-Gonzales, S.; de Frémont, P.; Noble, A. R.; Nolan, S. P. Angew. Chem., Int. Ed. 2006, 45, 1-5
- (4) (a) Buzas, A.; Istrate, F.; Gagosz, F. Org. Lett. 2006, 8, 1957–1959. (b) Buzas, A.; Gagosz, F. Org. Lett. 2006, 8, 515–518. (c) Mezailles, N.; Ricard, L.; Gagosz, F. Org. Lett. 2005, 7, 4133–4136.
 (5) Lopez, S.; Nieto-Oberhuber, C.; Echavarren, A. M. J. Am. Chem. Soc. 2005, 127, 6129. 6129.
- 2005, 127, 6178-6179.
- (6) Interestingly, aromatic groups do not interfere in the reaction (ref 3c).
- Allene 5a was obtained in 89% yield by reaction of 4a with 10% of
- AgNTf2 in CL_2 is obtained in CJ_2 includes the relation of 4a with 10% of 4a with 10% of 1 m CL_2 in CL_2 is room temperature for 30 min. In CL_2 in CL_2 in CL_2 is a non-temperature for 30 min. 4a, 17% 5a, 0% 6a. 10% AgNTf₂: 0% 4a, 89% 5a, 0% 6a.
- (9) The stereochemistry of the major endo isomer was assigned on the basis of NMR experiments.
- (10) The enantiomeric excess was determined by chiral HPLC analysis. However, the configuration of the major enantiomer was not determined.
- (11)A equilibrium between two conformers may explain the stereochemistry observed for 6d:

exo 6d

JA064223M