This article was downloaded by: [Case Western Reserve University] On: 05 November 2014, At: 01:00 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lcar20</u>

SYNTHESIS OF A CARBOHYDRATE-CENTERED C-GLYCOSIDE CLUSTER

Michael Dubber^a & Thisbe K. Lindhorst^b

^a Institut für Organische Chemie, Christian-Albrechts-Universität zu Kiel, Otto-Hahn-Platz 4, Kiel, D-24098, Germany

^b Institut für Organische Chemie, Christian-Albrechts-Universität zu Kiel, Otto-Hahn-Platz 4, Kiel, D-24098, Germany Published online: 16 Aug 2006.

To cite this article: Michael Dubber & Thisbe K. Lindhorst (2001) SYNTHESIS OF A CARBOHYDRATE-CENTERED C-GLYCOSIDE CLUSTER , Journal of Carbohydrate Chemistry, 20:7-8, 755-760

To link to this article: <u>http://dx.doi.org/10.1081/CAR-100108288</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

COMMUNICATION

SYNTHESIS OF A CARBOHYDRATE-CENTERED C-GLYCOSIDE CLUSTER¹

Michael Dubber and Thisbe K. Lindhorst*

Institut für Organische Chemie, Christian-Albrechts-Universität zu Kiel, Otto-Hahn-Platz 4, D-24098 Kiel, Germany

The carbohydrate moieties of glycoconjugates are involved in numerous recognition events in biological systems both in a physiological as well as a pathological context.^{2,3} In these processes, multivalency of molecular interactions⁴ plays an important role because certain sugar epitopes are frequently displayed in a polyvalent manner. With regard to the multiple-copy design of natural oligosaccharides, many means for clustering of carbohydrate ligands have been sought in order to provide synthetic mimetics which are more easily obtained than their natural counterparts, while possibly equally active.^{5, 4c} During our attempts to inhibit the adhesion of *Escherichia coli* bacteria⁶ to high mannose-type structures, we have employed different pathways for glycocluster synthesis comprising glycosidation,⁷ peptide coupling,⁸ thiourea bridging,⁹ and photoaddition reactions.¹⁰ In order to increase the stability of such compounds in a physiological environment, *C*-glycosides instead of *O*-glycosides can be employed in the design of carbohydrate-based antiadhesives.¹¹

Here we present the synthesis of the first carbohydrate-centered¹² *C*-glycoside cluster (5) using two spacer-modified carbohydrate derivatives (1 and 4) and peptide coupling in the ligation step. Two possible pathways for the synthesis of peptide bond-ligated glycoclusters can be envisaged. Either an oligocarboxylic acid is employed as the core molecule and coated with an amino-functionalized carbohydrate derivative, or an oligoamine core is peptide-coupled to a carboxylated saccharide epitope. In either of these two pathways, according to observations

^{*}Corresponding author.

ORDER		REPRINTS
-------	--	----------

DUBBER AND LINDHORST

made in our laboratory, a powerful coupling reagent has to be chosen to achieve multiple peptide coupling in good yields.

For the synthesis of glycodendrimers and glycoclusters using amide connectivities, HATU (N-[(dimethylamino)-1H-1,2,3-triazolo[4,5-b]pyridin-1-yl-methylene]-N-methylmethanaminium hexafluorophosphate N-oxide) was shown to be superior to other onium salts.¹³ In a peptide coupling reaction, HATU has to



Scheme 1. a. Bu₃SnH, AIBN, methyl acrylate, toluene; NaOMe, MeOH, 24%. b. LiOH, H₂O. c. HATU, DIPEA, DMF, 68% (two steps).





CARBOHYDRATE-CENTERED C-GLYCOSIDES

react first with the carboxylic acid component. This has to be a fast reaction in order to avoid decomposition of the expensive reagent.¹⁴ In order to prepare a cluster glycoside by peptide coupling, it is therefore not advisable to use an oligocarboxylic acid but rather a polyamine as the core molecule. The latter was reacted with a carboxy-functionalized C-glycoside (4) as the acid component (Scheme 1). C-Mannoside 4 (Scheme 1) was synthesized, starting from the glycosyl bromide 2 in a classical radical reaction using methyl acrylate.¹⁵ This reaction furnished methyl ester 3, which in turn was saponified¹⁶ to give the corresponding acid 4 in situ. This is a known compound,¹¹ which has, however, never been used in glycocluster synthesis. Here, it could be readily ligated to the glucose-centered pentaamino hydrochloride 1,¹⁰ which was selected as the core pentaamine in the peptide coupling reaction from a spectrum of different possibilities. 1.5 Equivalents of the acid 4 were used per amino group to obtain the C-glycosidically linked mannose cluster 5 in good yield. It is worthwhile to point out that due to the chiral core unit, NMR spectra of this type of molecules can be unequivocally solved (Scheme 2).

In conclusion, it was shown that a novel carbohydrate-centered *C*-mannoside glycocluster could be readily obtained by the peptide-coupling reaction of a branched oligoamine represented by **1** and a carboxylated *C*-mannoside such as **4**, involving a number of one-pot procedures en route to the product **5**. The method can be recommended for the multi-100 mg synthesis of the respective glycoclusters and will be further explored in our group.



757



ORDER		REPRINTS
-------	--	----------

EXPERIMENTAL

General methods. TLC was performed on silica gel plates (GF₂₅₄, Merck). Detection was effected by UV irradiation and subsequent charring with 10% sulphuric acid in ethanol followed by heat treatment. Flash chromatography was performed on silica gel 60 (230–400 mesh, particle size 0.040–0.063 mm, Merck). Optical rotations were measured on a Perkin-Elmer 241 polarimeter (sodium-D-line: 589 nm, length of cell 1 dm) in water. NMR spectra were recorded on Bruker AMX 400 (400.13 MHz for ¹H, 100.61 MHz for ¹³C) and DRX 500 (500.13 MHz for ¹H, 125.76 MHz for ¹³C). Assignment of the peaks was achieved with the aid of 2D-NMR techniques (¹H-¹H-COSY and HMQC). MALDI-TOF-mass spectra were recorded on a Bruker Biflex III with 19 kV acceleration voltage and DHB (2,4-di-hydroxy benzoic acid) as matrix ($c = 10 \ \mu g/\mu L$ in 40% acetonitrile/water). Ionisation was effected with a nitrogen laser at 337 nm. Elemental analyses were carried out in the Institute of Inorganic Chemistry, Christian-Albrechts-University Kiel.

Methyl 3-(α -D-mannopyranosyl)propanoate (3). A solution of 2,3,4,6tetra-O-acetyl- α -D-mannopyranosyl bromide (2) (17.72 g, 43.1 mmol) and methyl acrylate (19.5 mL, 215.2 mmol) in degassed toluene (50 mL) was stirred at 85 °C and a solution of Bu₃SnH (22.8 mL, 86.2 mmol) and AIBN (400 mg) in toluene (30 mL) was added dropwise over 1 h. Then stirring was continued for 0.5 h at 85 °C. The solution was concentrated and the residue dried in high vacuum. The residual syrup was dissolved in acetonitrile, washed with hexane, concentrated and purified by column chromatography (1:1 toluene-EtOAc; $R_f 0.42$). The crude acetylated product was dissolved in dry MeOH and stirred for 2 h with 1M NaOMe in MeOH. The solution was neutralized with Dowex-H⁺, filtered and purified by column chromatography (9:1 EtOAc-MeOH) to yield 1,5-anhydro-D-mannitol (470 mg, 7 %, R_f 0.06) as side product and the title compound 2 (2.62 g, 24 %, $R_f 0.15$) as a white solid: $[\alpha]_D + 18.7^{\circ} (c \ 1.4, H_2O); mp \ 77.7^{\circ}C; {}^{1}H NMR (400 \text{ MHz}, D_2O) \delta 3.80-3.95$ (m, 2H, H-1, H-2), 3.80–3.85 (m, 2H, H-3, H-6), 3.74 (dd≈t, 1H, J 6.1 Hz, H-4), 3.71 (s, 3H, OMe), 3.65 (dd≈t, 1H, J 9.4 Hz, H-6'), 3.51 (ddd, 1H, J 2.5, 6.1, 9.2 Hz, H-5), 2.43–2.58 (m, 2H, OCH₂CH₂CO₂Me), 2.11 (m, 1H, OCHHCH₂CO₂Me), 1.83 (m, 1H, OCHHCH₂CO₂Me) ppm; ¹³C-NMR (100.67 MHz, D₂O) δ 176.9 (CO₂Me), 78.0 (CH, C-1), 74.1 (C-5), 71.7 (C-2), 71.3 (C-3), 67.7 (C-4), 61.6 (C-6), 52.7 (OMe), 30.6 (OCH₂CH₂CO₂Me), 23.3 (OCH₂CH₂CO₂Me) ppm.

10-α-D-Mannopyranosyl-8-oxo-4-thia-7-azadecyl-2,3,4,6-tetra-O-[10α-D-mannopyranosyl-8-oxo-4-thia-7-azadecyl]-α-D-glucopyranoside (5). The octopus glycoside 3-(2-aminoethylthio)propyl-2,3,4,6-tetra-O-3(2-aminoethylthio)propyl-α-D-glucopyranoside (1)⁹ (54 mg, 0.057 mmol) and diisopropylethyl amine (DIPEA, 0.1 mL, 0.580 mmol) were dissolved in dry DMF (4 mL) and stirred for 1 d. In the meantime **3** (110 mg, 0.44 mmol) was dissolved in methanol (6 mL) and water (4 mL) and LiOH × H₂O (102 mg, 2.4 mmol) was added. The solution was stirred at 0 °C until ester cleavage was completed (TLC, MeOH). Additional water was added (20 mL) and the solution was neutralized at 0 °C with 2M hydrochloric acid and freeze dried to yield **4**, which was used as a crude product.



CARBOHYDRATE-CENTERED C-GLYCOSIDES

Then, a solution of 3 in DMF (4 mL) was added to the earlier prepared solution of 1. The coupling reagent HATU (N-[(dimethylamino)-1H-1,2,3-triazolo[4,5b]pyridin-1-yl-methylene]-N-methylmethanaminium hexafluorophosphate N-oxide, 167 mg, 0.44 mmol) and DIPEA (0.15 ml, 0.88 mmol) were added and the solution was stirred for 36 h at rt. For work-up, the solution was passed over a Sephadex LH-20 column with methanol as the eluent, to yield 5 (82 mg, 68 %) as a colorless syrup: $[\alpha]_{D}$ +22.0° (c 1.4, H₂O); ¹H NMR (D₂O) δ 4.95 (d, 1H, J_{1.2} 3.5 Hz, H-1_{glc}), 3.93–3.48 (m, 49H, 5 (glc)OCH₂, H-3_{glc}, H-5_{glc}, H-6_{glc}, H-6'_{glc}, 5 H-1_{man}, 5 H-2_{man}, 5 H-3_{man}, 5 H-4_{man}, 5 H-5_{man}, 5 H-6_{man}, 5 H-6'_{man}), 3.43–3.35 (m, 10H+MeOH, SCH₂CH₂N), 3.31-3.25 (m, 2H, H-2_{glc}, H-4_{glc}), 2.75-2.64 (m, 20H, 5 (glc) OCH₂CH₂CH₂SCH₂), 2.45–2.29 (m, 10H, 5 (man)CH₂CH₂), 2.13–2.01 (m, 5H, 5 (man)CHH), 2.00–1.80 (m, 15H, 5 (man)CHH, 5 (glc)OCH₂CH₂) ppm; ¹³C-NMR (125.77 MHz, D₂O): δ 99.1 (C-1_{glc}), 84.0 (C-3_{glc}), 83.0 (C-2_{glc}), 80.4 $(C-4_{glc})$, 78.7 (5×) (5 C-1_{man}), 77.4 (5×) (5 C-5_{man}), 74.0 (5×), 74.0 (5×) (5 C-2_{man}, 5 C-3_{man}), 72.9 (C-5_{glc}), 73.5, 72.0 (2×), 71.9, 71.3, 68.5 (C-6_{glc}, (glc)CH₂), 70.7 (5×) (5 C-4_{man}), 64.2 (5×) (5 C-6_{man}), 41.5 (5×) (5 SCH₂CH₂N), 34.5 (5×) (5 (man)CH₂CH₂), 33.3–33.2 (5×) (5 SCH₂CH₂N), 32.9, 32.5, 32.4, 32.0, 31.6 (5 $(glc)OCH_2CH_2)$, 30.7, 30.6 (2×), 30.5 (2×) (5 (glc)OCH_2CH_2CH_2S), 27.3 (5×) $(5(man)CH_2)$ ppm; MALDI-TOF-MS m/z 1879.4 $((M+Na)^+$ calcd 1855.8).

Anal.Calcd for C₇₆H₁₃₇N₅O₃₆S₅ (1857.24): C 49.15, H 7.44, N 3.77; Found C 48.61, H 7.24, N 3.36%.

ACKNOWLEDGMENTS

Support of this work by the Fonds der Chemischen Industrie (FCI) is gratefully acknowledged. We are indebted to Dr. V. Sinnwell and to Dr. C. Wolff for performing the NMR experiments.

REFERENCES

- Dedicated to Professor Dr. Joachim Thiem on the occasion of his 60th birthday. 1.
- 2. (a) Varki, A.; Cummings, R.; Esko, J.; Freeze, H; Hart, G; Marth, J., Eds.; Essentials of Glycobiology; Cold Spring Harbour Laboratory Press: Cold Spring Harbor, 1999; (b) Montreuil, J; Vliegenthart, J. F. G.; Schachter, H., Eds.; Glycoproteins and Disease; New Compr. Biochem. 30; Elsevier: Amsterdam, 1996.
- Kobata, A. A journey to the world of glycobiology. Glycoconjugate J. 2000, 17, 3. 443-464.
- 4. (a) Lee, Y. C.; Lee, R. T. Carbohydrate-protein interactions: basis of glycobiology. Acc. Chem. Res. 1995, 28, 321–327; (b) Kiessling, L. L.; Pohl, N. L. Strength in numbers: non-natural polyvalent carbohydrate derivatives. Chem. Biol. 1996, 3, 71-77; (c) Mammen, M.; Choi, D.-K.; Whitesides, G. M. Polyvalent interactions in biological systems: implications for design and use of multivalent ligands and inhibitors. Angew. Chem. 1998, 110, 2908–2953; Angew. Chem. Int. Ed. 1998, 37, 2754–2794; (d) Dimick, S. M.; Powell, S. C.; McMahon, S. A.; Moothoo, D. N.; Naismith, J. H.; Toone, E. J. On the meaning of affinity: cluster glycoside effects and concanavalin A.

759



Marcel Dekker, Inc.

270 Madison Avenue, New York, New York 10016

ORDER		REPRINTS
-------	--	----------

DUBBER AND LINDHORST

J. Am. Chem. Soc. **1999**, *121*, 10286–10296; (e) Lee, R. T.; Lee, Y. C. Affinity enhancement by multivalent lectin-carbohydrate interaction. Glycoconjugate J. **2000**, *17*, 543–551.

- (a) Roy, R. Glycodendrimers: A new class of biopolymers. Polymer News 1996, 21, 226–232; (b) Jayaraman, N.; Nepogodiev, S. A.; Stoddart, J. F. Synthetic carbohydrate-containing dendrimers. Chem. Eur. J. 1997, *3*, 1193–1199.
- (a) Sharon, N. Bacterial lectins, cell-cell recognition and infections disease. FEBS Lett. **1987**, *217*, 145–157; (b) Lindhorst, Th. K.; Kieburg, C.; Krallmann-Wenzel, U. Inhibition of type 1 fimbriae-mediated adhesion of *Escherichia coli* to erythrocytes by multiantennary α-mannosyl clusters: The effect of multivalency. Glycoconjugate J. **1998**, *15*, 605–613.
- Lindhorst, Th. K.; Dubber, M.; Krallmann-Wenzel, U.; Ehlers, S. Cluster-mannosides as inhibitors of type 1 fimbriae mediated adhesion of *Escherichia coli:* pentaerythritol derivatives as scaffolds. Eur. J. Org. Chem. 2000, 2027–2043.
- Kötter, S.; Krallmann-Wenzel, U.; Ehlers, S.; Lindhorst, Th. K. Multivalent ligands for the mannose-specific lectin on type 1 fimbriae of *Escherichia coli*: syntheses and testing of trivalent α-D-mannoside clusters. J. Chem. Soc., Perkin Trans. 1, **1998**, 2193–2200.
- Lindhorst, Th. K.; Kieburg C. Glycocoating of Oligovalent Amines: Synthesis of thiourea-bridged cluster glycosides from glycosyl isothiocyanates. Angew. Chem. 1996, 108, 2083–2086; Angew. Chem. Int. Ed. Engl. 1996, 35, 1953–1956.
- Dubber, M.; Lindhorst, Th. K. Synthesis of octopus glycosides: core molecules for the construction of glycoclusters and carbohydrate-centered dendrimers. Carbohydr. Res. 1998, 310, 35–41.
- 11. Bertozzi, C.; Bednarski, M. C-Glycosyl compounds bind to receptors on the surface of *Escherichia coli* and can target proteins of the organism. Carbohydr. Res. **1992**, 223, 243–253.
- (a) Kieburg, C.; Dubber, M.; Lindhorst, Th. K. A new type of carbohydrate clustering: synthesis of a pentavalent glycocluster based on a carbohydrate core. Synlett 1997, 1447–1449; (b) Fulton, D. A.; Stoddart, J. F. Toward the synthesis of large oligosaccharide-based dendrimers. Org. Lett. 2000, 2, 1113–1116; (c) Kitov, P. I.; Sadowska, J. M.; Mulvey, G.; Amstrong, G. D.; Ling, H.; Pannu, N. S.; Read, R. J.; Bundle, D. R. Shiga-like toxins are neutralized by tailored multivalent carbohydrate ligands. Nature 2000, 403, 669–672; (d) Furuike, T.; Aiba, S.; Nishimura, S.-I. A highly practical synthesis of cyclodextrin-based glycoclusters having enhanced affinity with lectins. Tetrahedron 2000, 56, 9909–9915.
- Patel, A.; Lindhorst, Th. K. A modular approach for the synthesis of oligosaccharide mimetics. J. Org. Chem. 2001, 66, 2674–2680.
- Albericio, F.; Bofill, J. M.; El-Faham, A.; Kates, S. A. Use of onium salt-based coupling reagents in peptide synthesis. J. Org. Chem. 1998, 63, 9678–9683.
- 15. Giese, B.; Dupuis, J.; Leising, M.; Nix, M.; Lindner, H. J. Synthesis of C-pento, hexo-, and -heptulopyranosyl compounds via radical C-C bond-formation reactions. Carbohydr. Res. **1987**, *171*, 329–341.
- 16. Kieburg, C.; Sadalapure, K; Lindhorst, Th. K. Glucose-based AB₂-building blocks for the construction of branched glycopeptidomimetics. Eur. J. Org. Chem. **2000**, 2035–2040.

Received June 7, 2001 Accepted August 21, 2001

Copyright © Marcel Dekker, Inc. All rights reserved



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the <u>U.S. Copyright Office</u> for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on <u>Fair Use in the Classroom</u>.

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our <u>Website</u> <u>User Agreement</u> for more details.

Order now!

Reprints of this article can also be ordered at http://www.dekker.com/servlet/product/DOI/101081CAR100108288