

Three New Sesquiterpenoid Glucosides of *Ficus pumila* Fruit

Junichi KITAJIMA,* Kaoru KIMIZUKA, and Yasuko TANAKA

Showa College of Pharmaceutical Sciences, Higashi-Tamagawagakuen 3, Machida, Tokyo 194–8543, Japan.

Received July 22, 1999; accepted September 14, 1999

As the glycosyl constituents of *Ficus pumila* L. fruits (Moraceae), three new sesquiterpenoid glucosides, pumilasides A, B and C were isolated together with benzyl β -D-glucopyranoside, (*E*)-2-methyl-2-butenyl β -D-glucopyranoside and rutin. Their structures were characterized as (1*S*,4*S*,5*R*,6*R*,7*S*,10*S*)-1,4,6-trihydroxyeudesmane 6-*O*- β -D-glucopyranoside, (1*S*,4*S*,5*S*,6*R*,7*R*,10*S*)-1,4-dihydroxyaalane 1-*O*- β -D-glucopyranoside and 10 α ,11-dihydroxycadin-4-ene 11-*O*- β -D-glucopyranoside by spectral and chemical methods.

Key words *Ficus pumila* fruit; sesquiterpenoid glycoside; eudesmane; aalane; cadinane; pumilaside

The fruit of *Ficus* (*F.*) *pumila* L. (Moraceae, *ōhitabi* in Japanese) has been used in Chinese folk medicine as antitumor, antiinflammatory and tonic medicament.¹⁾

In previous papers,²⁾ we reported on the sterol and triterpenoid components of this fruit. In this paper, we describe the isolation and characterization of three new sesquiterpenoid glucosides from the fruit, together with identification of the known glycosides.

The methanolic extract of the fresh fruit was suspended in water and then extracted with ether, ethyl acetate and *n*-butanol, successively. The *n*-butanol extract was treated as described in Experimental to isolate three new sesquiterpenoid glucosides, pumilaside A (**1**), pumilaside B (**2**) and pumilaside C (**3**), together with the known glycosides **4**–**6**, which were identified as benzyl β -D-glucopyranoside,³⁾ (*E*)-2-methyl-2-butenyl β -D-glucopyranoside⁴⁾ and rutin⁵⁾ by comparison of ¹H- and ¹³C-NMR spectra with those of authentic samples.

Pumilaside A (**1**, C₂₁H₃₈O₈, amorphous powder, $[\alpha]_D^{24}$ –28°) showed the $[M+K]^+$, $[M+Na]^+$, $[M+H]^+$ and $[M-C_6H_{12}O_6+H]^+$ ion peaks at *m/z* 457, 441, 419 and 239 on the positive FAB-MS. The ¹H-, ¹³C- and ¹³C–¹H correlation spectroscopy (COSY) NMR spectral data (Tables 1 and 2) showed the presence of one β -glucopyranosyl, two *tert*-methyls, two *sec*-methyls, four methylenes, five methines (two of them were oxygenated) and two quaternary carbons (one of them oxygenated). From the cross-peaks observed in the heteronuclear multiple bond correlation (HMBC) spectrum: H-1/C-9, C-10 and C-14; H-5/C-3, C-4, C-6, C-7, C-9, C-10 and C-14; H-6/C-7, C-10 and C-11; H-7/C-5 and C-12; H₂-9/C-5, C-7 and C-10; H₃-12/C-7, C-11 and C-14; H₃-13/C-7, C-11 and C-12; H₃-14/C-1, C-5, C-9 and C-10; H₃-15/C-3, C-4 and C-5, and ¹H–¹H COSY correlation data: H-1/H₂-2; H-6/H-5 and H-7; H-7/H-8 (δ 1.63) and H-11, a partial structure as described in Fig. 1 was obtained. Then, **1** was suggested to be a glucoside of eudesmane-type sesquiterpenoid having three hydroxyl groups at C-1, C-4 and C-6. The position of the glycosyl unit was ascertained to be C-6 from the HMBC correlation of glucosyl H-1/C-6, and from the observed nuclear Overhauser effect (NOE) interaction between the glucosyl H-1/H-6 in the nuclear Overhauser and exchange spectroscopy (NOESY) spectrum. The cross-peaks between H-6/H₃-14 and H₃-15; H₃-14/H₃-15 in the NOESY spectrum (Fig. 2) suggested that the orientation of H-6, H₃-14 and H₃-15 should be axial. Moreover, NOE interactions between H-5/H-1, H₃-12 and H₃-13 (Fig. 2), and the

small coupling constant (4.5 Hz) between Hax-6/H-7 suggested that the orientation of H-1, H-5 and the isopropyl group should be axial in the opposite direction to H-6, H₃-14 and H₃-15. So, **1** could be assumed to be a 6-*O*- β -D-glucopyranoside of 1 α ,4 β ,6 β -trihydroxyeudesmane or its enantiomer. Enzymatic hydrolysis of **1** gave an aglycone (**1a**, C₁₅H₂₈O₃, amorphous powder, $[\alpha]_D^{24}$ +6°) and D-glucose, and the absolute configuration at C-6 of **1** was indicated as *R* by the values of the glycosylation shift of the α - and the β -pro-*S*-side-carbons, and the chemical shift of the glucosyl anomeric carbon as shown in Table 3.⁶⁾ Thus, **1** was characterized as (1*S*,4*S*,5*R*,6*R*,7*S*,10*S*)-1,4,6-trihydroxyeudesmane 6-*O*- β -D-glucopyranoside as described in Fig. 2.

Pumilaside B (**2**, C₂₁H₃₆O₇, white powder [mp 195–197 °C (dec.)], $[\alpha]_D^{24}$ –19°) showed the $[M+K]^+$, $[M+Na]^+$, $[M-C_6H_{12}O_6+H]^+$ and $[M-C_6H_{12}O_6-H_2O+H]^+$ ion peaks at *m/z* 439, 423, 221 and 203 in the positive FAB-MS. The ¹H-, ¹³C- and ¹³C–¹H COSY NMR spectral data of **2** (Tables 1 and 2) showed the presence of one β -glucopyranosyl, four *tert*-methyls, four methylenes, four methines (one of them oxygenated), three quaternary carbons (one of them oxygenated). From the HMBC correlation data: H-5/C-1, C-3, C-4, C-6, C-9, C-10, C-11, C-14 and C-15; H-6/C-4, C-7, C-11, C-12 and C-13; H-7/C-5, C-6, C-9, C-11, C-12 and C-13; H₃-12/C-6, C-7, C-11 and C-13; H₃-13/C-6, C-7, C-11 and C-12; H₃-14/C-1, C-5, C-9 and C-10; H₃-15/C-3, C-4 and C-5, a partial structure as described in Fig. 1 was obtained and **2** was suggested to be a glucoside of aalane-type sesquiterpenoid having two hydroxyl groups at C-1 and C-4. The position of the glycosyl unit was ascertained to be C-1 in the same way as described for **1**. As the NOE interactions between the signals of H₃-14/Hax-2, H-6, H-7, Heq-9 and H₃-15; H₃-15/Heq-3 and H-6 were observed in the NOESY spectrum of **2** (Fig. 2), the orientation of H-6, H-7, H₃-14 and H₃-15 was concluded to be the same as **1**. Moreover, NOE interactions between H-5/H-1, Hax-3 and H₃-12 were observed in its NOESY spectrum (Fig. 2), and the orientation of H-1 and H-5 was opposite to H-6, H-7, H₃-14 and H₃-15. So, **2** was considered to be 1-*O*- β -D-glucopyranoside of 1 α ,4 β -dihydroxyaalane or its enantiomer. Enzymatic hydrolysis of **2** gave an aglycone (**2a**, C₁₅H₂₆O₂, mp 172–175 °C, $[\alpha]_D^{24}$ +10°) and D-glucose, and the values of the glycosylation shift of the α - and the β -pro-*S*-side-carbons, and the chemical shift of the glucosyl anomeric carbon (Table 3) suggested the absolute configuration at C-1 of **2** was *S*.⁶⁾ From these facts, **2** was determined as (1*S*,4*S*,5*S*,6*R*,7*R*,10*S*)-

* To whom correspondence should be addressed.

Table 1. ¹H-NMR Spectral Data for **1**, **2** and **3**

	1	2	3
H-1	3.69 (1H, t, <i>J</i> =7.0 Hz)	3.58 (1H, t, <i>J</i> =7.0 Hz)	1.71 (1H, ddd <i>J</i> =12.0, 10.0, 1.5 Hz)
H ₂ -2	1.90 (2H, m) ^{a)}	1.98 (1H, br ddd, <i>J</i> =13.0, 7.0, 3.0 Hz, H- α)	2.56 (1H, br ddd, <i>J</i> =15.0, 4.0, 1.5 Hz, H- α)
		2.42 (1H, dddd, <i>J</i> =13.0, 12.0, 7.0, 3.0 Hz, H- β)	1.40 (1H, br ddd, <i>J</i> =15.0, 12.0, 5.5 Hz, H- β)
H ₂ -3	1.93 (2H, m) ^{a)}	1.94 (1H, ddd, <i>J</i> =13.0, 12.0, 3.0 Hz, H- α)	2.11 (1H, br dd, <i>J</i> =15.0, 5.5 Hz, H- α)
		1.85 (1H, br dd, <i>J</i> =13.0, 3.0 Hz, H- β)	1.96 (1H, br dd, <i>J</i> =15.0, 4.0 Hz, H- β)
H-5	2.35 (1H, d, <i>J</i> =11.5 Hz)	1.38 (1H, d, <i>J</i> =6.0 Hz)	6.96 (1H, br s)
H-6	5.07 (1H, dd, <i>J</i> =11.5, 4.5 Hz)	0.85 (1H, dd, <i>J</i> =9.0, 6.0 Hz)	2.22 (1H, br dd, <i>J</i> =12.0, 10.0 Hz)
H-7	2.22 (1H, ddd, <i>J</i> =7.5, 4.5, 3.0 Hz)	0.56 (1H, t, <i>J</i> =9.0 Hz)	1.82 (1H, ddd, <i>J</i> =12.5, 12.0, 4.0 Hz)
H ₂ -8	1.63 (1H, m, H- α)	1.55 (1H, dd, <i>J</i> =15.0, 7.5 Hz, H- α)	1.79 (1H, dddd, <i>J</i> =12.5, 4.0, 3.5, 3.5 Hz, H- α)
	1.77 (1H, ddd, <i>J</i> =12.5, 6.0, 3.0 Hz, H- β)	1.75 (1H, m, H- β)	1.13 (1H, dddd, <i>J</i> =12.5, 12.5, 12.5, 3.5 Hz, H- β)
H ₂ -9	1.98 (1H, ddd, <i>J</i> =12.5, 3.0, 3.0 Hz, H- α)	2.44 (1H, m, H- α)	1.85 (1H, ddd, <i>J</i> =12.5, 12.5, 3.5 Hz, H- α)
	1.56 (1H, m, H- β)	0.91 (1H, ddd, <i>J</i> =13.0, 13.0, 7.5 Hz, H- β)	2.02 (1H, ddd, <i>J</i> =12.5, 3.5, 3.5 Hz, H- β)
H-11	2.48 (1H, m)		
H ₃ -12	1.44 (3H, d, <i>J</i> =6.5 Hz)	1.04 (3H, s)	1.494 (3H, s)
H ₃ -13	1.01 (3H, d, <i>J</i> =6.5 Hz)	1.07 (3H, s)	1.489 (3H, s)
H ₃ -14	1.24 (3H, s)	1.17 (3H, s)	1.32 (3H, s)
H ₃ -15	1.75 (3H, s)	1.45 (3H, s)	1.75 (3H, s)
Glc-1	5.18 (1H, d, <i>J</i> =7.5 Hz)	4.90 (1H, d, <i>J</i> =7.5 Hz)	5.13 (1H, d, <i>J</i> =7.5 Hz)
Glc-2	4.02 (1H, dd, <i>J</i> =9.0, 7.5 Hz)	4.04 (1H, dd, <i>J</i> =7.5, 7.0 Hz)	4.05 (1H, dd, <i>J</i> =9.0, 7.5 Hz)
Glc-3	4.27 (1H, t, <i>J</i> =9.0 Hz)	4.26 (1H, t, <i>J</i> =7.0 Hz)	4.25 (1H, t, <i>J</i> =9.0 Hz)
Glc-4	4.17 (1H, t, <i>J</i> =9.0 Hz)	4.27 (1H, t, <i>J</i> =7.0 Hz)	4.19 (1H, t, <i>J</i> =9.0 Hz)
Glc-5	4.05 (1H, m)	3.98 (1H, m)	3.92 (1H, m)
Glc-6	4.38 (1H, dd, <i>J</i> =12.5, 6.0 Hz)	4.42 (1H, dd, <i>J</i> =12.0, 5.0 Hz)	4.29 (1H, dd, <i>J</i> =12.0, 6.0 Hz)
	4.61 (1H, dd, <i>J</i> =12.5, 2.5 Hz)	4.55 (1H, dd, <i>J</i> =12.0, 2.5 Hz)	4.48 (1H, dd, <i>J</i> =12.0, 3.5 Hz)

Solvent: pyridine-*d*₅ (500 MHz). δ in ppm from TMS [coupling constants (*J*) in Hz are given in parentheses]. a) Assignment may be reversed.

Table 2. ¹³C-NMR Spectral Data for **1**, **1a**, **2**, **2a**, **3** and **3a**

	1	1a	2	2a	3	3a
Ag						
C-1	79.25	79.22	88.94	78.12	50.76	50.69
C-2	29.26	29.37	28.54	30.09	23.60	23.55
C-3	40.94	41.30	41.66	41.89	31.27	31.32
C-4	72.35	72.75	71.24	71.63	131.96	132.27
C-5	51.25	51.02	48.67	48.63	127.81	127.51
C-6	78.55	73.06	20.79	21.08	41.34	41.46
C-7	41.78	47.88	18.81	18.99	51.67	53.01
C-8	23.30	23.43	15.83	16.02	27.06	27.56
C-9	36.47	36.78	37.04	37.83	43.48	43.42
C-10	42.33	41.51	38.68	38.51	70.81	70.89
C-11	25.85	25.83	17.44	17.53	81.12	73.20
C-12	23.54	24.43	15.66	15.74	21.86	24.89
C-13	22.93	22.65	29.57	29.61	27.98	32.43
C-14	14.55	14.71	14.84	14.45	21.25	21.29
C-15	24.49	25.10	23.61	23.74	24.00	24.30
Glc						
C-1	100.30		106.69		98.43	
C-2	75.73		75.74		75.50	
C-3	78.87		78.69		79.21	
C-4	72.09		71.69		72.01	
C-5	78.57		78.21		77.77	
C-6	63.14		62.91		63.27	

Solvent: pyridine-*d*₅ (125 MHz). δ in ppm from TMS.

Table 3. Glycosylation Shift and Glucosyl C-1 Chemical Shift of **1** and **2**

Glucoside and carbon	$\Delta\delta$ (δ glucosyl – δ aglycone) or δ
1 α -Carbon (C-6)	$\Delta\delta$ +5.49 <i>R</i> -alcohols, about $\Delta\delta$ +5 to +8
2 α -Carbon (C-1)	$\Delta\delta$ +10.82 <i>S</i> -alcohols, about $\Delta\delta$ +10 to +11
1 β -pro- <i>S</i> -side-carbon (C-7)	$\Delta\delta$ –6.10 <i>R</i> -alcohols, about $\Delta\delta$ –4 to –5
2 β -pro- <i>S</i> -side-carbon (C-10)	$\Delta\delta$ +0.16 <i>S</i> -alcohols, about $\Delta\delta$ 0 to –2
1 Glucosyl C-1	δ 100.30 <i>R</i> -alcohols, about δ 102
2 Glucosyl C-1	δ 106.69 <i>S</i> -alcohols, about δ 106

1.85)/C-1, C-7, C-8, C-10 and C-14; H₃-12/C-7, C-11 and C-13; H₃-13/C-7, C-11 and C-12; H₃-14/C-1, C-9 and C-10; H₃-15/C-3, C-4 and C-5, and the ¹H–¹H COSY spectrum: H-6/H-7 showed the presence of a structure which is described in Fig. 1. Then, **3** was suggested to be a glucoside of cadinane-type sesquiterpenoid having a double bond at C-4(5) and two hydroxyl groups at C-10 and C-11. The comparison of ¹³C-NMR data with that of α -cadinol⁷⁾ supported this conclusion. The position of the glucosyl unit was ascertained to be C-11 in the same way as described for **1**. As the NOE interactions between the signals of H-6/H₃-12, H₃-14, and between the signals of H₃-14/Hax.-2, Hax.-8, Heq.-9, H₃-12 were observed in the NOESY spectrum of **3** (Fig. 2), the orientation of H₃-14, H-6 were suggested to be axial, the hydroxyisopropyl group attached to C-7 was equatorial, and the AB ring was indicated to be *trans*. Since D-glucose was obtained together with the aglycone **3a** by enzymatic hydrolysis, **3** was shown to have a 10 α ,11-dihydroxycadin-4-ene 11-*O*- β -D-glucopyranoside structure. But the absolute configuration of **3** could not be determined from available data.

This is the first report of the isolation of sesquiterpenoid

1,4-dihydroxymaaliane 1-*O*- β -D-glucopyranoside as described in Fig. 2.

Pumilaside C (**3**, C₂₁H₃₆O₇, amorphous powder, [α]_D²⁴ –17°) showed the [M+K]⁺, [M+Na]⁺ and [M–C₆H₁₂O₆+H]⁺ ion peaks at *m/z* 439, 423 and 221 in the positive FAB-MS. The ¹H-, ¹³C- and ¹³C–¹H COSY NMR spectral data of **3** (Tables 1 and 2) showed the presence of one β -glucopyranosyl, four *tert*-methyls, four methylenes, three methines, two oxygenated quaternary carbons and one trisubstituted double bond. The results of HMBC correlation: H-1/C-3, C-6, C-10 and C-14; H-5/C-1, C-3, C-6 and C-15; H-9 (δ




Fig. 1. Partial Structures of **1** to **3** Solved by HMBC (heavy lines) and ^1H - ^1H COSY (plot lines) Spectra




Fig. 2. Structures and NOE Interactions Observed in the NOESY Spectra of **1** to **3**

from plants of the *Ficus* genus.

Experimental

HPLC separation was carried out on a JASCO chromatography (980-system) with a JASCO 930 RI detector, JASCO OR-970 chiral detector and ODS-3251-D [Senshupak, column size, 8×250 mm], Symmetry Prep C₁₈ [Waters, column size, 7.8×300 mm], Megapak SIL C₁₈-10 [JASCO, column size, 7.5×250 mm], carbohydrate analysis [Waters, column size, 3.9×300 mm] column. The other instruments used and the experimental conditions for the spectral data and for chromatography were the same as in the preceding paper.^{2a)}

Extraction and Separation of 1 to 6 *F. pumila* L. was collected at Gushikawa City, Okinawa Prefecture, Japan, in March 1994. The fresh fruit (28 kg) was extracted with methanol (32 l) at room temperature. After evaporation of the solvent, the residue (987 g) was suspended with water and successively extracted with ether, ethyl acetate and *n*-butanol. Removal of the solvent from each phase gave an ether (43.1 g), ethyl acetate (6.5 g), *n*-butanol (35.1 g) and an aqueous (889 g) residue. The *n*-butanol residue was subjected to column chromatography on Amberlite XAD-II ($\text{H}_2\text{O} \rightarrow \text{MeOH}$) to afford water eluate (14.4 g) and methanol eluate (20.6 g). The methanol eluate fraction was chromatographed on Sephadex LH-20 (MeOH) which furnished four fractions. Fraction 2 (5.7 g) was purified by silica gel [CHCl_3 -MeOH (9:1 \rightarrow 8:2)] chromatography to afford six fractions. From the second fraction, **4** (2 mg) and **5** (3 mg) were isolated by Sephadex LH-20 (MeOH), Lobar RP-8 column (25% MeOH) and silica gel [CHCl_3 -MeOH (9:1)] chromatography, and HPLC using Symmetry Prep C₁₈ (20% MeOH) and ODS-3251-D (10% MeOH). From the third fraction (140 mg), **1** (13 mg) was isolated by Sephadex LH-20 (MeOH), Lobar RP-8 column (30% MeOH) silica gel [CHCl_3 -MeOH- H_2O (4:1:0.1)] chromatography and HPLC using carbohydrate analysis column (95% CH_3CN). From the fourth fraction (50 mg), **2** (22 mg) was isolated by Sephadex LH-20 (MeOH), silica gel [CHCl_3 -MeOH- H_2O (4:1:0.1)] chromatography and HPLC using Megapak SIL C₁₈ column (95% CH_3CN). From the fifth fraction (35 mg), **3** (11 mg) was isolated by Sephadex LH-20 (MeOH), silica gel [CHCl_3 -MeOH- H_2O (4:1:0.1)] chromatography and HPLC using Megapak SIL C₁₈ column (95% CH_3CN). Fraction 3 (3.2 g) was purified by Sephadex LH-20 (MeOH) and Lobar RP-8 column (40% MeOH) to get **6** (35 mg).

The following compounds were identified by comparison with authentic compounds.

Benzyl β -D-glucopyranoside (**4**), (*E*)-2-methyl-2-butenyl β -D-glucopyranoside (**5**) and rutin (**6**).

Pumilaside A (1) Amorphous powder, $[\alpha]_D^{24} -28^\circ$ ($c=1.0$, MeOH). Positive FAB-MS m/z : 457 $[\text{M}+\text{K}]^+$, 441.2417 $[\text{M}+\text{Na}]^+$ (base, Calcd for $\text{C}_{21}\text{H}_{38}\text{NaO}_8$: 441.2465), 419.2663 $[\text{M}+\text{H}]^+$ (Calcd for $\text{C}_{21}\text{H}_{39}\text{O}_8$: 419.2645), 399 $[\text{M}-\text{H}_2\text{O}+\text{H}]^+$ (base), 239 $[\text{M}-\text{C}_6\text{H}_{12}\text{O}_6+\text{H}]^+$.

Pumilaside B (2) White powder [mp $195-197^\circ\text{C}$ (dec.)], $[\alpha]_D^{24} -19^\circ$ ($c=1.8$, MeOH). Positive FAB-MS m/z : 439 $[\text{M}+\text{K}]^+$, 423.2314 $[\text{M}+\text{Na}]^+$ (Calcd for $\text{C}_{21}\text{H}_{36}\text{NaO}_7$: 423.2359), 221 $[\text{M}-\text{C}_6\text{H}_{12}\text{O}_6+\text{H}]^+$, 203 $[\text{M}-\text{C}_6\text{H}_{12}\text{O}_6-\text{H}_2\text{O}+\text{H}]^+$ (base).

Pumilaside C (3) Amorphous powder, $[\alpha]_D^{24} -17^\circ$ ($c=0.9$, MeOH). Positive FAB-MS m/z : 439 $[\text{M}+\text{K}]^+$, 423.2364 $[\text{M}+\text{Na}]^+$ (base, Calcd for $\text{C}_{21}\text{H}_{36}\text{NaO}_7$: 423.2359), 221 $[\text{M}-\text{C}_6\text{H}_{12}\text{O}_6+\text{H}]^+$.

Enzymatic Hydrolysis of 1, 2 and 3 Glycoside **1** (4 mg), **2** (6 mg) and **3** (3 mg) were each dissolved in water (5 ml) with hesperidinase (3 mg), and shaken in a water bath at 37°C for 2 weeks. The mixtures were evaporated *in vacuo* to dryness and the residues were chromatographed on silica gel [CHCl_3 -MeOH (9:1) \rightarrow CHCl_3 -MeOH- H_2O (7:3:0.5)] to give aglycone and the sugar fractions. The sugar fractions were passed through Sephadex LH-20 (MeOH) to give syrups. They were analyzed by HPLC [column, carbohydrate analysis (Waters: size, 3.9×300 mm), detector, JASCO RI-930 and OR-990 chiral detector: 85% CH_3CN , 2 ml/min; t_R 4.53 min] which revealed the presence of D-glucose. The aglycone fractions were purified by silica gel chromatography on silica gel [*n*-hexane-EtOAc (1:1)] to give aglycones (**1a**, 1.9 mg; **2a**, 2.0 mg; **3a**, 0.8 mg), respectively.

(1S,4S,5R,6R,7S,10S)-Trihydroxyeudesmane (1a) An amorphous powder, $[\alpha]_D^{24} +6^\circ$ ($c=0.2$, MeOH). Positive FAB-MS m/z : 256 $[\text{M}(\text{C}_{15}\text{H}_{28}\text{O}_3)+\text{H}]^+$. ^1H -NMR (pyridine- d_5) δ : 0.97 (3H, d, $J=6.5$ Hz, H₃-13), 1.24 (3H, s, H₃-14), 1.36 (3H, d, $J=6.5$ Hz, H₃-12), 1.62 (3H, s, H₃-15), 2.12 (1H, d, $J=11.5$ Hz, H-5), 2.29 (1H, ddd, $J=7.5$, 7.5, 4.5 Hz, H-7), 3.69 (1H, t, $J=7.0$ Hz, H-1), 4.67 (1H, dd, $J=11.5$, 4.5 Hz, H-6).

(1S,4S,5S,6R,7R,10S)-Dihydroxyeudesmane (2a) Colorless needles, mp $172-175^\circ\text{C}$, $[\alpha]_D^{24} +10^\circ$ ($c=0.2$, MeOH). Positive FAB-MS m/z : 239 $[\text{M}(\text{C}_{15}\text{H}_{26}\text{O}_2)+\text{H}]^+$. ^1H -NMR (pyridine- d_5) δ : 0.64 (1H, t, $J=9.0$ Hz, H-7), 0.91 (1H, dd, $J=9.0$, 6.0 Hz, H-6), 1.05 (3H, s, H₃-12), 1.10 (3H, s, H₃-13), 1.22 (3H, s, H₃-14), 1.42 (1H, d, $J=6.0$ Hz, H-5), 1.53 (3H, s, H₃-15), 3.59 (1H, t, $J=7.0$ Hz, H-1).

10 α ,11-Dihydroxycadin-4-ene (3a) Amorphous powder. ^1H -NMR (pyridine- d_5) δ : 1.34 (3H, s, H₃-14), 1.44 (6H, s, H₃-12 and H₃-13), 1.68 (3H, s, H₃-15).

Acknowledgments The authors thank Messrs. Y. Takase and H. Suzuki of the Central Analytical Department of this college for NMR and MS mea-

surements.

References

- 1) Kimura S., Konoshima M. (ed.), "Coloured Illustrations of Chinese Medical Plants," Yuhgonsha, Kyoto, 1986, pp. 412—413; Miyata S. (ed.), "Antitumoral Crude Drugs and their Prescriptions," Kagaku Shoin, Tokyo, 1981, pp. 25—27.
- 2) a) Kitajima J., Kimizuka K., Tanaka Y., *Chem. Pharm. Bull.*, **46**, 1408—1411 (1998); b) *Idem, ibid.*, **47**, 1138—1140 (1999).
- 3) Kitajima J., Ishikawa T., Tanaka Y., Ono M., Ito Y., Nohara T., *Chem. Pharm. Bull.*, **46**, 1587—1590 (1998).
- 4) Kitajima J., Ishikawa T., Tanaka Y., *Chem. Pharm. Bull.*, **46**, 1643—1646 (1998).
- 5) El-Kholy, Shaban M. A. M., *J. Chem. Soc., (C)*, **1966**, 1140—1142.
- 6) Kasai R., Suzuo M., Asakawa J., Tanaka O., *Tetrahedron Lett.*, **1977**, 175—178; Tori K., Seo S., Yoshimura Y., Arita Y., Tomita Y., *ibid.*, **1977**, 179—182; Kasai R., Okihara M., Asakawa J., Mizutani K., Tanaka O., *Tetrahedron*, **35**, 1427—1432 (1979); Mizutani K., Kasai R., Tanaka O., *Carbohydr. Res.*, **87**, 19—26 (1980); Kitajima J., Ishikawa T., Tanaka Y., *Chem. Pharm. Bull.*, **46**, 1643—1646 (1998); Ishikawa T., Kitajima J., Tanaka Y., Ono M., Ito Y., Nohara T., *ibid.*, **46**, 1738—1742 (1998); Kitajima J., Aoki Y., Ishikawa T., Tanaka Y., *ibid.*, **47**, 639—642 (1999).
- 7) Bottini A. T., Garfagnoli D. J., *J. Natural Products*, **50**, 732—734 (1987).