

Table 1. *Hydrogen-bonding geometry* (Å, °) for (I)

D—H...A	D—H	H...A	D—H...A
N16—H16...O11 ⁱ	0.92 (3)	2.25 (3)	166 (2)
C3—H3...O11 ⁱⁱ	0.90 (2)	2.69 (2)	176 (2)
C8—H8...O1 ⁱ	0.95 (2)	2.51 (2)	161 (2)
C15—H15A...F3 ⁱⁱⁱ	0.91 (3)	2.53 (3)	138 (2)

Symmetry codes: (i) $-x, -y, 1-z$; (ii) $1-x, -y, 2-z$; (iii) $x-1, y, z-1$.

Compound (II)

Crystal data

C₁₅H₁₄F₃NO₂ $M_r = 297.3$

Monoclinic

 $P2_1/n$ $a = 9.756$ (4) Å $b = 12.613$ (4) Å $c = 22.443$ (4) Å $\beta = 92.45$ (2)° $V = 2759.1$ (15) Å³ $Z = 8$ $D_x = 1.431$ Mg m⁻³ D_m not measured

Data collection

Rigaku AFC-6S diffractometer

 ω scans with profile analysis

Absorption correction:

 ψ scan (North *et al.*, 1968) $T_{\min} = 0.938$, $T_{\max} = 1.000$

5158 measured reflections

4856 independent reflections

Mo $K\alpha$ radiation $\lambda = 0.71069$ Å

Cell parameters from 25

reflections

 $\theta = 36.79$ – 42.39° $\mu = 0.1154$ mm⁻¹ $T = 296$ K

Prism

 $0.6 \times 0.5 \times 0.3$ mm

Yellow

2236 reflections with

 $I > 2\sigma(I)$ $R_{\text{int}} = 0.027$ $\theta_{\max} = 25.0^\circ$ $h = 0 \rightarrow 11$ $k = 0 \rightarrow 15$ $l = -26 \rightarrow 26$

3 standard reflections

every 150 reflections

intensity decay: 0.41%

Refinement

Refinement on F^2 $R(F) = 0.055$ $wR(F^2) = 0.127$ $S = 1.17$

4850 reflections

389 parameters

H atoms riding

 $w = 1/[\sigma^2(F_o^2) + (0.0548P)^2]$ $+ 1.0311P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.002$ $\Delta\rho_{\max} = 0.21$ e Å⁻³ $\Delta\rho_{\min} = -0.21$ e Å⁻³

Extinction correction: none

Scattering factors from

International Tables for Crystallography (Vol. C)

For both compounds, data collection: *MSC/AFC Diffractometer Control Software* (Molecular Structure Corporation, 1985); cell refinement: *MSC/AFC Diffractometer Control Software*; data reduction: *TEXSAN PROCESS* (Molecular Structure Corporation, 1988); program(s) used to solve structures: *MITHRIL* (Gilmore, 1984); program(s) used to refine structures: *SHELXL93* (Sheldrick, 1993); molecular graphics: *TEXSAN GRAPHICS*; software used to prepare material for publication: *TEXSAN FINISH*.

The authors wish to acknowledge the donors of the Petroleum Research Fund, administered by the American Chemical Society, for support of this research, and the National Science Foundation Research in Undergraduate Institutions Instrumentation Program (grant

No. 8818307) for creation of the New England Molecular Structure Center at Keene State College.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: BK1297). Services for accessing these data are described at the back of the journal.

References

- Gilmore, C. J. (1984). *J. Appl. Cryst.* **17**, 42–46.
 Jasinski, J. P. & Woudenberg, R. C. (1993). *Acta Cryst.* **C49**, 1965–1967.
 Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
 Molecular Structure Corporation (1985). *TEXSAN. TEXRAY Structure Analysis Package*. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
 Molecular Structure Corporation (1988). *MSC/AFC Diffractometer Control Software*. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
 Motherwell, W. D. S. & Clegg, W. (1978). *PLUTO. Program for Plotting Molecular and Crystal Structures*. University of Cambridge, England.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
 Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.

Acta Cryst. (1998). **C54**, 412–415

1,4-Bis(triphenylmethoxy)-2-butyne, Rhombohedral with $Z' = \frac{7}{6}$

MARÍA DEL ROSARIO BENITES, ROBERT P. HAMMER AND FRANK R. FRONCZEK

Department of Chemistry, Louisiana State University, Baton Rouge, LA 70803-1804, USA. E-mail: fronz@chxray1.chem.lsu.edu

(Received 7 October 1997; accepted 13 November 1997)

Abstract

The title compound, C₄₂H₃₄O₂, crystallizes in the space group $R\bar{3}$ with one molecule disordered at a crystallographic $\bar{3}$ site and another in a general position. The conformation of the linkage of the trityl groups to the central butyne group is such that the O—C...C—O grouping forms a torsion angle of magnitude 87.2 (2)°. The C≡C length in the ordered molecule is 1.176 (3) Å.

Comment

The ditritylated title compound, (1), was isolated as a minor side-product during the synthesis of 4-tri-





Fig. 1. ORTEP (Johnson, 1976) drawing of the ordered molecule of (1), with 30% probability ellipsoids and H atoms of arbitrary radii.

phenylmethoxy-2-butyne-1-ol from 2-butyne-1,4-diol. The structure determination was carried out in order to confirm its identity.



Compound (1) crystallizes in the rhombohedral space group $R\bar{3}$, with the uncommon value of $Z = 21$ ($Z' = \frac{7}{6}$). There are two independent molecules in the asymmetric unit, one lying in a general position (Fig. 1) and another disordered at a crystallographic $\bar{3}$ site rendering geometric parameters for this (disordered) molecule unreliable. The triply-bonded C atom (C2A) and the central C atom of the trityl group (C3A) doubtless also exhibit unresolved disorder, rendering geometric parameters for this molecule unreliable. The conformation of the central O—C—C $\equiv$ C—C—O group of the ordered molecule is non-planar, with the O1, C1, C2 and O2, C4, C3 planes almost perpendicular to one another, forming a dihedral angle of $87.2(2)^\circ$. Similar conformations have been observed in 2-butyne-1,4-diol and its inclusion compound with α -cyclodextrin (Steiner, 1996), in which the corresponding dihedral angles are $94.7(3)$ and 83° , respectively. The angles involving the C $\equiv$ C bond (C1—C2—C3 and C2—C3—C4) are close to linearity. The *O*-trityl groups are *anti* with respect to the C1—C2 and C3—C4 bonds [C5—O1—C1—C2 torsion angle $-169.62(13)$

and C24—O2—C4—C3 $-170.5(2)^\circ$]. The phenyl rings in the trityl groups do not present a symmetrical propeller conformation as does triphenylmethanol (Ferguson *et al.*, 1992), but a less symmetric arrangement, with the O1—C5—C—C and O2—C24—C—C torsion angles having magnitudes in the range $5.2(3)$ – $78.8(2)^\circ$. The C(trityl)—O bond lengths (C5—O1 and C24—O2) are within the range observed for the equivalent C—O bonds in triphenylmethanol (1.44 Å), tris(4-methoxyphenyl)methanol [1.441(3)–1.459(3) Å; Ferguson *et al.*, 1996] and substituted diphenylmethanols [1.421(4)–1.462(4) Å; Ferguson *et al.*, 1995].

A search of the Cambridge Structural Database (Allen & Kennard, 1993; Version of April 1997; *ca* 168 000 entries) for $Z = 21$ yielded but three hits, all of which are rhombohedral. CCl₄ at 229 K (Rudman & Post, 1968) is reported to be rhombohedral with $Z = 21$, but the space group is unknown. Tris(tricyclohexylphosphine)platinum heptane solvate (Immirzi *et al.*, 1977) is reported to be ordered in space group $R\bar{3}$. The lone precedent for $R\bar{3}$ with $Z = 21$ is dodecamethylcyclotetrasilazane (Edwards *et al.*, 1976), which also has one ordered molecule in a general position and one disordered on a $\bar{3}$ site.

Experimental

The title compound was prepared by treatment of triphenylmethyl chloride with a large excess of 2-butyne-1,4-diol in the presence of catalytic amounts of 4-dimethylaminopyridine in CH₂Cl₂. Purification by flash chromatography (hexane–ethyl acetate 1:1) afforded compound (1) as a minor side-product in 2% yield. Crystals were grown by layering a CH₂Cl₂ solution of (1) with CH₃OH.

*Crystal data*C₄₂H₃₄O₂*M_r* = 570.74

Trigonal

*R*³*a* = 36.120 (2) Å*c* = 14.787 (1) Å*V* = 16 707 (2) Å³*Z* = 21*D_x* = 1.191 Mg m⁻³*D_m* not measuredCu *K*α radiation*λ* = 1.54184 Å

Cell parameters from 25 reflections

θ = 22–38°*μ* = 0.52 mm⁻¹*T* = 295 K

Fragment

0.40 × 0.25 × 0.23 mm

Colorless

Data collection

Enraf–Nonius CAD-4 diffractometer

θ/2*θ* scans

Absorption correction:

ψ scans (North *et al.*, 1968)*T_{min}* = 0.86, *T_{max}* = 0.89

8031 measured reflections

7651 independent reflections

5705 reflections with

I > *σ*(*I*)*R_{int}* = 0.017*θ_{max}* = 75°*h* = −45 → 0*k* = 0 → 39*l* = −18 → 18

3 standard reflections

frequency: 120 min

intensity decay: 0.8%

*Refinement*Refinement on *F*²*R*(*F*) = 0.057*wR*(*F*²) = 0.059*S* = 1.206

5705 reflections

476 parameters

H atoms calculated

w = 4*F*_o²/[*σ*²(*F*_o²)+ 0.0004*F*_o⁴](Δ/*σ*)_{max} = 0.008Δ*ρ*_{max} = 0.34 e Å⁻³Δ*ρ*_{min} = −0.06 e Å⁻³

Extinction correction:

isotropic (Zachariasen, 1963)

Extinction coefficient:

5.4 (2) × 10⁻⁷Scattering factors from *International Tables for X-ray**Crystallography* (Vol. IV)Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters (Å²)
$$U_{eq} = (1/3) \sum_i \sum_j U^{ij} a_i^* a_j^* \mathbf{a}_i \cdot \mathbf{a}_j$$

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U_{eq}</i>
O1	0.24736 (3)	0.04764 (4)	0.49284 (8)	0.0455 (6)
O2	0.24019 (4)	0.08196 (4)	0.14049 (8)	0.0445 (6)
C1	0.21143 (5)	0.04739 (6)	0.4529 (1)	0.0564 (9)
C2	0.21407 (5)	0.04435 (5)	0.3544 (1)	0.0461 (8)
C3	0.21516 (5)	0.04302 (5)	0.2751 (1)	0.0452 (8)
C4	0.21575 (6)	0.03997 (5)	0.1764 (1)	0.0498 (9)
C5	0.24599 (5)	0.04299 (5)	0.5904 (1)	0.0414 (8)
C6	0.21343 (5)	−0.00324 (5)	0.6162 (1)	0.0427 (8)
C7	0.19888 (6)	−0.01427 (6)	0.7044 (1)	0.054 (1)
C8	0.17303 (7)	−0.05648 (7)	0.7286 (1)	0.064 (1)
C9	0.16119 (7)	−0.08843 (7)	0.6663 (2)	0.067 (1)
C10	0.17497 (7)	−0.07812 (6)	0.5793 (2)	0.064 (1)
C11	0.20085 (6)	−0.03597 (6)	0.5543 (1)	0.053 (1)
C12	0.23771 (5)	0.07703 (5)	0.6316 (1)	0.0490 (8)
C13	0.27090 (6)	0.11882 (6)	0.6394 (2)	0.067 (1)
C14	0.26350 (8)	0.15115 (7)	0.6664 (2)	0.088 (1)
C15	0.22331 (8)	0.14286 (7)	0.6851 (2)	0.103 (1)
C16	0.18956 (7)	0.10185 (7)	0.6768 (2)	0.099 (1)
C17	0.19702 (6)	0.06897 (6)	0.6511 (1)	0.071 (1)
C18	0.29026 (5)	0.04992 (5)	0.6156 (1)	0.0430 (8)
C19	0.30282 (6)	0.05430 (6)	0.7057 (1)	0.059 (1)
C20	0.34193 (6)	0.05946 (7)	0.7297 (2)	0.069 (1)
C21	0.36969 (6)	0.06074 (6)	0.6652 (2)	0.067 (1)
C22	0.35778 (6)	0.05617 (6)	0.5759 (2)	0.065 (1)
C23	0.31829 (5)	0.05049 (6)	0.5510 (1)	0.0525 (9)

C24	0.23833 (5)	0.08529 (5)	0.0430 (1)	0.0426 (8)
C25	0.27306 (5)	0.13131 (5)	0.0208 (1)	0.0460 (9)
C26	0.29582 (5)	0.16071 (5)	0.0869 (1)	0.0488 (9)
C27	0.32768 (6)	0.20173 (6)	0.0645 (2)	0.057 (1)
C28	0.33666 (6)	0.21369 (6)	−0.0233 (2)	0.064 (1)
C29	0.31397 (8)	0.18517 (7)	−0.0901 (2)	0.077 (1)
C30	0.28240 (8)	0.14416 (7)	−0.0686 (1)	0.065 (1)
C31	0.24876 (5)	0.05304 (5)	0.0008 (1)	0.0456 (8)
C32	0.29102 (6)	0.06225 (6)	−0.0012 (1)	0.061 (1)
C33	0.30133 (6)	0.03191 (7)	−0.0276 (2)	0.084 (1)
C34	0.26972 (7)	−0.00803 (6)	−0.0528 (2)	0.089 (1)
C35	0.22802 (7)	−0.01777 (6)	−0.0502 (2)	0.078 (1)
C36	0.21716 (6)	0.01243 (6)	−0.0238 (1)	0.059 (1)
C37	0.19514 (5)	0.08063 (5)	0.0161 (1)	0.0486 (9)
C38	0.17984 (6)	0.07211 (7)	−0.0716 (1)	0.070 (1)
C39	0.14273 (7)	0.07270 (8)	−0.0941 (2)	0.096 (1)
C40	0.12077 (7)	0.08174 (7)	−0.0306 (2)	0.094 (1)
C41	0.13527 (7)	0.08924 (7)	0.0558 (2)	0.085 (1)
C42	0.17208 (6)	0.08910 (6)	0.0792 (2)	0.063 (1)
O1A†	0.01588 (9)	0.0043 (2)	0.1890 (2)	0.060 (2)
C1A†	−0.0011 (2)	0.0192 (2)	0.1308 (4)	0.070 (4)
C2A	0	0	0.0379 (2)	0.0655 (8)
C3A	0	0	0.2871 (2)	0.0620 (8)
C4A	0.04568 (6)	0.03162 (7)	0.3199 (1)	0.068 (1)
C5A	0.05367 (8)	0.05933 (9)	0.3898 (2)	0.086 (2)
C6A	0.09442 (9)	0.0859 (1)	0.4232 (3)	0.105 (2)
C7A	0.12747 (9)	0.0857 (1)	0.3879 (3)	0.117 (2)
C8A	0.12252 (7)	0.0587 (1)	0.3205 (2)	0.134 (2)
C9A	0.08113 (8)	0.03132 (9)	0.2854 (2)	0.131 (2)

† Site occupancy = 0.33.

Table 2. Selected geometric parameters (Å, °)

O1—C1	1.422 (3)	C1—C2	1.467 (3)
O1—C5	1.450 (2)	C2—C3	1.176 (3)
O2—C4	1.422 (2)	C3—C4	1.463 (3)
O2—C24	1.451 (2)		
C1—O1—C5	115.7 (1)	C1—C2—C3	176.7 (3)
C4—O2—C24	115.5 (1)	C2—C3—C4	178.3 (2)
O1—C1—C2	108.6 (2)	O2—C4—C3	108.5 (1)
C5—O1—C1—C2	−169.6 (1)	O1—C5—C12—C13	−78.8 (2)
C1—O1—C5—C18	−173.2 (1)	O1—C5—C18—C23	−12.4 (2)
C24—O2—C4—C3	−170.5 (2)	O2—C24—C25—C26	−5.2 (3)
C4—O2—C24—C25	−170.0 (2)	O2—C24—C31—C32	−76.3 (2)
O1—C5—C6—C11	20.9 (3)	O2—C24—C37—C42	22.3 (2)

The H atoms on the disordered C1A atom were not located. Atoms C1A and O1A were assigned a population of $\frac{1}{3}$. For the calculation of H-atom positions, a C—H distance of 0.95 Å was assumed. Isotropic displacement parameters for H atoms were assigned as *B_{iso}* = 1.3*B_{eq}* of the bonded C atom.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1994). Cell refinement: *CAD-4 Software*. Data reduction: *PROCESS* in *MolEN* (Fair, 1990). Program(s) used to solve structure: *MULTAN80* (Main *et al.*, 1980). Program(s) used to refine structure: *LSFM* in *MolEN*. Software used to prepare material for publication: *BTABLE PTABLE CIF IN* in *MolEN*.

The purchase of the diffractometer was made possible by a National Science Foundation chemical instrumentation grant, which we gratefully acknowledge. Improvements to the LSU X-ray Crystallography Facility were supported by Grant No. LEQSF(1996–97)–ENH-TR-10, administered by the Louisiana Board of Regents.


Supplementary data for this paper are available from the IUCr electronic archives (Reference: FG1398). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H. & Kennard, O. (1993). *Chem. Des. Autom. News*, **8**, 31–37.
- Edwards, B. P. E., Harrison, W., Nowell, I. W., Post, M. L., Shearer, H. M. M. & Trotter, J. (1976). *Acta Cryst.* **B32**, 648–649.
- Enraf–Nonius (1994). *CAD-4 Software*. Version 5.1/1.2. Enraf–Nonius, Delft, The Netherlands.
- Fair, C. K. (1990). *MolEN. An Interactive Intelligent System for Crystal Structure Analysis*. Enraf–Nonius, Delft, The Netherlands.
- Ferguson, G., Carroll, C. D., Glidewell, C., Zakaria, C. M. & Lough, A. J. (1995). *Acta Cryst.* **B51**, 367–377.
- Ferguson, G., Gallagher, J. F., Glidewell, C., Low, J. N. & Scrimgeour, S. N. (1992). *Acta Cryst.* **C48**, 1272–1275.
- Ferguson, G., Glidewell, C. & Patterson, I. L. J. (1996). *Acta Cryst.* **C52**, 420–423.
- Immirzi, A., Musco, A. & Mann, B. E. (1977). *Inorg. Chim. Acta*, **21**, L37–L38.
- Johnson, C. K. (1976). *ORTEP*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Main, P., Fiske, S. J., Hull, S. E., Lessinger, L., Germain, G., Declercq, J.-P. & Woolfson, M. M. (1980). *MULTAN80. A System of Computer Programs for the Automatic Solution of Crystal Structures from X-ray Diffraction Data*. Universities of York, England, and Louvain, Belgium.
- North, A. C. T., Phillips, D. C., & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Rudman, R. & Post, B. (1968). *Mol. Cryst.* **5**, 95–110.
- Steiner, T. (1996). *Acta Cryst.* **C52**, 2885–2887.
- Zachariasen, W. H. (1963). *Acta Cryst.* **16**, 1139–1144.

Comment

The structure of the title salt has been investigated by X-ray diffraction as a part of a systematic study (Castellari & Sabatino, 1994, 1996; Castellari & Ottani, 1995, 1996, 1997) of diclofenac [2-(2,6-dichlorophenylamino)-phenylacetic acid] salts, an important group of non-steroidal anti-inflammatory drugs. Alterations in solubility and chemical stability have important pharmaceutical consequences since they can modify the bioavailability of a drug. Such alterations depend on the presence of polymorphic forms, crystalline stoichiometry and water molecules in the solid. Thus, the successful design of a dosage form of the drug depends strongly on knowledge of its solid-state chemistry. The asymmetric unit of the title compound, $(\text{H}_2\text{PZ})\text{D}_2 \cdot 2\text{H}_2\text{O}$ [where H_2PZ is the piperazinium dication and D is the 2-(2,6-dichlorophenylamino)phenylacetate anion] is shown in Fig. 1.



Acta Cryst. (1998). **C54**, 415–417

Stereochemistry of Anti-Inflammatory Drugs. VI. Piperazinium 2-(2,6-Dichlorophenylamino)phenylacetate Dihydrate, $(\text{H}_2\text{PZ})\text{D}_2 \cdot 2\text{H}_2\text{O}$

CARLO CASTELLARI^a AND STEFANO OTTANI^b

^a*Dipartimento di Chimica 'G. Ciamician', Università di Bologna, Via Selmi 2, 40126 Bologna, Italy, and* ^b*Centro Studi Fisica Macromolecole, c/o Dipartimento di Chimica 'G. Ciamician', Università di Bologna, Via Selmi 2, 40126 Bologna, Italy. E-mail: stefano@frodo.ciam.unibo.it*

(Received 31 July 1997; accepted 30 October 1997)

Abstract

The asymmetric unit of the title compound, $\text{C}_4\text{H}_{12}\text{N}_2^{2+} \cdot 2\text{C}_{14}\text{H}_{10}\text{Cl}_2\text{NO}_2^- \cdot 2\text{H}_2\text{O}$, consists of two piperazinium dications located around two non-equivalent centers of symmetry, two 2-(2,6-dichlorophenylamino)phenylacetate anions in general positions and two water molecules. The solid-state structure is strongly hydrogen bonded.

The asymmetric unit contains two bivalent piperazinium cations located around the symmetry centers $(0,0,\frac{1}{2})$ and $(\frac{1}{2},\frac{1}{2},\frac{1}{2})$, two 2-(2,6-dichlorophenylamino)phenylacetate anions and two water molecules. The structure is characterized by the presence of several hydrogen bonds (Table 2). The cations, anions and water molecules are connected by intermolecular hydrogen bonds to form a two-dimensional network in the *ab* plane. No short contact is detected along the *c* direction of the cell. The carboxylate groups, cations and water molecules are arranged along the $[010]$ direction to form a hydrophilic tunnel. Water molecules interact with both H_2PZ and D ions, but not with each other.

The H_2PZ cations form salt bridges with the carboxylate groups. The $\text{N} \cdots \text{O}$ distances are short, as expected for hydrogen bonds in which both donor and acceptor atoms carry a formal charge. However, while both the ammonium H atoms (H11 and H12) at each end of the cation at $(0,0,\frac{1}{2})$ interact with carboxylate O atoms, only one H atom (H12A) from each end of the cation at $(\frac{1}{2},\frac{1}{2},\frac{1}{2})$ forms a salt bridge. As detailed in Table 2, the other ammonium H atom (H11A) is strongly bonded to the O3 water molecule. As a consequence, O3 is more closely packed than the O3A water molecule [mean contact distances 2.853(2) and 2.994(2) Å for O3 and O3A, respectively]. This feature is probably related to the simultaneous donor and acceptor character of O3 in the hydrogen-bonding