

433.4, 422.2, 358.7, 337.5, 283.5, 239.1, 223.7, 198.3, 177.4, 150.4; UV-vis [CH₃COCH₃, λ (nm)]: 444, 402, 340, 315; ¹H NMR [CD₃CODD₃, δ (p.p.m.)]: 7.20–7.60 (*m*, 10H, SPH), 3.46 (*t*, 8^aH, C^aH₂CH₂CH₂CH₃), 1.83 (*m*, 8^bH, CH₂C^bH₂CH₂CH₃), 1.47 (*m*, 8^cH, CH₂CH₂C^cH₂CH₃), 1.01 (*t*, 12^dH, CH₂CH₂CH₂C^dH₃).

Crystal data

(C₁₆H₃₆N)[Mo₂I-
(C₆H₅S)₂(CO)₆]

M_r = 947.62

Monoclinic

*P*2₁/*n*

a = 9.8951 (4) Å

b = 16.2430 (7) Å

c = 25.0787 (10) Å

β = 90.444 (1)°

V = 4030.7 (3) Å³

Z = 4

D_x = 1.562 Mg m⁻³

D_m not measured

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 6187 reflections

θ = 1.49–25.02°

μ = 1.531 mm⁻¹

T = 293 (2) K

Needle

0.40 × 0.15 × 0.10 mm

Black

Data collection

Siemens SMART CCD diffractometer

ω scans

Absorption correction:

empirical (SADABS;

Sheldrick, 1996)

T_{min} = 0.698, *T_{max}* = 0.894

13 287 measured reflections

6816 independent reflections

4719 reflections with

I > 2σ(*I*)

R_{int} = 0.038

$\theta_{\max}$ = 25.02°

h = -11 → 11

k = 0 → 19

l = 0 → 29

Intensity decay: none

Refinement

Refinement on *F*²

R [*F*² > 2σ(*F*²)] = 0.051

wR (*F*²) = 0.089

S = 1.054

6816 reflections

415 parameters

H atoms not refined

w = 1/[σ²(*F_o*²) + (0.0168*P*)² + 2.8450*P*]

where *P* = (*F_o*² + 2*F_c*²)/3

(Δ/σ)_{max} = 0.001

Δρ_{max} = 0.384 e Å⁻³

Δρ_{min} = -0.428 e Å⁻³

Extinction correction: none

Scattering factors from

International Tables for Crystallography (Vol. C)

Table 1. Selected geometric parameters (Å, °)

I—Mo1	2.9460 (6)	Mo1—Mo2	2.8824 (7)
I—Mo2	2.9599 (6)	Mo2—C6	1.947 (7)
Mo1—C3	1.919 (7)	Mo2—C5	1.992 (6)
Mo1—C1	2.004 (7)	Mo2—C4	2.000 (7)
Mo1—C2	2.017 (7)	Mo2—S1	2.4783 (15)
Mo1—S1	2.4675 (15)	Mo2—S2	2.4836 (15)
Mo1—S2	2.4714 (15)		
Mo1—I—Mo2	58.43 (2)	Mo1—S1—Mo2	71.29 (4)
S1—Mo1—S2	108.84 (5)	Mo1—S2—Mo2	71.14 (4)
S1—Mo2—S2	108.10 (5)		

Data were collected over a hemisphere of reciprocal space by a combination of three sets of exposures. Each set had a different φ angle for the crystal and each exposure of 10 s covered 0.3° in ω . The crystal-to-detector distance was 4.95 cm. Coverage of the unique set was over 99% complete to at least 25° in θ . Crystal decay was monitored by measurement of duplicate

reflections. H atoms were all located theoretically and not refined.

Data collection: SMART (Siemens, 1996). Cell refinement: SMART and SAINT (Siemens, 1994b). Data reduction: XPREP in SHELXTL (Siemens, 1994a). Program(s) used to solve structure: SHELXTL. Program(s) used to refine structure: SHELXTL. Molecular graphics: SHELXTL. Software used to prepare material for publication: SHELXTL.

We are grateful to NNSF and SKLSC for financial support of this work.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: TA1215). Services for accessing these data are described at the back of the journal.

References

- Pan, G., Zhuang, B. & Chen, J. (1998). *Acta Cryst.* **C54**, 1815–1816.
Sheldrick, G. M. (1996). *SADABS. Program for Absorption Corrections*. University of Göttingen, Germany.
Siemens (1994a). *SHELXTL*. Release 5.03. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
Siemens (1994b). *SAINT. Software Reference Manual*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
Siemens (1996). *SMART. Software Reference Manual*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
Smith, D. A., Zhuang, B., Newton, W. E., McDonald, J. W. & Schultz, F. A. (1987). *Inorg. Chem.* **26**, 2524–2531.
Zhuang, B., Huang, L. & He, L. (1995). *Chin. J. Struct. Chem.* **14**, 359–363.
Zhuang, B., McDonald, J. W., Schultz, F. A. & Newton, W. E. (1984). *Organometallics*, **3**, 943–945.

Acta Cryst. (1999). **C55**, 298–300

A dimolybdenum(I) carbonyl compound with thiolate and carboxylate bridges: tetrabutylammonium bis(μ-benzenethiolato-*S:S*)hexacarbonyl-μ-pivalate-*O:O'*-dimolybdenum(Mo—Mo)

GUOHUA PAN, BOTAO ZHUANG* AND JIUTONG CHEN

Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, State Key Laboratory of Structural Chemistry, Fuzhou, Fujian 350002, People's Republic of China. E-mail: zbt@ms.fjirsm.ac.cn

(Received 10 June 1998; accepted 18 September 1998)


Abstract

In the anion of the title compound, (C₁₆H₃₆N)[Mo₂-(C₅H₉O₂)(C₆H₅S)₂(CO)₆], each Mo atom and three terminal carbonyl groups form a *fac*-Mo(CO)₃ fragment, and two Mo atoms and two benzenethiolate bridging

ligands form a planar Mo_2S_2 unit, with the two phenyl rings in an *anti* configuration. The pivalate acts as a conventional μ, η^2 -ligand bridge to two Mo atoms, to form an approximately planar five-membered ring. The two $[\text{MoOS}_2(\text{CO})_3]$ octahedra share a common S—S edge.

Comment

We have been investigating the syntheses and structures of dimolybdenum(I) compounds containing bridging thiolate, as well as other inorganic and organic ligands. It has been found that the phenyl rings of two benzenethiolate ligands adopt either *syn* or *anti* configurations when various bridging ligands are introduced to substitute the axial CO in the parent compound, $[\text{Mo}_2(\text{C}_6\text{H}_5\text{S})_2(\text{CO})_6]$, (I) (Pan *et al.*, 1998). The introduction of a halogen-ion bridging ligand leads to a *syn* configuration, while a carboxylate ligand leads to an *anti* configuration. Despite the greater steric hindrance of the pivalate in the title compound, (II), the two phenyl rings are in an *anti* configuration.



The structure of the anion of (II) is shown in Fig. 1. Each Mo atom has a distorted octahedral geometry, involving three C atoms from carbonyls, two S atoms from benzenethiolate bridging groups and one O atom from the pivalate ligand. Two *fac*- $\text{Mo}(\text{CO})_3$ fragments are linked together by two benzenethiolate and one pivalate, forming an edge-sharing bi-octahedral structure.

The structure of the Mo_2S_2 unit in (II) is similar to that in compound (I) (Zhuang *et al.*, 1995). The introduction of a pivalate ligand causes the Mo_2S_2 unit to contract slightly, resulting in smaller Mo—S—Mo angles and a shorter Mo—Mo distance, but the Mo—S distances are essentially equal to those of (I). The *trans*-Mo—C distances are considerably shorter than those *cis* to the pivalate ligand. This indicates that a tetrabridged compound substituting the axial CO in compound (II) further, with a pivalate, would be difficult to prepare.

The pivalate ligand substitutes two axial carbonyls of (I), forming an approximately planar five-membered Mo—O—C—O—Mo ring. The Mo—Mo distance is long at 2.9071 (8) Å, leading to a slightly larger O—C—O angle [125.1 (6)°], and slightly more acute O8—Mo1—Mo2 [81.13 (11)°] and O7—Mo2—Mo1 [81.45 (11)°] angles in the five-membered ring. It is worth noting that the Mo1—O8 distance [2.215 (4) Å] is slightly shorter than




Fig. 1. View of the anion of (II), with displacement ellipsoids shown at the 30% probability level and H atoms omitted for clarity.

the Mo2—O7 distance [2.234 (4) Å], but the O7—C31 distance [1.257 (7) Å] is equal to the O8—C31 distance [1.260 (7) Å].

Experimental

The title compound was synthesized by the reaction of $\text{Mo}_2(\text{CO})_8(\text{C}_6\text{H}_5\text{S})_2$ (Zhuang *et al.*, 1984; Smith *et al.*, 1987) with Bu_4NBr and $(\text{CH}_3)_3\text{CCOONa}$ (1:1:1) in acetone, and crystallized by adding *i*-PrOH to the filtrate and allowing it to stand at a temperature below 273 K for several days. Analysis calculated for $\text{C}_{39}\text{H}_{55}\text{Mo}_2\text{NO}_8\text{S}_2$: C 50.75, H 5.97, N 1.52%; found: C 50.76, H 6.01, N 1.54%; IR (KBr, cm^{-1} , ν_{CO}): 1997.9 (s), 1949.6 (s), 1930.4 (s), 1905.3 (s), 1889.9 (s), 1847.5 (s), 1820.5 (s); ν_{COO} : 1540.9 (s), 1417.4 (s).

Crystal data

$(\text{C}_{16}\text{H}_{36}\text{N})[\text{Mo}_2(\text{C}_5\text{H}_9\text{O}_2)(\text{C}_6\text{H}_5\text{S})_2(\text{CO})_6]$
 $M_r = 921.84$
 Monoclinic
 $P2_1/c$
 $a = 14.4791(2)$ Å
 $b = 12.9825(2)$ Å
 $c = 24.4658(3)$ Å
 $\beta = 104.790(1)^\circ$
 $V = 4446.58(11)$ Å³
 $Z = 4$
 $D_x = 1.377$ Mg m⁻³
 D_m not measured

Mo $K\alpha$ radiation
 $\lambda = 0.71073$ Å
 Cell parameters from 8192 reflections
 $\theta = 1.45\text{--}23.26^\circ$
 $\mu = 0.704$ mm⁻¹
 $T = 293(2)$ K
 Block
 $0.36 \times 0.32 \times 0.28$ mm
 Green

Data collection

Siemens SMART CCD diffractometer

4577 reflections with $I > 2\sigma(I)$

ω scans
 Absorption correction:
 empirical (SADABS;
 Sheldrick, 1996)
 $T_{\min} = 0.562$, $T_{\max} = 0.862$
 16 882 measured reflections
 6373 independent reflections

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.135$
 $S = 1.011$
 6373 reflections
 384 parameters
 H atoms not refined
 $w = 1/[\sigma^2(F_o^2) + (0.0512P)^2 + 11.5194P]$
 where $P = (F_o^2 + 2F_c^2)/3$

$R_{\text{int}} = 0.038$
 $\theta_{\max} = 23.26^\circ$
 $h = -16 \rightarrow 15$
 $k = 0 \rightarrow 14$
 $l = 0 \rightarrow 27$
 Intensity variation: none
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.817 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.622 \text{ e } \text{\AA}^{-3}$
 Extinction correction: none
 Scattering factors from
International Tables for
Crystallography (Vol. C)

Siemens (1994a). *SAINT. Software Reference Manual*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Siemens (1994b). *SHELXTL*. Release 5.03. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Siemens (1996). *SMART. Software Reference Manual*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Smith, D. A., Zhuang, B., Newton, W. E., McDonald, J. W. & Schultz, F. A. (1987). *Inorg. Chem.* **26**, 2524–2531.
 Zhuang, B., Huang, L. & He, L. (1995). *Chin. J. Struct. Chem.* **14**, 359–363.
 Zhuang, B., McDonald, J. W., Schultz, F. A. & Newton, W. E. (1984). *Organometallics*, **3**, 943–945.

Acta Cryst. (1999). **C55**, 300–302

trans-Diaquabis(quinoline-2-carboxylato-N,O)cobalt(II)–water–ethanol (1/2/2)

NOBUO OKABE AND TOMOMI MAKINO

Faculty of Pharmaceutical Sciences, Kinki University,
 Kowakae 3-4-1, Higashiosaka, Osaka 577, Japan. E-mail:
 okabe@phar.kindai.ac.jp

(Received 26 May 1998; accepted 19 October 1998)

Abstract

The title compound, [Co(C₁₀H₆NO₂)₂(H₂O)₂].2H₂O.2C₂H₅OH, contains a six-coordinate Co^{II} ion at a center of symmetry. The Co^{II} ion displays distorted octahedral coordination geometry defined by the two quinoline N atoms, two O atoms of the carboxylate groups and two O atoms of the water molecules. All of the corresponding pairs of ligand atoms lie in *trans* positions with respect to each other. Molecules are linked together by an intermolecular hydrogen-bonding network involving the uncoordinated water and ethanol molecules.

Comment

2-Quinolinecarboxylic acid, (I), is an intermediate tryptophan metabolite and is known to chelate transition metal ions (Martell & Smith, 1974). Crystal structures of complexes of 2-quinolinecarboxylic acid have been determined for several metal ions, including Cu^{II} (Haendler, 1986), Mn^{II} (Haendler, 1996; Okabe & Koizumi, 1997) and Fe^{II} (Okabe & Makino, 1998). We have carried out the structural analysis of the Co^{II} complex, (II), and report the results here.




Table 1. Selected geometric parameters (Å, °)

Mo1—C6	1.932 (7)	Mo2—C2	2.009 (8)
Mo1—C4	1.995 (8)	Mo2—C3	2.013 (8)
Mo1—C5	1.998 (7)	Mo2—O7	2.234 (4)
Mo1—O8	2.215 (4)	Mo2—S2	2.460 (2)
Mo1—S1	2.477 (2)	Mo2—S1	2.464 (2)
Mo1—S2	2.482 (2)	O7—C31	1.257 (7)
Mo1—Mo2	2.9071 (8)	O8—C31	1.260 (7)
Mo2—C1	1.936 (8)		
S1—Mo1—S2	107.36 (6)	Mo2—S1—Mo1	72.08 (5)
O8—Mo1—Mo2	81.13 (11)	Mo2—S2—Mo1	72.06 (5)
S2—Mo2—S1	108.45 (6)	O7—C31—O8	125.1 (6)
O7—Mo2—Mo1	81.45 (11)		

Data were collected over a hemisphere of reciprocal space by a combination of three sets of exposures. Each set had a different φ angle for the crystal and each exposure of 10 s covered 0.3° in ω . The crystal-to-detector distance was 4.95 cm. Coverage of the unique set was over 99% complete to at least 23° in θ . Crystal decay was monitored by measurement of duplicate reflections. H atoms were all located theoretically and were not refined. The lengths of some C—C bonds in the cation are shorter than is reasonable; this is due to high thermal motion and disorder.

Data collection: *SMART* (Siemens, 1996). Cell refinement: *SMART* and *SAINT* (Siemens, 1994a). Data reduction: *XPREF* in *SHELXTL* (Siemens, 1994b). Program(s) used to solve structure: *SHELXTL*. Program(s) used to refine structure: *SHELXTL*. Molecular graphics: *SHELXTL*. Software used to prepare material for publication: Siemens *SHELXTL*.

We are grateful to NNSF and SKLSC for financial support of this work.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: TA1216). Services for accessing these data are described at the back of the journal.

References

- Pan, G., Zhuang, B. & Chen, J. (1998). *Acta Cryst.* **C54**, 1580–1582.
 Sheldrick, G. M. (1996). *SADABS. Program for Absorption Corrections*. University of Göttingen, Germany.