

O4D†	0.5590 (7)	0.1900 (11)	−0.1014 (16)	0.112 (5)
C25D‡	0.5498 (6)	0.3052 (12)	−0.1053 (15)	0.094 (5)
C26D‡	0.5933 (6)	0.3475 (11)	−0.1157 (16)	0.107 (6)
C27D‡	0.6144 (6)	0.2610 (16)	−0.1601 (16)	0.123 (6)
C28D‡	0.5874 (7)	0.1710 (11)	−0.1596 (15)	0.121 (5)

† Occupancy of 0.40 (2). ‡ Occupancy of 0.60 (2).

Table 2. Selected geometric parameters (Å, °)

Cu1—N1	2.013 (3)	Cu1—O1	2.177 (2)
Cu1—O1 ⁱ	2.147 (2)	C3—C3 ⁱⁱ	1.480 (6)
Cu1—P1	2.1680 (11)		
N1—Cu1—O1 ⁱ	103.05 (10)	N1—Cu1—O1	98.77 (10)
N1—Cu1—P1	129.54 (8)	O1 ⁱ —Cu1—O1	73.82 (9)
O1 ⁱ —Cu1—P1	120.80 (7)	P1—Cu1—O1	115.56 (7)

Symmetry codes: (i) $\frac{1}{2} - x, \frac{1}{2} - y, -z$; (ii) $\frac{1}{2} - x, \frac{3}{2} - y, -z$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989). Cell refinement: *CAD-4 Software*. Data reduction: *CAD-4 Software*, *PROFILE* (Blessing, Coppens & Becker, 1974). Program(s) used to solve structure: *SHELXS86* (Sheldrick, 1990). Program(s) used to refine structure: *SHELXL93* (Sheldrick, 1993). Molecular graphics: *ORTEPII* (Johnson, 1971). Software used to prepare material for publication: *CIFTAB* in *SHELXL93*.

The authors thank the School of Chemical Sciences Materials Characterization Laboratory at the University of Illinois for X-ray data collection, Dr Scott R. Wilson for his assistance in the structure solution, the US Department of Energy through the Materials Research Laboratory at the University of Illinois (grant DEFG02-91-ER45439) and the National Science Foundation (grant CHE-94-23121) for financial assistance. Additional support from 3M Company and the Camille Dreyfus Teacher–Scholar Awards Program is gratefully acknowledged.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates, complete geometry and torsion angles have been deposited with the IUCr (Reference: FG1111). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Blessing, R. H., Coppens, P. & Becker, P. (1974). *J. Appl. Cryst.* **7**, 488–492.
 Cotton, F. A. & Wilkinson, G. (1988). *Advanced Inorganic Chemistry*, 5th ed., p. 1289. New York: Wiley Interscience.
 Enraf–Nonius (1989). *CAD-4 Software*. Version 5. Enraf–Nonius, Delft, The Netherlands.
 Gysling, H. J. & Kubas, G. J. (1979). *Inorg. Synth.* **19**, 92–94.
 Johnson, C. K. (1971). *ORTEPII*. ORNL-3794, revised. Oak Ridge National Laboratory, Tennessee, USA.
 MacGillivray, L. R., Subramanian, S. & Zaworotko, M. J. (1994). *J. Chem. Soc. Chem. Commun.* pp. 1325–1326.
 Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
 Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
 Yaghi, O. M. & Li, G. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 207–209.

Acta Cryst. (1996). **C52**, 2178–2180

(Cyclohexylamine-*N*){2-[(2-thiophenyl)iminomethyl]phenolato-*O,N,S*}nickel(II)

M. NAWAZ TAHIR,^a DİNÇER ÜLKÜ,^{a*} ORHAN ATAKOL^b
 AND ADNAN KENAR^b

^aDepartment of Engineering Physics, Hacettepe University, Beytepe 06532, Ankara, Turkey, and ^bDepartment of Chemistry, Faculty of Science, University of Ankara, Beşevler 06100, Ankara, Turkey. E-mail: dulku@eti.cc.hun.edu.tr


(Received 28 November 1995; accepted 13 May 1996)

Abstract

The crystal of the title compound is composed of discrete neutral [Ni(C₁₃H₉NOS)(C₆H₁₃N)] molecules. The Ni ion has a distorted square-planar coordination. The bond distances between the Ni atom and the donor N2, S1, N1 and O1 atoms are 1.948 (5), 2.148 (2), 1.874 (5) and 1.845 (5) Å, respectively. Although the bond angles around the Ni atom range from 83.4 (2) to 96.1 (2)°, their sum is 360.1°. The cyclohexyl ring is disordered.

Comment

N-(2-Hydroxyphenyl)salicylaldimine and its thio derivatives are interesting tridentate ligands. With metals of the first transition series such as Fe²⁺, Ni²⁺, Cu²⁺ and Co²⁺, these ligands form bridged dimeric complexes which have been the subject of attention from coordination chemists. If amine is present in the reaction, Ni²⁺ alone of these ions has a tendency to form mononuclear square-planar complexes instead of dimers. The crystal structures of Ni²⁺–Schiff base complexes with different ligands and amines have been reported previously (Elerman, Fuess & Paulus, 1991; Soriano-García, Toscano, Valdés-Martínez & Fernández-G., 1985; Elerman, Paulus & Fuess, 1991; Kabak, Elerman, Özbey & Atakol, 1995). We report here a new mononuclear Ni²⁺ complex, (cyclohexylamine-*N*){2-[(2-thiophenyl)iminomethyl]phenolato-*O,N,S*}nickel(II), (I).



The structure consists of discrete [Ni(C₁₃H₉NOS)-(C₆H₁₃N)] molecules (Fig. 1). The Ni²⁺ ion is located in a slightly distorted square-planar environment.

The S1—Ni—N1 and S1—Ni—N2 angles are 89.6(2) and 91.0(2)°, respectively, while the O1—Ni—N2 and O1—Ni—N1 angles, with their respective values of 83.4(2) and 96.1(2)°, deviate a little more from 90°. The sum of the bond angles around the Ni atom is 360.1°. The Ni—N distances [1.874(5) and 1.948(5) Å], as well as the Ni—S1 [2.148(2) Å] and Ni—O1 [1.845(5) Å] distances, are not significantly different from those observed in similar compounds (Soriano-García, Toscano, Valdés-Martínez & Fernández-G., 1985; Gyepes, Pavelčík & Beno, 1981; Kabak, Elerman, Özbey & Atakol, 1995). The observation of two different Ni—N distances is an indication of the already well established fact (Curtis, 1979) that the Ni—N distance depends on the nature of the N-donor atom and also the number of atoms in the chelate rings. The dihedral angle between the chelate rings is 6(1)°. The phenyl rings make a dihedral angle of 10.9(8)° with one another and are practically coplanar with their respective chelate rings. Planes A (C14A, C15A, C17A, C18A) and B (C14B, C15B, C17B, C18B) make dihedral angles of 113.7(4) and 85.4(4)°, respectively, with the coordination plane around the Ni atom. The dihedral angle between planes A and B is 28(1)°. The distances of the two *para* positioned chair atoms, C16A/C19A and C16B/C19B, from their respective planes are 0.52(2)/−0.29(1) and −0.39(2)/0.56(2) Å.




Fig. 1. An ORTEP (Johnson, 1965) drawing of the title complex with the atom-numbering scheme. For clarity, only one component of the disordered cyclohexyl ring is shown; the C atoms of the other component carry suffix A instead of B. The displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small circles of arbitrary radii.

Experimental

The Schiff base ligand was obtained from salicylaldehyde and 2-aminothiophenol by a condensation reaction in ethanol solution. The ligand (0.115 g, 0.0005 mol) was dissolved in hot acetonitrile (50 ml) and cyclohexylamine (1.0 ml) was added to the resulting solution. A solution of $[\text{Ni}(\text{CH}_3\text{COO})_2] \cdot 4\text{H}_2\text{O}$ (0.125 g, 0.0005 mol) in hot methanol (30 ml) was then added and the resulting solution was mixed and set aside for 48 h. The precipitated crystals were filtered and used for X-ray data collection.

Crystal data

$[\text{Ni}(\text{C}_{13}\text{H}_9\text{NOS})(\text{C}_6\text{H}_{13}\text{N})]$
 $M_r = 385.17$
 Monoclinic
 $P2_1/n$
 $a = 14.293(1) \text{ Å}$
 $b = 5.980(1) \text{ Å}$
 $c = 21.660(2) \text{ Å}$
 $\beta = 106.58(4)^\circ$
 $V = 1774.4(6) \text{ Å}^3$
 $Z = 4$
 $D_x = 1.442 \text{ Mg m}^{-3}$
 D_m not measured

Mo $K\alpha$ radiation
 $\lambda = 0.71073 \text{ Å}$
 Cell parameters from 25 reflections
 $\theta = 10.05\text{--}18.08^\circ$
 $\mu = 1.22 \text{ mm}^{-1}$
 $T = 295 \text{ K}$
 Needle
 $0.45 \times 0.15 \times 0.10 \text{ mm}$
 Red

Data collection

Enraf–Nonius CAD-4 diffractometer
 $\omega/2\theta$ scans
 Absorption correction: empirical via ψ scans (MolEN; Fair, 1990)
 $T_{\min} = 0.913$, $T_{\max} = 0.999$
 2942 measured reflections
 2782 independent reflections

2074 observed reflections
 $[I > 3\sigma(I)]$
 $R_{\text{int}} = 0.012$
 $\theta_{\max} = 24.31^\circ$
 $h = -15 \rightarrow 16$
 $k = -6 \rightarrow 0$
 $l = -25 \rightarrow 0$
 3 standard reflections
 frequency: 120 min
 intensity decay: 0.6%

Refinement

Refinement on F
 $R = 0.052$
 $wR = 0.058$
 $S = 1.04$
 2074 reflections
 211 parameters
 Weighting scheme: if $F \geq 95.32$, $w = (95.32/F)^2$; if $F^2 < 3\sigma F^2$, $w = 0$

$(\Delta/\sigma)_{\max} = 0.0003$
 $\Delta\rho_{\max} = 0.70 \text{ e Å}^{-3}$
 $\Delta\rho_{\min} = -0.56 \text{ e Å}^{-3}$
 Extinction correction: none
 Atomic scattering factors from *International Tables for X-ray Crystallography* (1974, Vol. IV)

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters (Å^2)

$$B_{\text{eq}} = (8\pi^2/3) \sum_i \sum_j U_{ij} a_i^* a_j^* \cdot \mathbf{a}_i \cdot \mathbf{a}_j$$

	<i>x</i>	<i>y</i>	<i>z</i>	<i>B</i> _{eq}
Ni	0.55660(6)	0.0501(2)	0.36678(4)	3.66(2)
S1	0.5021(1)	0.2648(3)	0.42812(8)	4.75(4)
O1	0.6107(3)	−0.1490(8)	0.3216(2)	4.3(1)
N1	0.5035(3)	0.2426(9)	0.2974(2)	3.6(1)
N2	0.6097(4)	−0.156(1)	0.4379(2)	4.4(1)
C1	0.6145(4)	−0.133(1)	0.2624(3)	3.9(1)
C2	0.6627(4)	−0.300(1)	0.2373(3)	4.7(2)
C3	0.6734(5)	−0.283(1)	0.1766(3)	5.3(2)
C4	0.6354(5)	−0.100(1)	0.1373(3)	5.5(2)

C5	0.5866 (5)	0.061 (1)	0.1601 (3)	5.2 (2)
C6	0.5741 (4)	0.050 (1)	0.2221 (3)	3.9 (1)
C7	0.5209 (4)	0.223 (1)	0.2421 (3)	4.1 (1)
C8	0.4435 (4)	0.425 (1)	0.3066 (3)	3.7 (1)
C9	0.3932 (4)	0.572 (1)	0.2589 (3)	4.5 (2)
C10	0.3396 (5)	0.745 (1)	0.2727 (3)	5.1 (2)
C11	0.3350 (5)	0.776 (1)	0.3347 (3)	5.3 (2)
C12	0.3836 (5)	0.631 (1)	0.3829 (3)	4.8 (2)
C13	0.4376 (4)	0.452 (1)	0.3694 (3)	4.0 (1)
C14A†	0.7206 (6)	−0.147 (2)	0.4655 (4)	4.0 (2)
C15A†	0.757 (1)	−0.346 (4)	0.5085 (9)	4.7 (4)
C16A†	0.861 (1)	−0.311 (3)	0.5477 (7)	8.4 (4)
C17A†	0.884 (1)	−0.094 (3)	0.5771 (7)	8.1 (4)
C18A†	0.855 (2)	0.112 (4)	0.536 (1)	6.0 (5)
C19A†	0.7568 (8)	0.083 (2)	0.4875 (5)	5.7 (3)
C14B†	0.697 (1)	−0.103 (3)	0.4912 (8)	4.1 (4)
C15B†	0.7399 (7)	−0.305 (2)	0.5288 (5)	5.1 (2)
C16B†	0.832 (2)	−0.267 (4)	0.582 (1)	5.5 (5)
C17B†	0.903 (1)	−0.114 (3)	0.5653 (8)	4.1 (4)
C18B†	0.8735 (8)	0.071 (2)	0.5117 (5)	5.6 (2)
C19B†	0.776 (1)	0.003 (4)	0.463 (1)	5.5 (5)

† Atoms with suffix A have occupancies of 0.66 and those with suffix B have occupancies of 0.34.

Table 2. Selected geometric parameters (Å, °)

Ni—S1	2.148 (2)	Ni—N1	1.874 (5)
Ni—O1	1.845 (5)	Ni—N2	1.948 (5)
S1—Ni—O1	174.2 (1)	O1—Ni—N1	96.1 (2)
S1—Ni—N1	89.6 (2)	O1—Ni—N2	83.4 (2)
S1—Ni—N2	91.0 (2)	N1—Ni—N2	178.7 (2)

The H atoms of the Schiff base ligand were attached to their C atoms geometrically (C—H 0.95 Å). The H atoms of the cyclohexylamine ligand were not attached. Displacement parameters of all H atoms were constrained at 1.3 times those of the corresponding atoms and a riding model was used for all H atoms. During the early stages of refinement, disorder was evident for the cyclohexyl ring. Resolution of the disorder was made possible by restraining all the bonding C—C (1.54 Å) and the non-bonding C14...C18 and C15...C17 (2.51 Å) distances in the ring. After some isotropic refinement of the disordered C atoms, without the H atoms, new coordinates were obtained and these were kept fixed to refine the relative occupancy factors. At this point, a difference map started to show the second ring. The coordinates of the second ring were treated in the same way as described above. Finally, both components were refined isotropically with the coordinates of some atoms being constrained (C16B, C17A, C18A, C17B, C19B).

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1993). Data reduction: *MolEN* (Fair, 1990). Program(s) used to solve structure: *SIMPEL* in *MolEN*. Program(s) used to refine structure: *LSFM* in *MolEN*. Molecular graphics: *ORTEP* (Johnson, 1965) in *MolEN*. Software used to prepare material for publication: *MolEN*.

The authors acknowledge the purchase of the CAD-4 diffractometer under Grant DPT/TBAG1 of the Scientific and Technical Research Council of Turkey.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates, complete geometry and torsion angles have been deposited with the IUCr (Reference: BM1052). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Curtis, N. F. (1979). *Coordination Chemistry of Macrocyclic Compounds*, edited by G. A. Melson, pp. 219–344. New York: Plenum.
- Elerman, Y., Fuess, H. & Paulus, E. F. (1992). *Acta Cryst.* **C48**, 40–42.
- Elerman, Y., Paulus, E. F. & Fuess, H. (1991). *Acta Cryst.* **C47**, 70–72.
- Enraf–Nonius (1993). *CAD-4 EXPRESS*. Version 1.1. Enraf–Nonius, Delft, The Netherlands.
- Fair, C. K. (1990). *MolEN. An Interactive Intelligent System for Crystal Structure Analysis*. Enraf–Nonius, Delft, The Netherlands.
- Gyepes, E., Pavelčík, F. & Beno, A. (1981). *Collect. Czech. Chem. Commun.* **46**, 975–981.
- Johnson, C. K. (1965). *ORTEP*. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee, USA.
- Kabak, M., Elerman, Y., Özbey, S. & Atakol, O. (1995). *J. Chem. Crystallogr.* **25**, 259–262.
- Soriano-García, M., Toscano, R. A., Valdés-Martínez, J. & Fernández-G., J. M. (1985). *Acta Cryst.* **C41**, 498–500.

Acta Cryst. (1996). **C52**, 2180–2183

[N-(2-Amino-κN-ethyl)piperaziniumyl-N¹]-tetrachloroindium(III)

ANN M. CHIPPINDALE, SARAH J. BRECH, ANDREW R. COWLEY AND ALEXANDER N. CHERNEGA

Chemical Crystallography Laboratory, 9 Parks Road, Oxford, OX1 3PD, England. E-mail: kryst4@vax.ox.ac.uk

(Received 15 March 1996; accepted 8 May 1996)

Abstract

The crystal structure of the title compound consists of molecules of [InCl₄{(NH₂CH₂CH₂)C₄H₁₀N₂}] held together in buckled sheets by a network of hydrogen bonds. The In atom has a distorted octahedral coordination geometry with the *N*-(2-aminoethyl)piperazinium cation behaving as a bidentate ligand.

Comment

Few crystal structures of octahedral indium(III) mixed-ligand chloro complexes of the form [InCl_xL_{6-x}]^{3-x}, where *L* is a coordinating ligand with an N-donor atom, have been reported in the literature. Examples include, for *x* = 1, [HB(3,5-Me₂pz)₃In{H₂B(pz)₂}Cl] (pz is pyrazolyl) (Reger, Mason, Reger, Rheingold & Ostrander, 1994), for *x* = 2, [NHEt₃][InCl₂(dpt)₂] (dpt is 1,3-diphenyltriazene), [InCl₂(dpt)(bipy)] and [InCl₂(dpt)(3,5-Me₂py)₂] (bipy is bipyridyl, py is pyridyl) (Leman, Roman & Barron, 1992), for *x* = 3, [mer-InCl₃(py)₃].py (Jeffs, Small & Worrall, 1984) and [{HB(pz)₃}₂InCl₃][{HB(pz)₃}₂In].thf.CH₂Cl₂ (thf