



## Evidence for the Involvement of a Sulfurane Intermediate in the Oxidation of Simple Sulfides by Methyl(trifluoromethyl)dioxirane.

Gregorio Asensio,\* Rossella Mello, and Maria Elena González-Núñez

*Departamento de Química Orgánica. Facultad de Farmacia. Universidad de Valencia.*

*Avda. Vicente Andrés Estellés s.n. E-46100 Burjassot (Valencia) Spain*

**Abstract:** Methyl(trifluoromethyl)dioxirane reacts with sulfides to give preferentially sulfones, even in the presence of competing sulfoxides. The sulfoxide yield increases at the expense of the sulfone when 2,2,2-trifluoroethanol is used as co-solvent. The reaction of dioxirane **1b** with phenylmethyl sulfide in the presence of 1,1,1-trifluoropropanone-<sup>18</sup>O-hydrate (48% atom labelled), leads to the obtention of 23% atom <sup>18</sup>O-labelled sulfoxide and 6.1% atom <sup>18</sup>O-labelled sulfone. The involvement of a cyclic hypervalent sulfurane intermediate is proposed as reactive intermediate.

The oxidation of sulfides is one among the earliest studied O-transfer reactions of dioxiranes.<sup>1</sup> Dimethyldioxirane (**1a**) (hereafter DMDO) reacts with sulfides<sup>2,3</sup> to give sulfoxides, which can further react with DMDO (**1a**) to give sulfones in a sequential process (Equation 1). Since sulfides are better nucleophiles than sulfoxides, the oxidation can be controlled at the sulfoxide stage when the stoichiometric amount of DMDO (**1a**) is used. Now we report the first detailed study on the oxidation



**Equation 1**

of simple sulfides and sulfoxides by methyl(trifluoromethyl)dioxirane (**1b**) (hereafter TFDO), a dioxirane that exhibits a stronger electrophilic character.<sup>1,3</sup> The results of the oxidation of phenylmethyl sulfide (**2**) with TFDO (**1b**) to afford phenylmethyl sulfoxide (**3**) and phenylmethyl sulfone (**4**) at 0 °C in different solvents are shown in Table 1.<sup>4,5</sup> Surprisingly, the oxidation of **2** by TFDO (**1b**) produced **4** as the main product, even in the presence of a large excess of **2** relative to TFDO (**1b**). It should be noted that the ratio sulfone/sulfoxide (**4/3**) decreases in the presence of trifluoroethanol. The oxidation of dibenzyl sulfide and dibutyl sulfide by TFDO (**1b**) proceeded with similar results. Methyl or trifluoromethyl esters resulting from the radical chain decomposition of **1a**<sup>6</sup> or **1b**<sup>5</sup> were not detected in any case allowing us to disregard a radical mechanism for these oxidations. By analogy with the case of the oxidations carried with DMDO, the oxidation of sulfides to sulfones by TFDO were expected to proceed in a sequential fashion as depicted in Eq. 1.<sup>3</sup> In that case, from our results, phenylmethyl sulfoxide (**3**) appears to react with TFDO (**1b**) faster than sulfide **2**.<sup>7</sup> In order to clarify this surprising conclusion, we performed a series of competitive oxidations of a pair sulfide/sulfoxide bearing minor structural differences, namely phenyl(trideuteriomethyl) sulfide (**2-d<sup>3</sup>**) and phenylmethyl sulfoxide (**3**), with TFDO (**1b**). Reactions were carried out as described and the products analyzed by GC-MS. The results are shown in Table 2.

The reaction of an equimolar mixture of **2-d<sup>3</sup>** and **3** with DMDO (**1a**) yielded the sulfoxide **3-d<sup>3</sup>** and small amounts of the sulfones **4-d<sup>3</sup>** and **4** while most of the sulfoxide **3** remained unchanged (entry 1, Table 2). By contrast, when a mixture of sulfide **2-d<sup>3</sup>** and sulfoxide **3** reacted with TFDO (**1b**), *sulfone 4-d<sup>3</sup> was the main oxidation product from sulfide 2-d<sup>3</sup>* even in the presence of excess of sulfoxide **3** (entries 2 and 3, Table 2).<sup>8</sup> We should remark that, although sulfoxide **3** is also oxidized

by TFDO (**1b**) to sulfone **4**, oxygen transfer occurs

*preferentially* to sulfide **2-d<sup>3</sup>**. The same trend is

observed in trifluoroethanol solution, but in this

case the **4-d<sup>3</sup>/3-d<sup>3</sup>** ratio decreases (entry 4, Table

2). These data allow us to conclude that the

conversion of sulfide **2** into sulfone **4** by TFDO (**1**)

cannot occur *only* through the oxidation of the

intermediate sulfoxide **3** as depicted in Eq. 1, and

suggest the involvement of a sulfide-derived

reactive intermediate that, *i*) is oxidized by TFDO

(**1b**) faster than either sulfides or sulfoxides and, *ii*)

is readily converted into a sulfoxide by the action of

protic acidic solvents such as trifluoroethanol.

Then, we designed <sup>18</sup>O-tracer experiments

addressed to trap the suspected intermediate (**5**).

We carried out the oxidation of 0.031 M

phenylmethyl sulfide (**2**) at 0 °C under inert

**Table 1.** Oxidation of Phenyl Methyl Sulphide (**2**) by TFDO (**1b**)<sup>a</sup>

| run | 2:1b <sup>b</sup> | solvent <sup>c</sup> | reaction mixture (%) <sup>d</sup> |    |    | 4/3 ratio |
|-----|-------------------|----------------------|-----------------------------------|----|----|-----------|
|     |                   |                      | 2                                 | 3  | 4  |           |
| 1   | 1:1               | DC                   | 42                                | 12 | 46 | 3.8       |
| 2   | 10:1              | DC                   | 93                                | 3  | 4  | 1.3       |
| 3   | 2:1               | DC                   | 68                                | 9  | 23 | 2.6       |
| 4   | 2:1               | DC/AC 1:19           | 71                                | 8  | 21 | 2.6       |
| 5   | 2:1               | DC/AN 1:19           | 68                                | 9  | 23 | 2.6       |
| 6   | 2:1               | DC/TFE 1:7           | 66                                | 18 | 16 | 0.9       |

<sup>a</sup> The reactions were carried out at 0 °C with an initial concentration of **1b** ranging between 0.030 to 0.063 M; identical results were obtained when TFDO solution was added dropwise. <sup>b</sup> Molar ratio. <sup>c</sup> DC: dichloromethane; AC: acetone; AN: acetonitrile; TFE: trifluoroethanol; <sup>d</sup> Values are the average of at least two identical runs within a standard error of ±2%.

atmosphere with the equimolar amount of TFDO (**1b**) in 4:1 CH<sub>3</sub>CN/CH<sub>2</sub>Cl<sub>2</sub> in the presence of a 100-fold excess of 1,1,1-trifluoroacetone-<sup>18</sup>O-hydrate (**6-<sup>18</sup>O**, 49 atom % labelled<sup>9</sup>), a suitable acidic reagent for trapping of a polar intermediate. Effectively, under these conditions GC-MS analysis of the reaction mixture revealed<sup>10</sup> that a 23.0 atom % of


**Table 2.** Competitive Oxidations of PhSCD<sub>3</sub> (**2-d<sup>3</sup>**) and PhSOCH<sub>3</sub> (**3**) by Dioxiranes **1**.<sup>a</sup>

| entry | reagents                    | 1         | molar ratio            | solvent <sup>b</sup> | reaction mixture (%) <sup>c</sup> |                               |                               | O-transfer <sup>d</sup> | ratio                              |
|-------|-----------------------------|-----------|------------------------|----------------------|-----------------------------------|-------------------------------|-------------------------------|-------------------------|------------------------------------|
|       |                             |           |                        |                      | 2-d <sup>3</sup>                  | 3-d <sup>3</sup> ( <b>3</b> ) | 4-d <sup>3</sup> ( <b>4</b> ) |                         |                                    |
|       |                             |           | 1: 2-d <sup>3</sup> :3 |                      |                                   |                               |                               | 2-d <sup>3</sup> / 3    | 4-d <sup>3</sup> /3-d <sup>3</sup> |
| 1     | 2-d <sup>3</sup> + <b>3</b> | <b>1a</b> | 1:1:1                  | DC/AC 1:1            | 13                                | 80 (93)                       | 7 (7)                         | 13.4                    | 0.1                                |
| 2     | 2-d <sup>3</sup> + <b>3</b> | <b>1b</b> | 1:1:1                  | DC                   | 58                                | 11 (66)                       | 31 (34)                       | 2.1                     | 2.8                                |
| 3     | 2-d <sup>3</sup> + <b>3</b> | <b>1b</b> | 1:1:3                  | DC                   | 73                                | 9 (80)                        | 18 (20)                       | 2.3                     | 2.0                                |
| 4     | 2-d <sup>3</sup> + <b>3</b> | <b>1b</b> | 1:1:1                  | DC/TFE 1:9           | 53                                | 21 (70)                       | 26 (30)                       | 2.4                     | 1.2                                |

<sup>a</sup> Reactions carried out at 0 °C with a concentration of the substrates ranging between 0.03-0.06 M. <sup>b</sup>DC: Dichloromethane; AC: acetone; TFE: trifluoroethanol. <sup>c</sup> Mixture composition have been calculated separately for CD<sub>3</sub>- and CH<sub>3</sub>- substituted compounds and the values are the average of at least two identical runs within a standard error of ±2%. <sup>d</sup> Calculated as [(3-d<sup>3</sup>) + 2x(4-d<sup>3</sup>)]/[4].


the sulfoxide (34 % yield) and a 6.1 atom % of the sulfone (66% yield) were  $^{18}\text{O}$ -labelled with a 56% sulfide conversion. In control experiments it was ascertained that sulfoxide **3** and sulfone **4** do not undergo oxygen exchange with 1,1,1-trifluoropropanone- $^{18}\text{O}$ -hydrate ( $6\text{-}^{18}\text{O}$ ) under identical conditions.

To explain all the above observations we propose the involvement of the cyclic sulfurane<sup>11</sup> **5** in the oxidation of sulfides to sulfones (Scheme 1). Sulfurane **5** would lead to sulfoxide **3** or sulfone **4** either by  $\beta$ -elimination of



**Scheme 1**

1,1,1-trifluoropropanone (*path a*, Scheme 1) or by further oxidation by TFDO (*path b*, Scheme 1). The influence of the solvent in the product distribution and the  $^{18}\text{O}$ -incorporation in the tracer experiments are explained by the known ligand exchange reactivity shown by sulfuranes<sup>13</sup> when reacting with active hydrogen substrates, which derives from their basic character due to the negative charge density localised on the oxygen atoms. So, the reaction of **5** with the acidic solvent trifluoroethanol (run 6, Table 1) would promote ring opening to afford sulfoxide **3** and trifluoroacetone (*path a*, Scheme 1). On the other hand, the formation of  $7\text{-}^{18}\text{O}$  by ligand exchange of **5** with  $6\text{-}^{18}\text{O}$  (Scheme 2) accounts for the sulfoxides



**Scheme 2**

$3\text{-}^{18}\text{O}$  and **3** upon  $\beta$ -elimination of 1,1,1-trifluoropropanone or 1,1,1-trifluoropropanone- $^{18}\text{O}$ , respectively. Since 23%  $^{18}\text{O}$ -labelled sulfoxide is formed, the sulfoxide should derive almost exclusively from the sulfurane  $7\text{-}^{18}\text{O}$ , from which the expected maximum statistical  $^{18}\text{O}$ -label incorporation in the sulfoxide is 24.5%. Additional alternative routes are suggested in Scheme 2 to account for the formation of sulfone.

In conclusion, the oxidation of simple sulfides by TFDO (**1b**) to give sulfones does not proceed, at least as the main route, through oxidation of an initially formed sulfoxide. Furthermore, solvent effects and  $^{18}\text{O}$ -labelling techniques suggest for this process the involvement of a highly reactive intermediate, most likely the hypervalent sulfurane **5**. The mechanistic

details of the oxygen transfer step from dioxirane **1b** to sulfuranes **5** and/or **7** to give sulfone are presently under investigation.

**Acknowledgement.** This work was supported by the Spanish Dirección General de Investigación Científica y Técnica (PB93-0681). We thank the S.C.S.I.E. de la Universidad de Valencia for the use of NMR and GC-MS facilities. We wish to thank Prof. J. O. Edwards and Prof. R. Curci for useful discussions.

## REFERENCES AND NOTES

- Adam, W.; Hadjirapoglou, L. P.; Curci, R.; Mello, R. in *Organic Peroxides*; Ando, W. Ed.; Wiley Interscience, Chichester, U.K., 1992, Chapter 4, pp195-219. (b) Murray, R. W. *Chem. Rev.* **1989**, *89*, 1187.
- (a) Murray, R. W.; Jeyaraman, R.; Pillay, M. K. *J. Org. Chem.* **1987**, *52*, 746. (b) Murray, R. W.; Jeyaraman, R. *J. Org. Chem.* **1985**, *50*, 2847.
- (a) Adam, W.; Golsch, D. *Chem. Ber.* **1994**, *127*, 1111. (b) Ballistreri, R. P.; Tomaselli, G. A.; Toscano, R. M.; Bonchio, M.; Conte, V.; Di Furia, F. *Tetrahedron Lett.* **1994**, *35*, 8041.
- Reactions were carried out at 0 °C under inert atmosphere by adding in one batch in the dark a thermostated dioxirane **1b** solution (0.3-0.6 M) in ketone-free methylene chloride<sup>5</sup> to a stirred solution of the sulfide in the selected solvent. The analysis of the reaction mixtures was performed by GC and <sup>1</sup>H NMR.
- Adam, W.; Curci, R.; González-Núñez, M. E.; Mello R. *J. Am. Chem. Soc.* **1991**, *113*, 7654.
- (a) Bravo, A.; Fontana, F.; Fronza, G.; Mele, A.; Minisci, F. *J. Chem. Soc. Chem. Commun.* **1995**, 1573. (b) Bravo, A.; Fontana, F.; Fronza, G.; Minisci, F.; Serri, A. *Tetrahedron Lett.* **1995**, *36*, 6945.
- McDouall, J. J. W. *J. Org. Chem.* **1992**, *57*, 2861.
- A relative reaction rate value  $k_{if}/k_D$  of 0.94 ± 3% was obtained in the competitive oxidation of sulfoxides **3**:**3-d** by TFDO.
- 1,1,1-Trifluoroacetone-<sup>18</sup>O-hydrate (6-<sup>18</sup>O) was generated *in situ* by adding to a solution of 1,1,1-trifluoroacetone in CH<sub>3</sub>CN/CH<sub>2</sub>Cl<sub>2</sub>, under inert atmosphere at 0 °C, an equimolar amount of H<sub>2</sub><sup>18</sup>O (98 atom %, supplied by Aldrich). MS analysis showed that the incorporation of H<sub>2</sub><sup>18</sup>O into the hydrate 6-<sup>18</sup>O was complete.
- <sup>18</sup>O-Labelled % was calculated by applying the equation  $100 \times [(I+2)-(I+2)_{natural}] / [I+(I+2)]$  to the normalized relative intensities corresponding to the peaks 125 (M<sup>+</sup>-CH<sub>3</sub>) and 140 (M<sup>+</sup>) for the sulfoxide **3**/**3-<sup>18</sup>O**, and to the peaks 141 (M<sup>+</sup>-CH<sub>3</sub>) and 156 (M<sup>+</sup>) for the sulfone **4**/**4-<sup>18</sup>O**, respectively, and obtaining the average value. Mass spectra were recorded at least three times. The estimated error was less than 1%.
- The involvement of an hypervalent sulfur adducts has been proposed in the oxidation of sulfides with a variety of peroxidic oxidants.<sup>12</sup> By analogy with dioxetanes,<sup>12c</sup> the intermediacy of a sulfurane intermediate has been suggested in the oxidation of sulfides by dioxiranes,<sup>1</sup> although it has not been demonstrated up to now. An alternative zwitterionic structure such as PhMeS<sup>+</sup>-OCCH<sub>3</sub>CF<sub>3</sub>O<sup>-</sup> could be depicted.
- (a) Watanabe, Y.; Kuriki, N.; Ishiguro, K.; Sawaki, Y. *J. Am. Chem. Soc.* **1991**, *113*, 2677. (b) Clennan, E. L.; Yang, K. *J. Am. Chem. Soc.* **1990**, *112*, 4044. (c) Campbell, B. S.; Denney, D. B.; Denney, D. Z.; Shih, L. S. *J. Am. Chem. Soc.* **1975**, *97*, 3850.
- (a) Hayes, R. A.; Martin, J. C. In *Organic Sulfur Chemistry*, Bernardi, F.; Csizmadia, I. G.; Mangini, A.; Eds.; Elsevier: Amsterdam, 1985. (b) Adzima, L. J.; Duesler, E. N.; Martin, J. C. *J. Am. Chem. Soc.* **1977**, *99*, 4001. (c) Martin, L. D.; Martin, J. C. *J. Am. Chem. Soc.* **1977**, *99*, 3511. (d) Martin, J. C.; Franz, J. A. *J. Am. Chem. Soc.* **1975**, *97*, 583. (e) Martin, J. C.; Franz, J. A.; Arhart, R. J. *J. Am. Chem. Soc.* **1974**, *96*, 4604. (f) Franz, J. A.; Martin, J. C. *J. Am. Chem. Soc.* **1973**, *95*, 2017. (g) Martin, J. C.; Arhart, R. J. *J. Am. Chem. Soc.* **1972**, *94*, 5003.

(Received in UK 19 January 1996; revised 8 February 1996; accepted 9 February 1996)