

CONDENSED HETEROAROMATIC RING SYSTEMS. XXIII.¹

A CONCISE SYNTHESIS OF HIPPADINE

Takao Sakamoto,*^{a)} Akito Yasuhara,^{a)} Yoshinori Kondo,^{a)} and Hiroshi Yamanaka^{b)}

a) Pharmaceutical Institute, Tohoku University, Aobayama, Aoba-ku, Sendai 980, Japan; b) Department of Education, Yamagata University, 1-4-12, Koshirakawa-cho, Yamagata 990, Japan

Abstract—Hippadine, a pyrrolophenanthridine alkaloid, was synthesized in 39% overall yield of three steps using the palladium-catalyzed cross-coupling reaction of 2,6-dibromoaniline with (*Z*)-1-tributylstannyl-2-ethoxyethene as a key step.


Previously, we reported that the palladium-catalyzed reaction of aryl halides with (*Z*)-1-tributylstannyl-2-ethoxyethene yielded (*Z*)-2-ethoxyethenylarenes which are arylacetaldehyde equivalent derivatives.^{2,3} Application of the palladium-catalyzed reaction to *ortho*-substituted haloaromatics made the construction of various condensed heterocycles possible. Especially, pyrrole ring-fused heterocycles such as indole,³ pyrrolopyridine,¹ and pyrrolopyrimidine⁴ were easily synthesized by this methodology.

Utilizing the indole formation as a key reaction, we report here a concise synthesis of hippadine, a pyrrolophenanthridine alkaloid isolated from *Crinum* species (*Amaryllidaceae*).⁵

The palladium-catalyzed reaction of commercially available 2,6-dibromoaniline (**1**) with (*Z*)-1-tributylstannyl-2-ethoxyethene (**2**) under the reported conditions^{2,3} followed by cyclization of the coupling product (**3**) with oxalic acid gave 7-bromoindole (**4**) in 96% yield. To the best of our

knowledge, this reaction is the simplest and most high yield synthetic method of 7-bromoindole from the commercially available material at present.

The *N*-acylation of **4** with 2-bromo-3,4-methylenedioxybenzoyl chloride gave the 1-benzoylated indole (**6**) in 72% yield, which was cyclized to hippadine under Stille's conditions⁶ in 68% yield. The overall yield from the commercially available starting material (**1**) was 39%.



Scheme 1

A synthesis of hippadine was reported by some groups.⁷⁻¹⁰ For example, recent Snieckus's synthesis from *N*-acetylindoline (5 steps, 24% overall yield) used palladium-catalyzed arylation of 7-iodo-*N*-acetylindoline with a benzeneboric acid derivative as a key reaction.

EXPERIMENTAL

7-Bromoindole (**4**)

A mixture of 2,6-dibromoaniline (1.00 g, 4 mmol), (*Z*)-1-tributylstannyl-2-ethoxyethene (**2**) (1.44 g, 4 mmol), Pd(PPh₃)₂Cl₂ (0.16 g, 0.23 mmol), Et₄NCl (0.66 g, 4 mmol), and MeCN (20 ml) was refluxed for 4 h. To the reaction mixture was added (COOH)₂·2H₂O (200 mg) and the mixture was

refluxed for 3 h. After removal of the solvent, the residue was purified by silica gel column chromatography using hexane/CH₂Cl₂ (1:1) to give colorless needles which were recrystallized from hexane. mp 44°C. Lit.,¹¹ mp 45-45.5°C. 60 MHz ¹H-Nmr (CDCl₃) δ (ppm): 4.2-4.4 (1H, br), 6.5-7.4 (5H, m).

7-Bromo-1-(2-bromo-4,5-methylenedioxybenzoyl)indole (6)

A mixture of 2-bromo-4,5-methylenedioxybenzoic acid (0.86 g, 3.3 mmol) and SOCl₂ (5 ml, 70 mmol) was refluxed for 1 h. After evaporation of the SOCl₂, the residue (5) was dissolved in THF (10 ml). A THF (5 ml) solution of 4 (0.59 g, 3 mmol) was dropwise added to a THF suspension of 60% NaH (0.14 g, 3.6 mmol), and the mixture was stirred at room temperature for 0.5 h. To this mixture was added the THF solution of 5 described above, and the whole mixture was stirred at room temperature for 3 h. The reaction mixture was poured into ice-water and extracted with CHCl₃. The CHCl₃ extract was dried over MgSO₄, and the CHCl₃ was evaporated. The residue was purified by silica gel column chromatography using hexane/CH₂Cl₂ (1:1) to give colorless needles which were recrystallized from acetone. mp 125-126°C. Ir ν (KBr) cm⁻¹: 1714. 300 MHz ¹H-Nmr (CDCl₃) δ (ppm): 6.10 (2H, s), 6.60 (1H, d, J=3.7 Hz), 7.06 (1H, s), 7.11 (1H, d, J=3.7 Hz), 7.14 (1H, s), 7.18 (1H, dd, J=7.7, 7.7 Hz), 7.56 (1H, d, J=7.7 Hz), 7.58 (1H, d, J=7.7 Hz). Ms m/z: 423 (M⁺). Anal. Calcd for C₁₆H₉NO₃Br₂: C, 45.42; H, 2.14; Br, 37.77; N, 3.31. Found: C, 45.33; H, 2.29; Br, 37.70; N, 3.31.

Hippadine (7)

A mixture of 6 (0.50g, 1.46 mmol), (Bu₃Sn)₂ (0.67 g, 1.46 mmol), Bu₄NBr (0.71 g, 2.19 mmol), Li₂CO₃ (0.11 g, 1.46 mmol), Pd(PPh₃)Cl₂ (50 mg, 0.07 mmol), and toluene (20 ml) was refluxed for 12 h. After removal of the solvent, the residue was purified by silica gel column chromatography using CH₂Cl₂ as an eluent to give pale yellow needles, which were recrystallized from hexane-acetone. mp 215-217°C. Lit.,⁸ mp 217-218°C. Ir ν (CHCl₃) cm⁻¹: 1675. 300 MHz ¹H-Nmr (CDCl₃) δ (ppm): 6.17 (2H, s), 6.90 (1H, d, J=3.7 Hz), 7.48 (1H, dd, J=7.7, 7.7 Hz), 7.66 (1H, s), 7.76 (1H, d, J=7.3 Hz), 7.92 (1H, d, J=7.3 Hz), 7.98 (1H, s), 8.05 (1H, d, J=3.7 Hz). Ms m/z: 263 (M⁺). Anal. Calcd for C₁₆H₉NO₃: C, 73.00; H, 3.45; N, 5.32. Found: C, 72.78; H, 3.52; N, 5.29.

REFERENCES

1. T. Sakamoto, C. Satoh, Y. Kondo, and H. Yamanaka, *Heterocycles*, 1992, **34**, 2379.
2. T. Sakamoto, Y. Kondo, A. Yasuhara, and H. Yamanaka, *Heterocycles*, 1990, **31**, 219.
3. T. Sakamoto, Y. Kondo, A. Yasuhara, and H. Yamanaka, *Tetrahedron*, 1991, **47**, 1877.
4. T. Sakamoto, C. Satoh, Y. Kondo, and H. Yamanaka, *Chem. Pharm. Bull.*, 1993, **41**, 81.
5. A. A. Ali, M. K. Mesbah, and A. W. Frahm, *Planta Med.*, 1981, **43**, 407.
6. a) A. M. Echavarren and J. K. Stille, *J. Am. Chem. Soc.*, 1987, **109**, 5478; b) T. R. Kelly, Q. Li, and V. Bhushan, *Tetrahedron Lett.*, 1990, **31**, 161.
7. S. Prabhakar, A.M. Lobo, and M. M. Marques, *J. Chem. Res. (S)*, 1987, 167.
8. K. Hayakawa, T. Yasukouchi, and K. Kanematsu, *Tetrahedron Lett.*, 1987, **28**, 5895.
9. D. St. C. Black, P. A. Keller, and N. Kumar, *Tetrahedron Lett.*, 1989, **30**, 5807.
10. M. A. Siddiqui and V. Snieckus, *Tetrahedron Lett.*, 1990, **31**, 1523.
11. M. Somei Y. Saida, T. Funamoto, and T. Ohta, *Chem. Pharm. Bull.*, 1987, **35**, 3146.

Received, 6th July, 1993