



ALLYLIC SUBSTITUTIONS IN THE 2-BENZALINDAN,
2-BENZYLINDENE SERIES

Georges Maury and Norman H. Cromwell¹

Avery Laboratory, University of Nebraska, Lincoln, Nebraska 68508

(Received in USA 27 February 1969; received in UK for publication 2 April 1969)

In previous papers^{2,3}, it was reported that the allylic substitutions of 2-(α -aminobenzyl)-1-indenones and 3-halogeno-2-benzal-1-indanones with primary and secondary amines occur only with inversion of the allylic system by a bimolecular mechanism. The orientation and the ease of the preceding reactions have been associated with the presence of the electron-withdrawing carbonyl group. Similarly, in the 3-(α -haloalkyl)benzo [b] thiophene 1,1-dioxide series, the sulfonyl group was claimed to increase the reactivity of the allylic system toward amines in reducing the electron density in the C=C bond and in delocalizing the negative charge developed at the β -position in the transition state.⁴ Therefore, a study of the corresponding indene-indan systems 1 and 2 (X=halogen) was planned. It has now been found that the reaction of 2-(α -chlorobenzyl)-indene, 1 (X=Cl), with amines in acetonitrile gives both the direct substitution product and the "abnormal" substitution product.



The reduction of trans 2-benzal-1-indanone with NaBH₄ in aqueous methanol occurred exclusively by 1,2-addition⁵ and gave trans 2-benzal-1-indanol, mp=111-112⁶; in the presence of LiAlH₄ a trans 2-benzyl-1-indanol was formed⁷. The nmr spectrum⁸ of the prepared indanol is in agreement with the proposed structure, showing 9H aromatic between 2.3 and 2.8 τ , 1H vinyl at 3.10 τ (quadruplet, J allyl = 2.3 Hz), 1H benzyl at 4.40 τ (doublet, J = 9Hz), 2H methylene at 6.12 τ and OH at 7.57 τ (doublet, J = 9Hz). The uv spectrum of the alcohol showed absorptions at 258 and 270 (Sh) m μ (ϵ =20,000, 12,500). The structure of the allylic alcohol was further proved by reducing trans 2-deutero-2-benzal-1-indanone under the same conditions; the signal of the vinyl proton is absent in the nmr spectrum of the deuterated alcohol.

Treatment of trans 2 (X=OH) with thionyl chloride in benzene or ether at room temperature yielded only one allyl chloride (mp = 56-58⁰); its nmr spectrum showed a multiplet between 2.5

and 3.0 τ (9H aromatic), a quadruplet at 3.32 τ (1H vinyl, $J_{\text{allyl}} \sim 1.7$ Hz), a singlet at 4.07 τ (1H benzyl), and a singlet at 6.60 τ (2H methylene). The allyl chloride has the structure 1 (X=Cl) as shown by repeating the preceding experiment with trans 2-deuterobenzal-1-indanol; the corresponding H benzyl signal is absent in the nmr spectrum of the deutero-product, the H vinyl signal is a triplet ($J_{\text{allyl}} = 1.7$ Hz) and the methylene signal is a doublet ($J_{\text{allyl}} = 1.7$ Hz). The allyl chloride absorbs in the uv at 267 m μ ($\epsilon=13,700$).

The reaction of five molar equivalents of t-butylamine with 1 (X=Cl) in acetonitrile at 25°C gave very readily two amino allyl isomers isolated by chromatography on florisil: 22% of 1 (X=t-butyl) and 78% of 2 (X=t-butyl). The isomer 1 shows nmr signals between 2.4 and 3.0 τ (9H aromatic), at 3.21 τ (1H vinyl, quadruplet, $J_{\text{allyl}} \sim 1.5$ Hz), at 5.10 τ (1H benzyl), at 6.73 τ (2H methylene), at 8.60 τ (NH), and at 8.91 τ (9H t-butyl); uv absorptions were found at 260 and 265 m μ ($\epsilon=15,800$ and 15,200). The isomer 2 shows nmr signals between 2.3 and 2.8 τ (H aromatic), at 3.14 τ (H vinyl, quadruplet, $J_{\text{allyl}} = 2.6$ Hz), at 5.13 τ (H benzyl), at 6.11 τ (methylene) and at 8.67 τ (NH and t-butyl); the uv absorption is at 251 m μ ($\epsilon=26,300$). The nmr analysis of the reaction product of 2-(α -chlorodeuterobenzyl)-indene with t-butylamine confirms the proposed structures of the two allylamines; all the signals of the two non-deuterated compounds are present except the H vinyl signal of 2-deuterobenzal-1-t-butylamino-indan (the H benzyl and H methylene signals are broad singlets: $J < 1$ Hz) and the H benzyl signal of 2-(α -t-butylamino-deuterobenzyl)-indene (the H vinyl signal is a triplet and the H methylene signal a doublet: $J_{\text{allyl}} = 1.6$ Hz). It must be noted that the allylic coupling constant between H vinyl and H benzyl is expected to be larger in the rigid molecule 2 than in the compound 1 where H benzyl of the lateral chain is probably near the plane of the molecule⁹; this relation is in agreement with the preceding conclusions based on deuteration, and it can be used to assign a structure to the isomers 1 and 2 (X=NH₂) when R is relatively bulky. The nmr absorptions of compounds 1 and 2 are consistent with those of parent compounds⁷. The similarities in the uv and nmr spectra of trans 2-benzal-1-indanol and the allylamines 2 (X=NR₂) suggest that the latter have the structure trans in which the A^(1,3) strain¹⁰ is smaller than in the cis isomer.

When the concentration of 1 (X=Cl) was held constant, it was found that the relative proportions of isomers 1 (X=t-butyl) and 2 (X=t-butyl) at completion of reaction were constant when the number of molar equivalents of amine were increased from 2 to 13 in acetonitrile solvent. The ratio 1/2 also depends on the nature of the reacting amine: it increases in the following sequence of amines: diisopropylamine (15/85), t-butylamine (28/72), isopropylamine (35/65), and piperidine (52/48). The allyl chloride 1 (X=Cl) is stable at room temperature in solvent aceto-

nitrile-d₃; at 60°C, decomposition occurs slowly and without rearrangement. These results suggest that SN₂ and SN₂' reactions occur simultaneously; however an SNi' mechanism (prior to substitution⁴) cannot be ruled out a priori. Kinetic studies are being carried out to check these hypothesis. On the other hand, the reaction of thionyl chloride with trans 2-benzal-1-indanol in ether is believed to be SNi' because of the relatively low reactivity of the chloride ion in displacement reactions under these conditions¹¹, and because of the favorable steric conditions in the transition state.

Attempts are being made to prepare the isomer 2 (X=halogen) and to compare its reactivity toward amines with the reactivities of 1 (X=halogen) and 3-halogeno-2-benzal-1-indanone.

Acknowledgement. This work was supported in part by a grant from the National Cancer Institute of the U. S. Public Health Service (Grant No. 02931) and in part by an award from the Research Council of the University of Nebraska.

References

1. All correspondence concerning this article should be addressed to this author.
2. G. Maury, E. Ming-Wu, N. H. Cromwell, J. Org. Chem., 33, 1900 (1968).
3. G. Maury, E. Ming Wu, N. H. Cromwell, J. Org. Chem., 33, 1907 (1968).
4. F. G. Bordwell, D. A. Schexnayder, J. Org. Chem., 33, 3236 and 3240 (1968).
5. See S. Munavalli, G. Ourisson, Bull. Soc. Chim., 1964, 3103, who have shown that KBH₄ reduces a 2-isopropylidene-1-indanone to a 2-isopropylidene-1-indanol.
6. All new compounds have satisfactory analysis.
7. C. R. Ganellin, J. M. Loynes, H. F. Ridley, R. G. W. Spickett, J. Med. Chem., 10, 827 (1967).
8. The nmr spectra have been run in deuterochloroform, and the uv spectra in n-hexane.
9. N. S. Bhacca and D. H. Williams "Applications of NMR Spectroscopy in Organic Chemistry", Holden-Day Inc. 1964, p. 108.
10. F. Johnson, S. K. Malhotra, J. Am. Chem. Soc., 87, 5492 (1965).
11. R. H. DeWolfe, W. G. Young, Chem. Rev., 56, 814 (1956).