

Dimethyl Δ^3 -1,2-Diazetine-1,2-dicarboxylate: A New Four-membered 6π -Ring System

By E. E. NUNN and R. N. WARRENER*

(*Department of Chemistry, Australian National University, Canberra, A.C.T. 2600, Australia*)

Summary Dimethyl Δ^3 -1,2-diazetine-1,2-dicarboxylate has been prepared by photolysis of the 1:1 adduct of 2,5-dimethyl-3,4-diphenylcyclopenta-2,4-dienone and 2,3-dimethoxycarbonyl-2,3-diazabicyclo[2,2,0]hex-5-ene; this


new ring system formally satisfies the Hückel ($4n + 2$) rule, but no pronounced stability was observed, and the absence of a ring current suggests that it is not aromatic by this criterion.

Δ^3 -1,2-DIAZETINE (and derivatives thereof) is the π -excessive heterocyclic analogue isoelectronic with cyclobutadiene dianion† and formally satisfies the simple Hückel classification of aromaticity. Conflicting reports on its predicted aromaticity have appeared.³ Derivatives of 1,2-dithiete, the sulphur analogue of this potential 6π -system have been prepared,⁴ but their properties may be atypical owing to the contribution from the sulphur d -orbitals.

We were initially interested in the synthesis of the Δ^3 -1,2-diazetene system in view of the successful conversion of the bicyclo[2,2,0]hexene (1) into the cyclobutene (2),⁵ via our 1,2-photoaromatisation reaction.⁶ Similarly, the diaza-analogue (5) has been converted into the diazetene (9) as follows: Reaction of cyclobutadiene [formed by oxidation of its iron tricarbonyl complex (3) with ceric ammonium nitrate] with (4) at 0° formed the diazabicyclohexene (5) (31%; identical with the product from u.v. irradiation of 1,2-dimethoxycarbonyl-1,2-diazacyclohexa-3,5-diene⁷). Cycloaddition of (5) with 2,5-dimethyl-3,4-diphenylcyclopenta-2,4-dienone formed a 1:1 adduct (6), m.p. 177.5–178° (¹H n.m.r. spectrum consistent with formulation). Irradiation‡ of (6) in CDCl₃ at 0° caused elimination of carbon monoxide to form (7) (undetected) which yielded, in the 1,2-photoaromatisation step, the aromatic hydrocarbon (8), together with a relatively unstable substance which was isolated by t.l.c. (95% purity). The ¹H n.m.r. spectrum [δ 6.57 (2H, s, vinylic-H) and 3.88, (6H, OMe) p.p.m.] and mass spectrum [m/e 172 (C₆H₈N₂O₄)] were entirely consistent with the Δ^3 -1,2-diazetene structure (9). Further, hydrogenation (Pd-C) yielded the dihydro-derivative (10)§ [¹H n.m.r. δ 4.34 (4H, s, cyclobutyl-H) and 3.87 (6H, s, OMe) p.p.m.; m/e 174].

The Δ^3 -1,2-diazetene was thermally unstable [$t_{1/2}$ (20°) 6.9 h; $t_{1/2}$ (34°) 1.8 h] even at ambient temperatures and isomerised to the ring-opened product (11) [¹H n.m.r. δ 8.36 (2H, s, imino-aldehydic-H) and 3.98 (6H, s, OMe) p.p.m.]. Rapid polymerisation on attempted isolation frustrated further characterisation of this product.

The question of aromaticity is always a difficult one. Certainly HMO calculations show only a small stabilisation energy for (9), and suggest that it is not aromatic to a significant extent. This is further supported by the ¹H n.m.r. spectrum of (9) (*vide supra*) which indicates no diamagnetic ring current [*cf.* α -vinylic proton resonance in



N-ethoxycarbonylpyrrole (heteroaromatic, δ 7.36 p.p.m.); *N*-ethoxycarbonylazonine (nonaromatic, ^b δ 6.4 p.p.m.)). Thus while the evidence suggests that compound (9) is nonaromatic, we defer a more general comment on this ring system until other members of the series have been prepared.¶

(Received, 5th April 1972; Com. 557.)

† In the 4-membered alicycles, the 2π -system (cyclobutadiene dication) has been formed at -75° and characterised by ¹H n.m.r. spectroscopy,¹ but attempts to form the 6π -equivalent (cyclobutadiene dianion) have been unsuccessful.^{2a} Pettit and his co-workers have now obtained evidence for its intermediacy.^{2b}

‡ American Hanovia (450 W) medium-pressure Hg lamp, 0°, Vycor filter, N₂ atmosphere, 137 mg/0.5 mg in CDCl₃.

§ Contrary to that reported,⁸ we could find no evidence (mass-spectral) for the formation of any 1:1 adduct on attempted thermal cycloaddition of dimethyl azodicarboxylate on ethylene. Indeed ¹H n.m.r. studies on the crude product showed no resonances corresponding to (10).

¶ Anastassiou and his co-workers^{9a} have recently reported that the aromatic properties of the azonine ring system are particularly sensitive to the *N*-substituent.

¹ G. A. Olah, J. M. Bollinger, and A. M. White, *J. Amer. Chem. Soc.*, 1969, **91**, 3667.

^{2(a)} W. Adam, *Tetrahedron Letters*, 1963, 1387; A. T. Blomquist and V. J. Hruby, *J. Amer. Chem. Soc.*, 1964, **86**, 5041. (b) J. S. McKennis, L. Brener, J. R. Schweiger, and R. Pettit, *J.C.S. Chem. Comm.* 1972, 365.

³ M. E. Vol'pin, *Russian Chem. Rev.*, 1960, **29**, 153; A. T. Balaban and Z. Simon, *Rev. Roumaine Chem.*, 1965, **10**, 1059.

⁴ C. G. Krespan, B. C. McKusick, and T. L. Cairns, *J. Amer. Chem. Soc.*, 1960, **82**, 1515.

⁵ E. E. Nunn, Ph.D. Thesis, Australian National University, 1972.

⁶ C. M. Anderson, J. B. Bremner, H. H. Westberg, and R. N. Warrener, *Tetrahedron Letters*, 1969, 1585; E. E. Nunn, W. S. Wilson and R. N. Warrener, *ibid.*, 1972, 175; E. E. Nunn and R. N. Warrener, *Synth. Comm.*, 1972, **2**, 67.

⁷ L. J. Altman, M. F. Semmelhack, R. B. Hornby, and J. C. Vederas, *Chem. Comm.*, 1968, 686. We thank Professor Altman for providing experimental details on this photochemical isomerisation.

⁸ M. A. Englin, A. S. Filatov, and N. F. Sirotenkova, *J. Org. Chem. U.S.S.R.*, 1969, **5**, 1555.

⁹ (a) A. G. Anastassiou, S. W. Eachus, R. P. Cellura, and J. H. Gebrian, *Chem. Comm.*, 1970, 1133. (b) A. G. Anastassiou and J. H. Gebrian, *J. Amer. Chem. Soc.*, 1969, **91**, 4011.