

CLEAVAGE REACTIONS OF THE LEAD-CARBON BOND II. THE REACTION OF TETRAORGANOLEAD COMPOUNDS WITH 1-CHLORO-1,2,3-BENZOTRIAZOLE

BHUVAN C. PANT* AND JAN G. NOLTES

Institute for Organic Chemistry TNO, Utrecht (The Netherlands)


(Received September 13th, 1971)

SUMMARY

1-(Triorganoplumbyl)-1,2,3-benzotriazoles were synthesized in excellent yields by the cleavage of an organic group from tetraorganolead derivatives with 1-chloro-1,2,3-benzotriazole in benzene at 50–60°. The order of ease of cleavage of organic groups by 1-chloro-1,2,3-benzotriazole in the case of mixed-tetraorganolead compounds is discussed.

INTRODUCTION

Nitrogen-bonded heterocyclic lead compounds have been known for a number of years^{1,2,3}. Willemsens and van der Kerk⁴ reported a variety of such compounds obtained in good yield by the reaction of triorganolead hydroxides with five-membered 1,3-diazaheterocycles, *e.g.*:




The observation of a facile and specific cleavage of lead-carbon bonds in tetraorganolead compounds by *N*-bromosuccinimide⁵ has led us to explore reactions of other compounds containing nitrogen-bonded "positive" halogen with organolead derivatives as a possible source of organolead compounds containing lead-nitrogen bonds. Such a study, moreover, might contribute to our knowledge of the reactivity of the lead-carbon bond. In the present paper we report the synthesis of 1-(triorganoplumbyl)-1,2,3-benzotriazoles by cleavage of an organic group from tetraorganolead compounds with 1-chloro-1,2,3-benzotriazole.

* Present address: School of Molecular Sciences, University of Warwick, Coventry, Warwickshire CV47AL, England.

TABLE I


REACTION OF ORGANOLEAD COMPOUNDS WITH 1-CHLORO-1,2,3-BENZOTRIAZOLE

Organolead compound used	Organic chloride formed	1-(Triorgano-plumbyl)-1,2,3-benzotriazole obtained	Yield (%)	M.p. ^a (°C)	Analytical data ^b , found (calcd.) (%)			
					C	H	N	Pb
								
(C ₆ H ₅) ₄ Pb	C ₆ H ₅ Cl	(C ₆ H ₅) ₃ PbX	88	280	51.95 (51.79)	3.45 (3.42)	7.57 (7.55)	37.22 (37.23)
(p-CH ₃ C ₆ H ₄) ₄ Pb	p-CH ₃ C ₆ H ₄ Cl	(p-CH ₃ C ₆ H ₄) ₃ PbX	93	235-240	54.09 (54.18)	4.53 (4.18)	7.09 (7.02)	
(2-C ₄ H ₉ S) ₄ Pb	2-C ₄ H ₉ SCl	(2-C ₄ H ₉ S) ₃ PbX	80	225	37.91 (37.63)	2.14 (2.26)	7.25 (7.32)	36.24 (36.06)
(C ₆ H ₅) ₃ PbCH ₂ C ₆ H ₅	C ₆ H ₅ Cl	(C ₆ H ₅) ₂ PbX CH ₂ C ₆ H ₅	80	218	52.74 (52.63)	3.78 (3.68)	7.42 (7.37)	36.97 (36.32)
(2-C ₄ H ₉ S) ₃ PbCH ₂ C ₆ H ₅	2-C ₄ H ₉ SCl	(2-C ₄ H ₉ S) ₂ PbX CH ₂ C ₆ H ₅	97	150-160	42.81 (43.30)	3.05 (2.89)	7.31 (7.22)	
(C ₆ H ₅) ₃ Pb-i-C ₃ H ₇	C ₆ H ₅ Cl	(C ₆ H ₅) ₂ PbX CH ₂ C ₆ H ₅	96	195-200	48.29 (48.28)	4.09 (4.02)	8.13 (8.05)	39.53 (39.65)
(C ₆ H ₅) ₃ PbCH ₂ CH=CH ₂	ClCH ₂ CH=CH ₂	(C ₆ H ₅) ₃ PbX i-C ₃ H ₇	88	280	51.86 (51.79)	3.52 (3.42)	7.58 (7.55)	
(C ₆ H ₅) ₃ PbC≡CC(CH ₃) ₃	ClC≡CC(CH ₃) ₃	(C ₆ H ₅) ₃ PbX	90	280	51.82 (51.79)	3.54 (3.42)	7.59 (7.55)	
(CH ₃) ₃ PbC ₆ H ₅	C ₆ H ₅ Cl	(CH ₃) ₃ PbX	89	175	29.10 (29.19)	3.71 (3.51)	11.40 (11.35)	56.09 (55.94)
(C ₆ H ₅) ₃ Pb(C ₆ H ₄ CH ₃ -p)	C ₆ H ₅ Cl + p-CH ₃ C ₆ H ₄ Cl	Mixture						

^a Compounds turned brown or black at this temperature. ^b Analyses were carried out in this Institute by Mr. W. J. Buis.

RESULTS AND DISCUSSION


1-Chloro-1,2,3-benzotriazole⁶ is a "positive-halogen" compound of the same general type as *N*-bromosuccinimide, and thus it was thought worthwhile to study cleavage reactions of tetraorganolead compounds using this reagent. It was found that 1-chloro-1,2,3-benzotriazole readily reacts with tetraorganolead compounds in benzene solution at 50–60° according to:



The various 1-(triorganoplumbyl)-1,2,3-benzotriazoles obtained in this manner are listed in Table 1.

This type of reaction offers a convenient synthesis of 1-(triorganoplumbyl)-1,2,3-benzotriazoles in excellent yields. No formation of side products is observed. The 1-chloro-1,2,3-benzotriazole starting material which is readily obtained by the reaction of 1,2,3-benzotriazole with sodium hypochlorite in acetic acid⁶ is a stable material readily soluble in benzene.

No reaction was found to occur between tetramethyllead and 1-chloro-1,2,3-benzotriazole even in refluxing benzene. Similarly no dephenylation occurred in the case of tetraphenyltin, although the phenyl group was cleaved in trimethylphenyltin yielding 1-(trimethylstannyl)-1,2,3-benzotriazole in quantitative yield:



The relative ease of cleavage of different organic groups attached to the lead atom in mixed-tetraorganolead derivatives with 1-chloro-1,2,3-benzotriazole was also studied. It was found that:

- (i). In the series $(C_6H_5)_3PbR$ where $R = CH_3$; $i-C_3H_7$; $C_6H_5CH_2$, the phenyl group was cleaved exclusively.
- (ii). In the compounds of the type $(C_6H_5)_3PbR$ where $R = -CH_2CH=CH_2$; $-C\equiv C-C(CH_3)_3$, no phenyl group cleavage was observed and the group R was found to be exclusively cleaved.
- (iii). In the case of $(C_6H_5)_3Pb(C_6H_4CH_3-p)$ both the phenyl and *p*-tolyl groups were simultaneously cleaved with approximately the same rate (found by GLC technique).

The results are analogous to those obtained in the cleavage reactions of mixed-tetraorganolead compounds with *N*-bromosuccinimide⁵.

The reactions of various other types of organolead compounds in particular those containing lead-hetero atom bonds with 1-chloro-1,2,3-benzotriazole are being investigated.

EXPERIMENTAL

1-Chloro-1,2,3-benzotriazole, prepared as previously described⁶, was recrystallized from 1/1 methylene chloride/light petroleum 40–60° (yield 77%; m.p. 106°, lit.⁶ 105–106°).

As identical procedures were used for the preparation of 1-(triorganoplumbyl)-1,2,3-benzotriazoles only a few typical examples are described (cf. Table 1).

Reaction between tetraphenyllead and 1-chloro-1,2,3-benzotriazole

To a suspension of tetraphenyllead (5.1 g) in benzene (50 ml) at 50° was added a benzene solution (25 ml) of 1-chloro-1,2,3-benzotriazole (1.5 g) over a period of 15 min. The mixture was stirred at 50–60° for a period of 4 h. The white precipitate was filtered, washed with benzene, and dried. The solid obtained was dissolved in warm dimethylformamide. Pure 1-(triphenylplumbyl)-1,2,3-benzotriazole (4.9 g) was precipitated from this solution with 96% ethanol, filtered and dried at 50°/0.1 mm for 2 h.

Reaction between trimethylphenyllead and 1-chloro-1,2,3-benzotriazole

A solution of 1-chloro-1,2,3-benzotriazole (0.8 g) in benzene (25 ml) was added to a well-stirred solution of trimethylphenyllead (1.6 g) in benzene (25 ml) at 50–60°. The mixture was stirred at this temperature for 4 h. Removal of benzene and recrystallization of the crude material from boiling acetone containing a small amount of dimethylformamide gave pure 1-(trimethylplumbyl)-1,2,3-benzotriazole (1.6 g).

Reaction between trimethylphenyltin and 1-chloro-1,2,3-benzotriazole

A benzene solution (25 ml) of trimethylphenyltin (1.5 g) was allowed to react with a solution of 1-chloro-1,2,3-benzotriazole (1.0 g) in benzene (25 ml) at 60° for 3 h. Evaporation of benzene and recrystallization of the residue from boiling acetone containing a small amount of dimethylformamide gave pure 1-(trimethylstannyl)-1,2,3-benzotriazole (1.6 g; 94% yield). M.p. 224–226° (lit. value⁷: 221.5–223°).

ACKNOWLEDGEMENT

The authors wish to thank the International Lead Zinc Research Organization (Dr. S. F. Radtke), New York, U.S.A. for financial support, and Dr. L. C. Willemsens for discussions and for the supply of organolead starting materials.

REFERENCES

- 1 R. HEAP AND B. C. SAUNDERS, *J. Chem. Soc.*, (1949) 2983.
- 2 B. C. SAUNDERS, *J. Chem. Soc.*, (1950) 684.
- 3 W. B. LIGETT, R. D. CLOSSON AND C. N. WOLF, *U. S. Pat.* 2,654,689; Oct. 6, (1953); *Chem. Abstr.*, 48 (1954) 942.
- 4 L. C. WILLEMSSENS AND G. J. M. VAN DER KERK, *Investigations in the field of organolead chemistry*, Inst. Org. Chem. TNO, Utrecht, The Netherlands; ILZRO, New York, 1965, p. 70.
- 5 B. C. PANT, *J. Organometal. Chem.*, 24 (1970) 697.
- 6 C. W. REES AND R. C. STORR, *Chem. Commun.*, (1968) 1305.
- 7 J. G. A. LUITEN AND G. J. M. VAN DER KERK, *Recl. Trav. Chim. Pays-Bas*, 82 (1963) 1181.

J. Organometal. Chem., 36 (1972)