


Conversion of Ketones into Lactones with Benzeneseleninic Acid and Hydrogen Peroxide (Benzeneperoxyseleninic Acid): a New Reagent for the Baeyer-Villiger Reaction

By PAUL A. GRIECO,* YUUSAKU YOKOYAMA, SYDNEY GILMAN, and YASUFUMI OHFUNE
(Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260)

Summary Benzeneperoxyseleninic acid generated *in situ* from benzeneseleninic acid and hydrogen peroxide constitutes a new reagent for the Baeyer-Villiger reaction.




THE ability of benzeneperoxyseleninic acid to epoxidize olefins has recently been reported.¹ We report herein that benzeneperoxyseleninic acid (1) constitutes a new reagent for use in the Baeyer-Villiger reaction. The reagent is generated *in situ* from the readily available benzeneseleninic acid² and an excess of 30% hydrogen peroxide [equation (1)]. The reaction can be carried out under either homo-

After 5 min, 1.0 mmol of 2-methylcyclohexanone in 1.0 ml of CH_2Cl_2 was added to the vigorously stirred heterogeneous reaction mixture. The reaction was quenched after 1 h with saturated NaHCO_3 solution giving, after purification, pure lactone in 83% yield.



geneous [tetrahydrofuran (THF)] or heterogeneous (CH_2Cl_2) conditions. In general, 1.25 mol. equiv. of PhSe(O)OH and 10 mol. equiv. of 30% H_2O_2 are employed. In order to suppress hydrolysis to the corresponding hydroxycarboxylic acid, the reaction can be buffered at pH 7 with phosphate buffer.

The following experimental procedure illustrates the method. A suspension of 1.25 mmol of PhSe(O)OH in 1.5 ml of CH_2Cl_2 and 1.0 ml of water was treated at room temperature with 10 mmol of *ca.* 30% aqueous H_2O_2 .

















The potential of this new reaction was made evident from the Baeyer-Villiger oxidation of the ketone (2) which could not be oxidized to either the lactone (3) or the hydroxy acid (4a) employing a variety of procedures³ (40% peroxyacetic acid, *m*-chloroperoxybenzoic acid, anhydrous peroxyacetic acid, and selenium dioxide-hydrogen peroxide). Initial attempts at oxidation of (2) with (1) [1.25 mol. equiv. PhSe(O)OH, 10 mol. equiv. 30% H₂O₂] in THF gave, after 4 h, a *ca.* 15% yield of (3) with 50% recovery of the starting

ketone. However, heating the above reaction mixture for 13 h at 45 °C gave, after work-up and esterification, a 63% isolated yield of the pure hydroxy ester (4b).†

Cyclopentanones can be slowly converted into δ -lactones employing the conditions given in the Table. For example, estrone methyl ether gave the corresponding δ -lactone in 80% yield after 24 h. The synthesis of natural psilostachyin C from damsine was carried out [equation (2)] employing (1) [2.0 mol. equiv. PhSe(O)OH, 15 mol. equiv. 30% H₂O₂] in Bu^tOH. Natural psilostachyin C, m.p. 224–225 °C (lit.⁴ m.p. 224–226 °C), was obtained in 55% isolated yield. *Ca.* 20% of the corresponding hydroxycarboxylic acid, which resulted from hydrolysis of the δ -lactone during the course of the reaction or the work-up, was isolated.

TABLE. Reaction of ketones with benzeneperoxyseleninic acid.^a

Ketone	Product	Solvent	Time/h	% Yield ^b
		THF CH ₂ Cl ₂	0.75 1	60 ^c 83
		CH ₂ Cl ₂	2	83
		CH ₂ Cl ₂	18	67
		CH ₂ Cl ₂	2.5	79 ^d
		CH ₂ Cl ₂	2	50 ^e
		THF	24	80
		CH ₂ Cl ₂	96	60 ^f

^a All reactions were carried out in either THF or CH₂Cl₂, as indicated, using 1.25 mol equiv. of PhSe(O)OH and 10.0 mol. equiv. of 30% aqueous H₂O₂. ^b All compounds were fully characterized by spectral methods. Yields reported are for isolated, chromatographically pure substances. ^c *Ca.* 8% of the corresponding hydroxy acid isolated as its methyl ester was found. ^d This reaction was carried out in the presence of pH 7 phosphate buffer. ^e *Ca.* 45% of the starting ketone was recovered. ^f Starting material was recovered in 35% yield.


† This reaction was carried out by Dr. M. Uemura.

¹ P. A. Grieco, Y. Yokoyama, S. Gilman, and M. Nishizawa, *J. Org. Chem.*, 1977, **42**, 2034.


² J. D. McCullough and E. S. Gould, *J. Amer. Chem. Soc.*, 1949, **71**, 674.

³ P. McCurry, personal communication.

⁴ H. B. Kagan, H. E. Miller, W. Renold, M. V. Lakshimkantham, L. R. Tether, W. Herz, and T. H. Mabry, *J. Org. Chem.*, 1966, **31**, 1629.



During an attempt to reduce the amount of H₂O₂ employed in the examples cited in the Table, we observed a novel reaction with the ketone (5). Upon treatment of (5) in THF with 1.25 mol. equiv. of PhSe(O)OH and 1.50 mol. equiv. of 30% H₂O₂, we isolated, after 16 h, a 60% yield of the pure enone (6) and a 12% yield of the keto selenide (7). There was no lactone formation under the above conditions.



We thank the National Institutes of Health, Glidden Organics, and Shell Development Company for support of this research, Professor E. Rodriguez for generous samples of natural damsine and natural psilostachyin C, and Professor P. McCurry, Carnegie-Mellon University, for a sample of the ketone (2).

(Received, 6th July 1977; Com. 684.)