

Fluorination of Diazo-ketones with Fluoroxytrifluoromethane

By CLAUDE WAKSELMAN* and JACQUES LEROY


(C.N.R.S.-C.E.R.C.O.A., 2 rue Henri Dunant, Thiais 94320, France)

Summary Fluorination of diazo-ketones with CF_3OF produces a mixture of $\alpha\alpha$ -difluoro-ketones and α -trifluoromethoxy- α -fluoro-ketones by the initial electrophilic attack on the diazo-group followed by nucleophilic attack with F^- or CF_3O^- .

It is possible to consider fluoroxy-compounds as 'pseudohalogen' derivatives of fluorine.¹ Their reactions with double bonds exhibit characteristics which are typical of electrophilic additions.² The α -fluorocarbocation initially formed is captured either by CF_3O^- or F^- ($\text{CF}_3\text{O}^- \rightleftharpoons \text{COF}_2 + \text{F}^-$).

Halogens or pseudohalogens are known to react readily with aliphatic diazo-compounds.³ The mechanism is considered to involve initial electrophilic attack on the diazo-alkane. In order to confirm the 'pseudohalogen' nature of CF_3OF , we examined its reaction with some α -diazo-ketones. We obtained mixtures containing mainly $\alpha\alpha$ -difluoro-ketones⁴ and α -trifluoromethoxy- α -fluoro-ketones, hitherto unknown compounds. This result can be explained by nucleophilic attack of the diazo-alkane upon the fluorine, followed by nucleophilic attack upon intermediate (2) (or its α -keto- α -fluorocarbonium ion), either by F^- or CF_3O^- .

For instance, α -diazoacetophenone, prepared from benzoyl chloride (0.02 mol) was treated in CFCl_3 with CF_3OF



a; R = Ph (3) : (4) = 58 : 42

b; R = PhCH_2 (3) : (4) = 62 : 38

c; R = cyclohexyl (3) : (4) = 66 : 34

(1.5 mol. equiv.) at -75°C . After work-up (aqueous NaHCO_3), ca. 2.5 g of crude oil was obtained. Compounds (3a) and (4a) form 85% of the crude mixture, in the ratio of 58 : 42, as shown by ^{19}F n.m.r. spectroscopy. The overall yields, from benzoyl chloride, of pure analytical samples (3a) and (4a) were 14 and 10%, respectively. I.r., n.m.r. and mass spectra are consistent with the structures given.

(Received, 6th April 1976; Com. 371.)

¹ D. H. R. Barton, L. S. Godinho, R. H. Hesse, and M. M. Pechet, *Chem. Comm.*, 1968, 804.

² D. H. R. Barton, L. J. Danks, A. K. Ganguly, R. H. Hesse, G. Tarzia, and M. M. Pechet, *J.C.S. Perkin I*, 1976, 101.

³ G. A. Olah and J. Welch, *Synthesis*, 1974, 896.

⁴ For other preparations of α -difluoro-ketones see: S. Nakanishi, R. L. Morgan, and E. V. Jensen, *Chem. and Ind.*, 1960, 1136; J. Cantacuzène and J. Leroy, *Tetrahedron Letters*, 1970, 3277; P. Crabbé, A. Cervantes, A. Cruz, E. Galeazzi, J. Iriarte, and E. Velarde, *J. Amer. Chem. Soc.*, 1973, **95**, 6655; T. B. Patrick and E. C. Hayward, *J. Org. Chem.*, 1974, **39**, 2120; C. Wakselman and M. Tordeux, *J.C.S. Chem. Comm.*, 1975, 956.