



tions using a large excess of cesium fluoride.


tert-Butyl alcohol does not react with **1** even at 80 °C. Thus it makes an excellent solvent for solid alcohols. When 2,2-dimethyl-1,3-propanediol (**4**, 3.65 g) is dissolved in *t*-BuOH (20 ml) containing 10 equiv of CsF and 1 g of **2**, a quantitative yield of **5** is obtained after 2 days.



Another feature of the general reaction is the formation of cyclic phosphates from diols. For example, if one starts with tris(trichloroethyl) phosphate (**6**) and compound **4** (1 g of **6** and 4.23 g of **4**) in *t*-BuOH (20 ml) along with 10 equiv of CsF at room temperature, the cyclic phosphate (**7**) is obtained in 95% yield. The remaining trichloroethyl group of **7** can be completely exchanged to give **8** by heating **7** at 80 °C in other alcohols (1 g of **7**/20 ml) with cesium fluoride (5 equiv) for 2 days.



As a final example, the exchange reaction occurs readily in nucleotide triesters where, for example, compound **9** is converted into **10** in 95% yield (100 mg of **8**/10 ml of MeOH and 30 equiv of CsF at room temperature for 2 days).



Thus the reaction described⁹ in this report is remarkably versatile and will benefit fields ranging from pesticides to phospholipids to nucleotides.

Acknowledgment. We wish to thank the National Research Council of Canada and the Province of Quebec for financial support.

References and Notes

- (1) R. L. Letsinger and W. B. Lunsford, *J. Am. Chem. Soc.*, **98**, 3655–3661 (1976).
- (2) F. Ramirez, J. F. Marecek, and I. Ugl, *J. Am. Chem. Soc.*, **97**, 3809–3817 (1975).
- (3) M. S. Poonian, E. F. Nowosiat, and A. L. Nussbaum, *J. Am. Chem. Soc.*, **94**, 3992–3996 (1972).
- (4) Prepared by reaction of the bis(2,2,2-trichloroethyl) phosphorochloridate with the alcohol in pyridine.
- (5) Reactions can be conveniently monitored by gas chromatography. The instrument used was a Hewlett-Packard Model 5711A gas chromatograph equipped with a 10 ft X 1/8 in. o.d. stainless steel column packed with 10% OV-1 on chromasorb W-HP. The off-column injector had a glass liner which must be changed frequently to prevent buildup of cesium fluoride resulting in incorrect analyses.
- (6) The presence of water in the alcohol slows the reactions to some extent. However, to make the synthesis as practical as possible the results reported here are based on laboratory grade absolute alcohol and spectrograde methanol (American Chemicals Ltd.).
- (7) All new products have been fully characterized including mass spectrometry on an AEI 902 high resolution instrument.
- (8) It is possible to exchange one group in **6** with other alcohols as well. For example, **1** (R = Et) is obtained in 77% yield after 4 days at room temperature when **6** is dissolved in ethanol (1 g/200 ml) along with 1 equiv of CsF. After this time 15% of **6** was unchanged and 8% of **2** (R = Et) was obtained.
- (9) The reactions apparently occur via initial attack of fluoride ion on phosphorus followed by rapid reaction of the phosphorofluoridate with the alcohol. The intermediate fluoride from **7** can be detected in *t*-BuOH. This aspect will be dealt with later in a full report.

Kelvin K. Ogilvie,* Serge L. Beaucage
Nicole Theriault, Douglas W. Entwistle

*Department of Chemistry, McGill University
Montreal, Quebec, Canada H3C 3G1
Received September 16, 1976*


Conversion of Aminoglycosidic Antibiotics: Novel and Efficient Approaches to 3'-Deoxyaminoglycosides via 3'-Phosphoryl Esters

Sir:


Semisynthetic 3'-deoxyaminoglycosidic antibiotics, including 3',4'-dideoxy derivatives, are remarkably effective against resistant strain bacteria producing phosphotransferases.^{1,2}

We now report new, simple methods for the selective dehydroxylation of aminoglycosides by a combination of enzymatic and chemical reactions; the former is phosphorylation of aminoglycosides by using enzymes from resistant strains,^{1,3} while the latter involves transformation of phosphates into 3'-deoxyaminoglycosides by treatment with silylating agents and subsequent hydrogenation. The procedures present a conceptionally new and promising approach to modifications of polyfunctional antibiotics. Typical experimental procedures for the transformations are available; see paragraph at end of paper regarding supplementary material.


Kanamycin B (**1**) was phosphorylated with the enzyme from *Pseudomonas aeruginosa* GN 573⁴ in the presence of ATP and MgSO₄ to its 3'-phosphate (**2**), mp 220–230 °C dec, [α]_D +106° (c 0.5, H₂O), in 99% yield.⁵ Reaction of **2** with trimethylchlorosilane (TMCS)–hexamethyldisilazane (9:4 by volume) in a mixture of pyridine and HMPA in the presence of triphenylphosphine⁶ in a sealed tube (120 °C, 30 h) yielded after hydrolysis, 3'-chloro-3'-deoxykanamycin B (**3**), mp 190–195 °C dec (from C₂H₅OH), [α]_D +126° (c 1.0, H₂O). The chloride **3** was hydrogenated with Raney nickel in the presence of triethylamine in water to afford 3'-deoxykanamycin B (**4**), identical with a natural product (tobramycin), in an overall yield of 47% based on **2**.^{2,7} Similarly, 3'-deoxyneamine (**6**, nebramine),^{2,7} 3'-deoxyxylostasin (**9**),⁸ mp 134–135 °C dec (from CH₃OH), [α]_D +28° (c 0.5, H₂O), 3'-deoxyribostamycin,⁹ 3'-deoxyparomycin I (lividomycin B),¹⁰ and 3'-deoxyneomycin B² were obtained from the corresponding aminoglycosides. The chlorination of butirosin A



- 1, X = OH; Y¹ = OH; Y² = 3AGlu; R = H
 2, X = OPO₃H₂; Y¹ = OH; Y² = 3AGlu; R = H
 3, X = Cl; Y¹ = OH; Y² = 3AGlu; R = H
 4, X = H; Y¹ = OH; Y² = 3AGlu; R = H
 5, X = Cl; Y¹ = OH; Y² = OH; R = H
 6, X = H; Y¹ = OH; Y² = OH; R = H
 7, X = Cl; Y¹ = Xyl; Y² = OH; R = H
 9, X = H; Y¹ = Xyl; Y² = OH; R = H
 10, X = OH; Y¹ = Xyl; Y² = OH; R = AHBA
 11, X = OPO₃H₂; Y¹ = Xyl; Y² = OH; R = AHBA
 13, X = H; Y¹ = Xyl; Y² = OH; R = AHBA



8, R = H
 12, R = AHBA



3'-phosphate (**11**)¹¹ proceeded in lower yield, presumably owing to an undesired side reaction on the acyl moiety. Kanamycin A 3'-phosphate¹ possessing a hydroxy group at the 2'-position was recovered unchanged in the chlorination reaction. The equatorial configuration of the chlorine atom at the 3'-position was established on the basis of the proton-proton coupling constants observed in **3** and 3'-chloro-3'-deoxyneamine (**5**) ($J_{2',3'} = 10$ and 11 Hz, respectively).

Another dehydroxylation procedure is the one involving aziridine derivatives. Treatment of **11** with bistrimethylsilylacetamide (BSA)-TMCS (5:1 by volume) in pyridine in a sealed tube (105 °C, 30 h) afforded, after hydrolysis, 2',3'-epimino-2'-deamino-3'-deoxybutirosin A (**12**), mp 212–214 °C (from CH₃OH), $[\alpha]_D +36^\circ$ (c 0.5, H₂O), in 58% yield with 80% conversion of **11**. The assigned structure for **12** was confirmed by the comparison of ¹³C NMR spectra of butirosin A (**10**)¹² and **12**. The signal of C-2' and C-3' forming the aziridine ring in **12** appeared at the higher field¹³ (chemical shifts for C-2' and C-3': 33.7–34.9 ppm in **12**; 56.4 and 74.0 ppm in **10**).¹⁴ Hydrogenation of **12** with Raney nickel in water at 70 °C, followed by separation by ligand exchange chromatography¹⁵ (Amberlite CG-50, Cu–NH₃ form) gave 3'-deoxybutirosin A (**13**), mp 204–208 °C (from CH₃OH), $[\alpha]_D +22^\circ$ (c 0.5, H₂O), in 57% yield in preference to the isomer (product ratio, 5:1). The structure of **13** was confirmed by hydrolysis leading to **6** and **9**. This method was successfully applied to 3'-phosphates of **1**, neamine,¹ xylostasin,¹⁶ and butirosin B¹⁷ to obtain **4**, **6**, **9**, and 3'-deoxybutirosin B,¹⁸ respectively, via the corresponding 2',3'-epiminoaminoglycosides.

3'-Chloro- and 2',3'-epiminoaminoglycosides thus obtained are interconvertible. For example, 2',3'-epimino-2'-deamino-3'-deoxyxylostasin (**8**) was converted into 3'-chloro-3'-deoxyxylostasin (**7**) in high yield under the chlorination condition, while treatment of **7** with BSA in pyridine in a sealed tube (120 °C, 25 h) gave **8** in moderate yield.¹⁹

These results, together with the aforementioned observation obtained in both the substitutions (2 → **3**, **11** → **12**), indicate that silylated epiminoglycosides are intermediates in the chlorination reaction.

Acknowledgment. The authors would like to thank Drs. E. Ohmura, H. Morimoto, T. Miki, M. Yoneda, and S. Igarasi for their advice and encouragement. Thanks are also due to Mr. H. Mabuchi and Mr. M. Yamamoto for their technical assistance.

Supplementary Material Available: Typical experimental procedures for the transformations (4 pages). Ordering information is given on any current masthead page.

References and Notes

- (1) H. Umezawa, "Advances in Carbohydrate Chemistry and Biochemistry", Vol. 30, R. S. Tipson and H. Horton, Ed., Academic Press, New York, N.Y., 1974, pp 183–225.
- (2) S. Umezawa, ref 1, pp 165–173.
- (3) O. K. Sebek, *Acta Microbiol. Acad. Sci. Hung.*, **22**, 381 (1975).
- (4) M. Kida, T. Asako, M. Yoneda, and S. Mitsuhashi, "Microbial Drug Resistance", S. Mitsuhashi and H. Hashimoto, Ed., University of Tokyo Press, Tokyo, 1975, pp 441–448.
- (5) All compounds described herein were separated by column chromatography on ion exchange resin Amberlite CG-50 (NH₄⁺ form), and characterized by microanalysis and thin layer chromatography, some compounds were also confirmed by mass spectra of their N-acetyl-O-trimethylsilyl derivatives, ¹H NMR, ¹³C NMR, and antibacterial spectra.
- (6) The presence of triphenylphosphine increased the yield of the desired products. It might have played the role of antioxidant.
- (7) K. F. Koch and J. A. Rhoades, *Antimicrob. Agents Chemother.*, **309** (1970).
- (8) For the preparation of xylostasin see, S. Horii, I. Nogami, N. Mizokami, Y. Arai, and M. Yoneda, *Antimicrob. Agents Chemother.*, **5**, 578 (1973).
- (9) D. Ikeda, T. Tsuchiya, S. Umezawa, and H. Umezawa, *J. Antibiot.*, **26**, 799 (1973).
- (10) T. Mori, Y. Kyotani, I. Watanabe, and T. Oda, *J. Antibiot.*, **25**, 149 (1972).
- (11) M. Yagisawa, H. Yamamoto, H. Naganawa, S. Kondo, T. Takeuchi, and H. Umezawa, *J. Antibiot.*, **25**, 748 (1972); K. Nakahama, H. Shirafuji, I. Nogami, M. Kida, and M. Yoneda, Annual Meeting of the Agricultural Chemical Society of Japan, 1976, Abstract Paper 262.
- (12) P. W. K. Woo and R. D. Westland, *Carbohydr. Res.*, **31**, 27 (1973).
- (13) G. E. Maciel and G. B. Savitsky, *J. Phys. Chem.*, **69**, 3925 (1965).
- (14) The δ values are from spectra of D₂O solutions containing dioxane as an internal standard. They were recorded in parts per million downfield from Me₄Si (dioxane, 67.4 ppm). Four peaks due to C-2', C-3', C-2, and C- β (N¹-acyl part) appeared at 33.7, 34.0, 34.7, and 34.9 ppm, but the exact assignment has not yet been established.
- (15) J. A. Navratil, E. Murgia, and H. F. Walton, *Anal. Chem.*, **47**, 122 (1975).
- (16) S. Igarasi, M. Kida, U.S. Patent 3 933 790, *Chem. Abstr.*, **82**, 140442 (1975).
- (17) Butirosin B 3'-phosphate was prepared in the same manner as **2** using the enzyme from *Escherichia coli* R 11.
- (18) D. Ikeda, F. Nagaki, S. Umezawa, T. Tsuchiya, and H. Umezawa, *J. Antibiot.*, **28**, 616 (1975).
- (19) Compound **7**, mp 180–185 °C dec (from C₂H₅OH-acetone), $[\alpha]_D +30^\circ$ (c 0.6, H₂O); compound **8**, mp 146 °C (from CH₃OH), $[\alpha]_D +44^\circ$ (c 1.1, H₂O).

Tetsuya Okutani, Tsunehiko Asako, Kouichi Yoshioka
 Kentaro Hiraga,* Makoto Kida

Central Research Division, Takeda Chemical Industries, Ltd.
 Yodogawa-ku, Osaka 532, Japan

Received August 23, 1976

Intrinsic Basicity Determination Using Metastable Ions

Sir:

We report a new method for the determination of intrinsic relative proton affinities. The procedure is sensitive to small differences in base strength and is simple in concept and in practice. It appears to be capable of generalization to the determination of affinities toward other ions. Present mass