J.C.S. Снем. Сомм., 1972

The Reactions of Copper(1) Phenylacetylide with Nitrones

By MANABU KINUGASA* and SHIZUNOBU HASHIMOTO

(Department of Applied Chemistry, Doshisha University, Karasuma-imadegawa, Kamikyo-ku, Kyoto, Japan)

Summary Reactions of copper(1) phenylacetylide with nitrones give cis- β -lactams.

WE have investigated the reactions of copper acetylides with 1,3-dipoles,¹ and have found that with nitrones β -lactams are formed.

The reactions of copper(I) phenylacetylide (1) with nitrones (2a—d) were performed in dry pyridine under a nitrogen atmosphere. After hydrolysis, β -lactams (3a—d) were obtained in good yield.

The structures of (3a-d) were confirmed by n.m.r. and i.r. spectra. Yields and physical data of (3a-d) are given in the Table.

The configurations of (3a) and (3c) were shown to be *cis* from the following information. (3a) had m.p. and i.r., n.m.r., and mass spectra in agreement with those of a *cis*- β -lactam described in the literature.² It has been reported that the β -lactam produced from the reaction of N-(o-methylbenzylidene)aniline with phenylketen, the isomer of (3c), has a m.p. of 138° ,³ and its configuration was shown



to be *trans* since the addition of a keten to an imine always gave a *trans*- β -lactam.⁴ Therefore, (3c), with m.p. 213—

214°, was deduced to exist in the *cis* form. It is possible that (3b) and (3d) formed in the same reaction, *i.e.*, "the

Yields and physical data of cis- β -lactams (3a-d)

Lactam ^a	Yield (%)	M.p. (°C)	$\nu(C=O)$ (Nujol)	HA	δ (CDCl _s) H _B	<i>J</i> ав (Нz)
(3a)	54.5	186 (182—183) ^b	1750 (5·73 μm) ^b	4·95 (4·96) ^b	5·43 (5·44) ^b	6·5 (7·0) ^b
(3b)	60.2	156	1750	`4·30 ′	`4 •95 [′]	3.0
(3c)	50.6	213 - 214	1740	4.20	5.25	$3 \cdot 0$
(3 ď)	51.2	190—192	1745	5.00	5.75	6.3

* Satisfactory analytical data were obtained on all β -lactams. b Values from ref. 2 are in parenthesis.

acetylide reaction," prefer the cis form. Also, the trans isomers of (3b) and (3d) have not been prepared by "the keten reaction."3

Although β -lactams have been synthesized in various ways,⁵ no reaction giving only a $cis-\beta$ -lactam has yet been reported. Hence, "the acetylide reaction" is useful as a stereoselective reaction and for the synthesis of $cis-\beta$ lactams.

(Received, 24th February 1972; Com. 303.)

¹ S. Hashimoto, W. Koike, and M. Kinugasa, Abstracts 24th Annual Meeting of the Chemical Society, Japan, No. 3, 1971, p. 1721.
² O. L. Chapman and W. R. Adams, J. Amer. Chem. Soc., 1968, 90, 2333.
⁸ R. Pfleger and A. Jöger, Chem. Ber., 1957, 90, 2460.
⁴ A. K. Bose, G. Spiegelmann, and M. S. Manhas, Tetrahedron Letters, 1971, 3167.
⁵ L. L. Muller and J. Hamer, '1,2-Cycloaddition Reactions,' Wiley, New York, 1967, p. 173, and references therein.