


Table 1. Relative Abundances of Acids Formed on Hydration of Ketenes with 95% H_2^{18}O in THF

acid	ketene			
	1 ^a	4 ^a	5 ^a	6 ^b
$\text{Ar}_2\text{CHC}^{16}\text{O}_2\text{H}$	(9.3) ^c	(22.4) ^c	(16.0) ^c	(77.8) ^c
$\text{Ar}_2\text{CHC}^{18}\text{O}^{16}\text{OH}$	100.0	100.0	100.0	100.0
$\text{Ar}_2\text{CHC}^{18}\text{O}_2\text{H}$	60.0	33.5	25.0	21.8

^a From the $\text{MH}^+ - \text{Ar}$ signals. ^b From the MH^+ signals. ^c See text.

in the hydration of the labeled ketene with H_2^{18}O . Consequently, the acid isolated from the experiment above was analyzed by CI mass spectrometry. The most relevant signals appeared as triads of signals at m/z 469, 467, and 465 ($\text{MH} + 4$, $\text{MH} + 2$, and MH) and at m/z 265, 263, and 261 ($\text{TipCHC}^{18}\text{O}_2\text{H}$, $\text{TipCHC}^{18}\text{O}^{16}\text{OH}$, and $\text{TipCHC}^{16}\text{O}_2\text{H}$). In both cases the ratio of intensities of the $^{18}\text{O}_2$ species to the $^{18}\text{O}^{16}\text{O}$ species was 64:100.¹³

The reversibility of hydration is not restricted to 1. The hydration of three other ketenes 4–6 with 60-fold excess H_2^{18}O in THF (conditions: 30 min at -18°C followed by 24 h at room temperature under argon) was also studied, and the relative abundances of the appropriate signals are given in Table 1.



Oxygen exchange of carboxylic acids under conditions related to ours is very slow,¹⁴ and likewise 3 and the acids in Table 1 were found (by CI-MS) not to exchange oxygen under the reaction conditions. Hence, the $^{18}\text{O}_2$ -labeled acids are obtained from the reaction of H_2^{18}O with the exchanged ketene 1- ^{18}O , formed by the reversible hydration. Since the unlabeled acid arises both from the reaction of unlabeled ketene with H_2^{16}O and from traces of the acid in the original ketenes, no significance is ascribed to its relative percentage.

Two interesting results arise from Table 1. First, even the hydration of the more reactive ketene, diphenylketene, whose


(13) The $M + 1$ signals of the three isotopic signals are not always in the expected intensity to the M signal based on the 1.1% ^{13}C isotope. $M - 1$ peaks may also contribute to these intensities. Hence, we regard our results only as qualitative.

(14) Samuel, D.; Silver, B. L. *Adv. Phys. Org. Chem.* **1965**, 3, 123.

reactions were extensively investigated, is reversible. Second, the extent of exchange qualitatively increases¹³ with increased stability of the intermediate enediol,^{1,12,15} which in turn seems to increase with the increased bulk of the ketene substituents.

The observation of exchange is relevant to the argument whether the mechanism of neutral hydrolysis of ketene proceeds via $\text{C}=\text{C}$ or $\text{C}=\text{O}$ hydration.^{2,3} Whereas formation of 2 and the 1,1-enediols derived from 4 and 5^{12,15} during the hydration suggest that the reaction proceeds via the 1,1-enediol, no 1,1-enediol intermediate was observed with diphenylketene, and the reversibility is the only evidence for its intermediacy.

A relevant observation is the reaction of a >90% enediol 2 solution (prepared by eq 1) with diazomethane at room temperature. Neither $\text{Tip}_2\text{C}=\text{C}(\text{OMe})_2$ nor $\text{Tip}_2\text{CHCO}_2\text{Me}$ was isolated. Instead a nearly quantitative yield of the conjugated bicyclic trienone 7 was obtained.¹⁶ Since independent reaction of 1 with diazomethane also gave 7, we believe that the reversibly formed ketene from 2 reacts at its double bond to give arylcyclopropanone 8,¹⁷ which undergoes a vinylcyclopropanone rearrangement involving one of the aromatic double bonds of 8 to give 7 (eq 3).



The results described above strongly indicate that ketene hydration is reversible. Further studies on the scope of this reaction are in progress.

Acknowledgment. The mass spectral analysis was conducted in the Mass Spectrometry Center at the Technion, Haifa, Israel. We are indebted to Prof. T. T. Tidwell for bringing ref 4c to our attention and to Professors H. H. Wasserman and J. A. Berson for comments regarding the 8 → 7 reaction.

JA943230K

(15) Frey, J.; Rappoport, Z. Unpublished results.

(16) The structure of 7 was corroborated by X-ray crystallography. We know of no precedent of such a facile rearrangement which is accompanied by a loss of aromaticity.

(17) For reaction of ketene with diazomethane, see: Turro, N. J.; Hammond, W. B. *Tetrahedron* **1968**, 24, 6017.