

## Standardised Tethering of Ru<sub>3</sub>—Ru<sub>6</sub> Clusters to High Surface Area Oxides

John Evans\* and Benjamin P. Gracey

*Department of Chemistry, The University, Southampton SO9 5NH, U.K.*

The complexes Ru<sub>3</sub>(CO)<sub>11</sub>L, H<sub>4</sub>Ru<sub>4</sub>(CO)<sub>11</sub>L, Ru<sub>5</sub>C(CO)<sub>14</sub>L, and Ru<sub>6</sub>C(CO)<sub>16</sub>L [L = PPh<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>Si(OEt)<sub>3</sub>] have been isolated and used to tether Ru<sub>3–6</sub> units to oxide surfaces (SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub>, and TiO<sub>2</sub>) in a standard manner.


Particle-size effects on the catalytic activity of materials derived from oxide-supported metal carbonyl clusters have been suggested.<sup>1</sup> However, testing this suggestion has not been possible since the designed tethering of clusters of differing nuclearity by a standard link has not been reported. By combining radical anion catalysed substitution reactions<sup>2</sup> with purification of complexes of the anchoring ligand PPh<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>Si(OEt)<sub>3</sub> (1) by flash chromatography<sup>3</sup> we have developed molecularity-specific routes to supported ruthenium clusters.




(1)




(2)



(3)



(4)



(5)

Ru<sub>3</sub>(CO)<sub>11</sub>{PPh<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>Si(OEt)<sub>3</sub>} (2) and H<sub>4</sub>Ru<sub>4</sub>(CO)<sub>11</sub>{PPh<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>Si(OEt)<sub>3</sub>} (3) were prepared by the Na–Ph<sub>2</sub>CO catalysed substitution of stoichiometric mixtures of (1) with Ru<sub>3</sub>(CO)<sub>12</sub> and H<sub>4</sub>Ru<sub>4</sub>(CO)<sub>12</sub> respectively in solution in tetrahydrofuran (thf). Ru<sub>5</sub>C(CO)<sub>14</sub>{PPh<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>Si(OEt)<sub>3</sub>} (4) and Ru<sub>6</sub>C(CO)<sub>16</sub>{PPh<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>Si(OEt)<sub>3</sub>} (5) were obtained by controlled substitution in solution by (1) into Ru<sub>5</sub>C(CO)<sub>15</sub> (at –40 °C) and Ru<sub>6</sub>C(CO)<sub>17</sub> (at 0 °C) respectively in methylene dichloride. All four complexes were purified by flash chromatography on Nagel Silica Gel 60 using light petroleum (b.p. 40–60 °C)–methylene dichloride solvent mixtures with the column pressurised by nitrogen (*ca.* 10 lb in<sup>–2</sup>)†. We have found this technique to have wide applicability in cluster chemistry. The <sup>1</sup>H n.m.r. spectra of complexes (2)–(5) contained the expected ligand resonances: *e.g.* (2) in CDCl<sub>3</sub>, δ 7.45 (m, 10H, Ph), 3.77 [q, *J*(HH) 7 Hz, 6H, O–CH<sub>2</sub>–], 2.54 (m, 2H, P–CH<sub>2</sub>–), 1.10 [t, *J*(HH) 7 Hz, 9H, O–CH<sub>2</sub>–CH<sub>3</sub>], and 0.48 (m, 2H, Si–CH<sub>2</sub>). Satisfactory elemental analyses and <sup>31</sup>P n.m.r.

† *E.g.* for complex (2): a slurry on silica gel was formed by removing the thf under reduced pressure and added to the top of a light petroleum–silica column. After elution of Ru<sub>3</sub>(CO)<sub>12</sub>, increasingly polar CH<sub>2</sub>Cl<sub>2</sub>–light petroleum solvent mixtures were used. Complex (2) can be separated with 20–30% CH<sub>2</sub>Cl<sub>2</sub> with the disubstituted derivative requiring a 40% CH<sub>2</sub>Cl<sub>2</sub> level.



Figure 1. I.r. spectra (2200—1900  $\text{cm}^{-1}$ ) of (2): (a) in cyclohexane, and (b) on alumina.




Figure 2. I.r. spectra (2200—1900  $\text{cm}^{-1}$ ) of (3): (a) in cyclohexane, and (b) on alumina.





Figure 3. I.r. spectra (2200—1900  $\text{cm}^{-1}$ ) of (4): (a) in cyclohexane, and (b) on alumina.

spectra were also obtained for these derivatives and the formulation of (2) was confirmed by a mass spectrum obtained by fast xenon atom bombardment of a solution in dipentylphenol. This included a group of peaks centred at 990 a.m.u. due to  $(M + H)^+$  as the highest masses present.

Complexes (2)–(5) were interacted with silica (Aerosil), alumina (aluminium oxide C), and titania (P25) in cyclohexane or  $\text{CH}_2\text{Cl}_2$  at room temperature for 4 days. These reactions were carried out under a carbon monoxide atmosphere to prevent decomposition. The oxides obtained exhibited the colours of the initial complexes and close agreement was found between the band maxima of their electronic spectra obtained by diffuse reflectance with those of the appropriate precursor complex in cyclohexane solution.<sup>‡</sup> Further strong evidence for the integrity of the supported clusters was obtained from their i.r. carbonyl fingerprints. Spectra for the  $\text{Ru}_3(\text{CO})_{11}\text{L}$ ,  $\text{H}_4\text{Ru}_4(\text{CO})_{11}\text{L}$ , and  $\text{Ru}_5\text{C}(\text{CO})_{14}\text{L}$  species [ $\text{L} = \text{PPh}_2\text{CH}_2\text{CH}_2\text{Si}(\text{OEt})_3$ ] in cyclohexane solution and on alumina (as a Nujol mull) are shown in Figures 1–3. The spectra of  $\text{Ru}_5\text{C}(\text{CO})_{16}\text{L}$  on the oxides were similar to those obtained by the reaction of  $\text{Ru}_5\text{C}(\text{CO})_{17}$  with phosphinated silica.<sup>4</sup> Probable molecular structures for the supported complexes are shown in Figure 4; these have been established by X-ray diffraction studies on the close analogues  $\text{Ru}_3(\text{CO})_{11}^-$

<sup>‡</sup> Electronic spectra ( $\lambda_{\text{max}}$ ), (2) in cyclohexane: 420, 300sh, 250sh, and 208 nm; on silica: 424, 376, 288sh, 244sh, and 208 nm; (3) in cyclohexane: 370, 300 w, sh, 240sh, and 210 nm; on silica: 364, 248, and 216 nm; (4) in cyclohexane: 530, 400 w, sh, 340sh, 270w, sh, and 208 nm; on silica: 530, 396sh, 336sh, and 220 nm; (5) in cyclohexane: 520w, sh, 420, 310sh, and 210 nm; on silica: 520w, 420, 308sh, and 210 nm.



**Figure 4.** Probable structures of the supported Ru<sub>3</sub>–Ru<sub>6</sub> cluster units.

(PPh<sub>3</sub>)<sub>5</sub>,<sup>5</sup> H<sub>4</sub>Ru<sub>4</sub>(CO)<sub>11</sub>{P(OMe)<sub>3</sub>}<sub>3</sub>,<sup>6</sup> Ru<sub>5</sub>C(CO)<sub>14</sub>(PPh<sub>3</sub>)<sub>7</sub>,<sup>7</sup> and Ru<sub>6</sub>C(CO)<sub>16</sub>(PPh<sub>2</sub>Et)<sub>4</sub>.<sup>4</sup>

Previous reports of tethering metal carbonyl clusters to oxide supports by means of an anchoring ligand have been isolated to particular systems for which specific syntheses had been devised, *e.g.* Os<sub>3</sub>(CO)<sub>11</sub>(PPh<sub>2</sub>R),<sup>8</sup> H<sub>4</sub>AuOs<sub>3</sub>(CO)<sub>10</sub>(PPh<sub>2</sub>R),<sup>9</sup> Ru<sub>6</sub>C(CO)<sub>16</sub>(PPh<sub>2</sub>R),<sup>4</sup> and Ru<sub>3</sub>(CO)<sub>9</sub>(PPh<sub>2</sub>R)<sub>3</sub>.<sup>10</sup> This report shows that procedures can be developed to synthesise series of

anchored complexes of different nuclearities with standardised metal–support links. Anchoring of preformed complexes like (2)–(5) by hydrolysis of the silyl group in the phosphine ligand (1) has been demonstrated by <sup>31</sup>P n.m.r. spectroscopy to yield more specific materials than by complexing the metals on to functionalised oxides.<sup>11</sup> We are now investigating the chemistry of this series of materials to probe for particle-size effects.

We thank the S.E.R.C. for a studentship (to B. P. G.), Mrs J. M. Street for the n.m.r. spectra, Mr R. Bedder of VG Update Limited for the fast atom bombardment mass spectrum, and Degussa Limited for the oxides.

Received, 8th November 1982; Com. 1266

## References

- 1 M. Ichikawa, *J. Chem. Soc., Chem. Commun.*, 1978, 566.
- 2 M. I. Bruce, D. C. Kehoe, J. G. Matison, P. H. Rieger, and M. L. Williams, *J. Chem. Soc., Chem. Commun.*, 1982, 442.
- 3 W. C. Still, M. Kahn, and A. Mitra, *J. Org. Chem.*, 1978, **43**, 2923.
- 4 S. C. Brown, J. Evans, and M. Webster, *J. Chem. Soc., Dalton Trans.*, 1981, 2263.
- 5 E. J. Forbes, N. Goodhand, D. L. Jones, and T. A. Hamor, *J. Organomet. Chem.*, 1979, **182**, 143.
- 6 R. D. Wilson, S. M. Wu, R. A. Love, and R. Bau, *Inorg. Chem.*, 1978, **17**, 1271.
- 7 D. H. Farrar, P. F. Jackson, B. F. G. Johnson, J. Lewis, J. N. Nicholls, and M. McPartlin, *J. Chem. Soc., Chem. Commun.*, 1981, 415.
- 8 S. C. Brown and J. Evans, *J. Chem. Soc., Chem. Commun.*, 1978, 1063.
- 9 R. Pierantozzi, K. J. McQuade, B. C. Gates, M. Wolf, H. Knözinger, and W. Ruhmann, *J. Am. Chem. Soc.*, 1979, **101**, 5436.
- 10 D. K. Liu and M. Wrighton, *J. Am. Chem. Soc.*, 1982, **104**, 898.
- 11 L. Bemi, H. C. Clark, J. A. Davies, D. Drexler, C. A. Fyfe, and R. Wasylshen, *J. Organomet. Chem.*, 1982, **224**, C5; L. Bemi, H. C. Clark, J. A. Davies, C. A. Fyfe, and R. Wasylshen, *J. Am. Chem. Soc.*, 1982, **104**, 438.