

Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

Preparation of 4,6-Diarylindazole Derivatives in Ionic Liquid under Solvent-free Conditions

Javad Safaei-Ghomi ^a, Mahdi Taheri ^a & Mohammad Ali Ghasemzadeh ^a

^a Department of Organic Chemistry, Faculty of Chemistry, University of Kashan, Kashan, I. R. Iran

Published online: 01 Oct 2010.

To cite this article: Javad Safaei-Ghomi, Mahdi Taheri & Mohammad Ali Ghasemzadeh (2010) Preparation of 4,6-Diarylindazole Derivatives in Ionic Liquid under Solvent-free Conditions, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 42:5, 485-489, DOI: [10.1080/00304948.2010.514795](https://doi.org/10.1080/00304948.2010.514795)

To link to this article: <http://dx.doi.org/10.1080/00304948.2010.514795>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

OPPI BRIEFS

Preparation of 4,6-Diarylindazole Derivatives in Ionic Liquid under Solvent-free Conditions

Javad Safaei-Ghomi, Mahdi Taheri,
and Mohammad Ali Ghasemzadeh

Department of Organic Chemistry, Faculty of Chemistry, University of Kashan,
Kashan, I. R. Iran

Ionic liquids have been the subject of considerable current interest as environmentally benign reaction media in organic synthesis because of their unique properties of non-volatility, non-flammability, recyclability, and ability to dissolve a wide range of organic and inorganic compounds.^{1–3} Ionic liquids are also thermally stable, have negligible vapor pressure and increased reaction rates, show selectivity and a tendency to immobilize starting materials and catalysts. Ionic liquids can play a significant role in controlling reactions as solvents or catalysts⁴ and have been applied in various organic reactions,^{5,6} such as the Friedel-Crafts,⁷ the Diels-Alder^{8,9} and Biginelli reactions,¹⁰ the Beckmann rearrangement,^{11,12} Michael addition¹³ and other reactions.^{14–16}


Among the new transformations, cyclocondensation reactions are attractive methodologies for the synthesis of heterocyclic compounds.¹⁷ Indazole derivatives exhibit anti-inflammatory,¹⁸ anti-depressant,¹⁹ anti-arthritic,²⁰ anti-tumor²¹ and analgesic²² activities. Most of the methods to prepare indazoles proceed from benzene derivatives on which the pyrazole moiety is attached by ring closure.²¹ We report a convenient synthesis of indazole derivatives from cyclohexenone derivatives in ionic liquid media.

A mixture of chalcone with ethyl acetoacetate and 1-butyl-3-methylimidazoliumbromide ([bmim]Br) was stirred at room temperature for the period of times shown in *Table 1*. The reaction mixture was then extracted with ether and the product was recrystallized from ethanol. The best results were obtained with a ratio of [1:1:0.7] mmols of chalcone, ethyl acetoacetate, [bmim]Br. Electron-withdrawing groups on phenyl rings led to shorter reaction times than electron-donating groups.

The present procedure catalyzed by a simple ionic liquid, [bmim]Br, provides an efficient and general methodology for Robinson Annulations. This method has the advantages

Submitted February 12, 2009.


Address correspondence to Javad Safaei-Ghomi, Department of Organic Chemistry, Faculty of Chemistry, University of Kashan, 51167 Kashan, I. R. Iran. E-mail: safaei@kashanu.ac.ir



a) $\text{R}_1 = \text{R}_2 = \text{H}$; b) $\text{R}_1 = \text{Cl}$, $\text{R}_2 = \text{H}$; c) $\text{R}_1 = \text{H}$, $\text{R}_2 = \text{Me}$; d) $\text{R}_1 = \text{Cl}$, $\text{R}_2 = \text{Me}$; e) $\text{R}_1 = \text{H}$, $\text{R}_2 = \text{Br}$;
f) $\text{R}_1 = \text{Cl}$, $\text{R}_2 = \text{Br}$; g) $\text{R}_1 = \text{H}$, $\text{R}_2 = \text{NO}_2$; h) $\text{R}_1 = \text{Cl}$, $\text{R}_2 = \text{NO}_2$

of shorter reaction times and higher yields compared to another representative method, catalyzed by K_2CO_3 (Table 1).²³

Treatment of cyclohexenones (3a-h) with hydrazine hydrate or phenylhydrazine in [bmim]Br as solvent and catalyst at 90°C for 3.5 h produced indazole derivatives. The results are given in Table 2. The conditions led to shorter reaction times and higher yields compared with another representative method in which acetic acid was used as catalyst (Table 2).²⁵



a) $\text{R} = \text{H}$, $\text{Ar} = \text{Ar}^1 = \text{C}_6\text{H}_5$; b) $\text{R} = \text{H}$, $\text{Ar} = \text{C}_6\text{H}_5$, $\text{Ar}^1 = 4\text{-Cl-C}_6\text{H}_4$; c) $\text{R} = \text{H}$, $\text{Ar} = 4\text{-Me-C}_6\text{H}_4$, $\text{Ar}^1 = \text{C}_6\text{H}_5$;
d) $\text{R} = \text{H}$, $\text{Ar} = 4\text{-Me-C}_6\text{H}_4$, $\text{Ar}^1 = 4\text{-Cl-C}_6\text{H}_4$; e) $\text{R} = \text{H}$, $\text{Ar} = 4\text{-Br-C}_6\text{H}_4$, $\text{Ar}^1 = \text{C}_6\text{H}_5$; f) $\text{R} = \text{H}$, $\text{Ar} = 4\text{-Br-C}_6\text{H}_4$, $\text{Ar}^1 = 4\text{-Cl-C}_6\text{H}_4$; g) $\text{R} = \text{H}$, $\text{Ar} = 4\text{-NO}_2\text{-C}_6\text{H}_4$, $\text{Ar}^1 = \text{C}_6\text{H}_5$; h) $\text{R} = \text{H}$, $\text{Ar} = 4\text{-NO}_2\text{-C}_6\text{H}_4$, $\text{Ar}^1 = 4\text{-Cl-C}_6\text{H}_4$;
i) $\text{R} = \text{Ar} = \text{Ar}^1 = \text{C}_6\text{H}_5$; j) $\text{R} = \text{C}_6\text{H}_5$, $\text{Ar} = \text{C}_6\text{H}_5$, $\text{Ar}^1 = 4\text{-Cl-C}_6\text{H}_4$; k) $\text{R} = \text{C}_6\text{H}_5$, $\text{Ar} = 4\text{-Me-C}_6\text{H}_4$, $\text{Ar}^1 = \text{C}_6\text{H}_5$;
l) $\text{R} = \text{C}_6\text{H}_5$, $\text{Ar} = 4\text{-Me-C}_6\text{H}_4$, $\text{Ar}^1 = 4\text{-Cl-C}_6\text{H}_4$; m) $\text{R} = \text{C}_6\text{H}_5$, $\text{Ar} = 4\text{-Br-C}_6\text{H}_4$, $\text{Ar}^1 = \text{C}_6\text{H}_5$; n) $\text{R} = \text{C}_6\text{H}_5$, $\text{Ar} = 4\text{-Br-C}_6\text{H}_4$, $\text{Ar}^1 = 4\text{-Cl-C}_6\text{H}_4$; o) $\text{R} = \text{C}_6\text{H}_5$, $\text{Ar} = 4\text{-NO}_2\text{-C}_6\text{H}_4$, $\text{Ar}^1 = \text{C}_6\text{H}_5$; p) $\text{R} = \text{C}_6\text{H}_5$, $\text{Ar} = 4\text{-NO}_2\text{-C}_6\text{H}_4$, $\text{Ar}^1 = 4\text{-Cl-C}_6\text{H}_4$

Table 1
Comparison of Two Methods in Robinson Annulation

Cmpd	mp(°C)	lit. mp. ²³	Stirred in K_2CO_3 ²³		Stirred in [bmim]Br	
			Time (min)	Yield (%)	Time (min)	Yield (%)
3a	109–111	111–112 ²⁴	60	85	35	95
3b	90–91	90–91 ²³	40	80	30	90
3c	141–143	139–142 ²⁴	50	85	35	95
3d	115–117	115–117 ²³	60	80	30	90
3e	153–155	153–155 ²³	30	80	20	90
3f	111–113	110–112 ²³	30	82	15	90
3g	107–109	105–107 ²⁴	20	60	15	85
3h	134–136	134–136 ²³	20	60	10	90

Table 2
Comparison of two Procedures in the Preparation of Indazoles

Cmpd	mp(°C)	lit. mp. ²⁵	Acetic Acid ^{a,25} Yield (%)	[bmim]Br ^b Yield (%)
5a	215–216	215–217	77	85
5b	123–125	123–125	72	80
5c	159–160	158–160	75	85
5d	120–122	120–121	72	83
5e	146–148	146–148	70	80
5f	141–143	141–143	71	78
5g	157–158	157–158	65	78
5h	167–168	167–168	66	80
5i	170–173	170–173	82	85
5j	166–167	167–168	70	78
5k	182–184	182–184	80	85
5l	186–188	186–188	75	83
5m	213–215	213–215	71	75
5n	167–168	167–168	72	78
5o	150–151	150–151	77	85
5p	160–161	161–162	75	83

a) 6 hours in Ref. 25; b) 3.5 hours.

One of the advantages of ionic liquids is their ability to function as a recyclable reaction medium. The catalyst [bmim]Br could be recovered and reused from the reaction medium easily by washing with ether and evaporating the solvent under vacuum.²⁶

Experimental Section

Yields refer to isolated pure products. The products were characterized by comparison of their spectral (IR, ¹H NMR, ¹³C NMR) and melting points with authentic samples. All of the reactions were carried out in a hood with strong ventilation. IR spectra were recorded on a Magna-550 IR spectrophotometer. Spectra of solids were carried out using KBr pellets. ¹H NMR and ¹³C NMR spectra were determined on a Bruker DPX-400 M Hz using [D₆]DMSO as solvent and TMS as internal standard.

Typical Procedure for the Robinson Annulation Reaction in [bmim]Br (3a–h)

A mixture of the chalcone (1mmol), ethyl acetoacetate (130 mg, 1 mmol) and [bmim]Br (153 mg, 0.7 mmol) was stirred at room temperature for the period of time required to complete the reaction (TLC). The reaction mixture was extracted with ether (3 × 10 mL) and separated from the ionic liquid phase. The residual ionic liquid may be reused in the next run if necessary after a 2 h heat treatment under vacuum. The ethereal layer was

evaporated under reduced pressure and the product was recrystallized from ethanol. The spectral data of all products were identical to those reported in a previous paper.²³

Typical Procedure for the Preparation of Indazole Derivatives (5a–p) in [bmim]Br

A mixture of **3a-h** (1 mmol) and [bmim]Br (153 mg, 0.7 mmol) was added to hydrazine hydrate or phenylhydrazine (1.5 mmol). The mixture was refluxed for 3.5 h at 90°C. The reaction mixture was extracted with ether (3×10 mL) and the ionic liquid phase was separated. The residual ionic liquid may be reused in the next run if necessary after a 2 h heat treatment under vacuum. The ethereal layer was evaporated under reduced pressure and the product was recrystallized from ethanol. The spectral data of all products were identical to those reported in a previous paper.²⁵

Acknowledgements

Financial support from the Research Affairs Office of the University of Kashan, Kashan, I. R. Iran is gratefully acknowledged.

References

1. T. Welton, *Chem. Rev.*, **99**, 2071 (1999).
2. P. Wasserscheid and W. Keim, *Angew. Chem. Int. Ed.*, **39**, 3773 (2000).
3. R. Sheldon, *Chem. Commun.*, 2399 (2001).
4. P. Wasserscheid and T. Welton. *Ionic Liquids in Synthesis*. Wiley-VCH Verlag, Stuttgart, Germany, 2002.
5. Q. W. Yao, *Org. Lett.*, **4**, 2197 (2002).
6. A. Arce, M. J. Earle, S. P. Katdare, H. Rodriguez and K. R. Seddon, *Chem. Commun.*, 2548 (2006).
7. A. Stark, B. L. Maclean and R. D. Singer, *J. Chem. Soc., Dalton Trans.*, 63 (1999).
8. T. Fischer, A. Sethi, T. Welton and J. Woolf, *Tetrahedron Lett.*, **40**, 793 (1999).
9. P. Ludley and N. Karodia, *Tetrahedron Lett.*, **42**, 2011 (2001).
10. J. J. Peng and Y. Q. Deng, *Tetrahedron Lett.*, **42**, 5917 (2001).
11. R. X. Ren, L. D. Zueva and W. Ou, *Tetrahedron Lett.*, **42**, 8441 (2001).
12. W. K. Su, Y. Zhang, J. J. Li and P. Li, *Org. Prep. Proc. Int.*, **40**, 543 (2008).
13. B. C. Ranu and S. S. Dey, *Tetrahedron*, **60**, 4183 (2004).
14. A. R. Hajipour, Z. Nasreesfahani and A. E. Ruoho, *Org. Prep. Proc. Int.*, **40**, 385 (2008).
15. A. Zare, A. Hasaninejad, M. Shekouhy and A. R. M. Zare, *Org. Prep. Proc. Int.*, **40**, 457 (2008).
16. J. Safaei-Ghomi and A. R. Hajipour, *J. Chin. Chem. Soc.*, **56**, 416 (2009); *Chem. Abstr.*, **151**, 313164 (2009).
17. A. R. Katritzky and C. W. Rees, *Comprehensive Heterocyclic Chemistry*, Vols. 1–8. Pergamon Press, New York, 1984.

18. M. N. A. Nasr and S. A. Said, *Arch. Pharm.*, **336**, 551 (2003).
19. J. P. Yeu, J. T. Yeh, T. Y. Chen and B. J. Uang, *Synthesis*, 1775 (**2001**).
20. C. D. Sommers, J. M. Thompson, J. A. Guzova, S. L. Bonar, R. K. Rader, S. Mathialagan, N. Venkatraman, V. W. Holway, L. E. Kahn, G. Hu, D. S. Garner, H.-C. Huang, P.-C. Chiang, J. F. Schindler, Y. Hu, D. M. Meyer and N. N. Kishore, *J. Pharmacol. Exp. Ther.*, **330**, 377 (2009).
21. R. Singh and H. Kaur, *Synthesis*, 2471 (**2009**).
22. A. Gomtsyan, E. K. Bayburt, R. G. Schmidt, C. S. Surowy, P. Honore, K. C. Marsh, S. M. Hannick, H. A. McDonald, J. M. Wetter, J. P. Sullivan, M. F. Jarvis, C. R., Faltynek, and C. H. Lee, *J. Med. Chem.*, **51**, 392 (2008).
23. J. Safaei-Ghomi and Z. Alishahi, *Org. Prep. Proc. Int.*, **38**, 417 (2006).
24. A. Garcia-Raso, J. Garcia-Raso, B. Campaner, R. Mestres and J. V. Sinisterra, *Synthesis*, 1037 (**1982**).
25. J. Safaei-Ghomi and Z. Alishahi, *Org. Prep. Proc. Int.*, **39**, 517 (2007).
26. A. Shaabani, E. Soleimani and A. Maleki, *Tetrahedron Lett.*, **47**, 3031 (2006).