

Novel Cycloaddition Reactions of Isocyanides with Diphenylacetylene Using Transition-metal Complexes

By YOSHIO SUZUKI* and TAKEO TAKIZAWA

(*Faculty of Pharmaceutical Sciences, University of Tokyo, Bunkyo-ku, Tokyo, Japan*)

Summary Reaction of tetrakis(aryl isocyanide)nickel complexes (Ia—c) with diphenylacetylene gave the diiminocyclobutenes (IIIa—c); on pyrolysis, the acetylene complex (Va) gave (IIIa) as sole product, while with (Vd) the major product was iminocyclopentadiene (VIId).

to give tri-iminocyclopentene and pyridine derivatives, whereas aryl and alkyl substituted acetylenes were unreactive, probably owing to the lower electron affinity of their triple bonds.¹ As the isocyano-group is isoelectronic with carbon monoxide and reactions of various metal carbonyls with carbon monoxide and reactions of various metal carbonyls with acetylenes are well known,² it was expected that the reactions of isocyanides with diphenyl-

WE previously reported the cycloadditions of isocyanide to acetylenes substituted with electron-withdrawing groups

⁷ R. Hüttel and H. J. Neugebauer, *Tetrahedron Letters*, 1964, 3541; P. M. Maitlis, D. Pollock, M. L. Games, and W. J. Pryde, *Canad. J. Chem.*, 1965, **43**, 470.