


New Synthesis of Pteridines from the Reaction of 6-Amino-1,3-dimethyl-5-nitrosouracil with Phenacylidetriphenylphosphoranes

By KEITARO SENG,*, HASHIME KANAZAWA, and SADA O NISHIGAKI

(Pharmaceutical Institute, School of Medicine, Keio University, 35 Shinanomachi, Shinjuku-ku, Tokyo 160, Japan)

Summary Treatment of 6-amino-1,3-dimethyl-5-nitrosouracil with phenacylidetriphenylphosphoranes gave the corresponding 7-substituted 1,3-dimethyl-lumazines.

A RECENT paper¹ described a new synthesis of purines by the reaction of 6-amino-1,3-dimethyl-5-nitrosouracil (I) with benzyldetriphenylphosphoranes. We now report a new, convenient synthesis of pteridines by treatment of (I) with phenacylidetriphenylphosphoranes.



SCHEME

triphenylphosphoranes (Wittig reagents) seems reasonable, since in the absence of base or Ph₃P no reaction occurred.

(Received, 13th May 1976; Com. 539.)

TABLE

Phenacyl halide	Product ^a	Yield/%
PhCOCH ₂ Br ..	(II)	67
<i>p</i> -MeC ₆ H ₄ COCH ₂ Br ..	(III)	39
<i>p</i> -MeO-C ₆ H ₄ COCH ₂ Br ..	(IV)	55
<i>p</i> -ClC ₆ H ₄ COCH ₂ Br ..	(V)	55
<i>p</i> -PhC ₆ H ₄ COCH ₂ Br ..	(VI)	64

^a None of the products melted below 300 °C.

To a pre-boiled (30 min) suspension of (I) (0.5 mmol), phenacyl bromide (1.5 equiv.) and Ph₃P (1.5 equiv.) in tetrahydrofuran-aqueous NaOH (10%, 0.5 ml) was added and the mixture was refluxed for 30 min. Evaporation *in vacuo*, followed by dilution with ethanol caused the separation of 1,3-dimethyl-7-phenyl-lumazine (II)² in good yield (Scheme). Other substituted phenacyl halides† provided the corresponding pteridines (see Table).‡

This new pteridine synthesis presumably proceeds through the initial formation of the pyrimidine anil§ by a type of Wittig reaction between the nitroso-group of (I) and phenacylidetriphenylphosphoranes and subsequent dehydrative cyclization. The formation of phenacylidene-

† In these instances, prolonged pre-boiling (*ca.* 1–2 h) is necessary for the completion of the formation of corresponding phenacylidene-triphenylphosphonium salts.

‡ Satisfactory analytical and spectral data were obtained for all products.

§ The reaction of diphenylmethyldetriphenylphosphorane or fluorenyldetriphenylphosphorane with nitrosobenzene has been reported to give benzophenone anil and fluorenone anil, respectively (A. Schönberg and K. H. Brosowski, *Chem. Ber.*, 1959, **92**, 2602).

¹ K. Senga, H. Kanazawa, and S. Nishigaki, *J.C.S. Chem. Comm.*, 1976, 155.

² G. P. G. Dick, H. C. S. Wood, and W. R. Logan, *J. Chem. Soc.*, 1956, 2131.