configuration XIII indicate a net charge of 0.20 e on the dioxygen ligand in the bent structure and 0.36 e in the perpendicular structure). These  $Mn^{II}(O_2)$ -type ( $Mn^{II}$ ,  $d^5$ , S = $\frac{3}{2}$  configurations have not been considered in ref 2, although there is no feature of the ESR spectra which in principle is incompatible with this formalism.<sup>30</sup> They also meet the requirement of no spin density on  $O_2$ . If the vast majority of  $Mn^{II}$ complexes are high spin (including Mn(TPP)), there are at least two systems of Mn<sup>11</sup> with a quartet ground state<sup>31,32</sup> including manganese phtalocyanine.<sup>32</sup> We also do not expect more refined calculations (of a double- $\zeta$  basis set quality or the introduction of configuration interaction between configurations of the same symmetry) to give a stationary value of the energy for the first configurations of the  $Mn^{IV}(O_2^{2-})$  type. However, they could modify the relative stabilities of the  $Mn^{II}(O_2)$ - and  $Mn^{III}(O_2^-)$ -type configurations. Such calculations are now in progress.

From Table I, it also appears that for all configurations but one the bent structure is more stable than the perpendicular one. The two structures are of comparable stability (to the precision of the calculations) for the configuration XIII. However, introduction of configuration interaction will mix this configuration with the configurations IV, VII, and X for the bent structure (since IV, VII, X, and XIII belong to the <sup>4</sup>A' symmetry) and with the configurations VI and X for the perpendicular structure (VI, X, and XIII belong to the <sup>4</sup>B<sub>1</sub> symmetry). Since the configurations IV, VII, and X of the bent structure are lower in energy than the configurations VI and X of the perpendicular structure, the same ordering is likely for the corresponding states  ${}^{4}A'$  and  ${}^{4}B_{1}$ . We can therefore predict that the bent structure is more stable than the perpendicular one. This result (opposite to the conclusions of ref 2) is not unexpected since we did not find any stable configuration with dioxygen bonded as a peroxo  $(O_2^{2-})$  ligand. If we compare the energy value of the most stable configuration for each of the two states  ${}^{4}A'$  and  ${}^{4}B_{1}$  (cf. Table I), the stabilization amounts to 33 kcal/mol.

In summary our conclusions are at odds with the interpretation of the experimental results for the  $Mn(TPP)O_2$  system. We look forward to a more detailed experimental study of the system,<sup>33</sup> especially an x-ray structural determination. We are also carrying more refined calculations on this system.

Acknowledgments. Calculations have been carried out at the Centre de Calcul du CNRS in Strasbourg-Cronenbourg. We thank the staff of the centre for their cooperation. We thank Drs. B. M. Hoffman and F. Basolo for preprints of their publications. This work has been supported through the A.T.P. No. 2240 of the CNRS.

#### References and Notes

- (1) C. J. Weschler, B. M. Hoffman, and F. Basolo, J. Am. Chem. Soc., 97, 5278 (1975).
- (2) B. M. Hoffman, C. J. Weschler, and F. Basolo, J. Am. Chem. Soc., . 98, 5473 (1976).
- (3) B. Gonzales, J. Kouba, S. Yee, C. A. Reed, J. F. Kirner, and W. R. Scheidt, J. Am. Chem. Soc., **97,** 3247 (1975).
- (4) The abbreviations used in this paper follow: M, metal; P, dianion of porphyrin; TPP, dianion of tetraphenylporphyrin; OEP, dianion of octaethylporphyrin; acacen, N,N'-ethylenebis(acetylacetonelminato); T(p-Tol)p, dianion of tetra(*p*-tolyl)porphyrin. L. Vaska, Acc. Chem. Res., **9**, 175 (1976).
- (6) R. Guilard, M. Fontesse, P. Fournari, C. Lecomte, and J. Protas, J. Chem. Soc., Chem. Commun., 161 (1976)
- (7) B. Chevrier, Th. Diebold, and R. Weiss, Inorg. Chim. Acta, 19, L57 (1976)
- (8) M. M. Rohmer, M. Barry, A. Dedieu, and A. Veillard, Proc. Int. Symp. At. Mol. Solid-State Theory, Collision Phenom. Comput. Methods, 1977, in oress.
- (9) X-ray crystal structures have been published for the dioxygen adduct of the iron picket fence porphyrin<sup>10</sup> and of several cobalt Schiff bases.<sup>11</sup>
- J. P. Coliman, R. R. Gagne, C. A. Reed, W. T. Robinson, and G. A. Rodley, *Proc. Natl. Acad. Sci. U.S.A.*, **71**, 1326 (1974).
  (11) (a) G. A. Rodley and W. T. Robinson, *Nature (London)*, **235**, 438 (1972); (b) M. Calligaris, G. Nardin, L. Randaccio, and G. Tauzher, *Inorg. Nucl. Chem.*
- Lett., 9, 419 (1973); (c) J. P. Collman, H. Tayaka, B. Winkler, L. Libit, S.

- (12) Ab initio calculations on the Fe(P)O<sub>2</sub>L (L = NH<sub>3</sub>, Im)<sup>13,14</sup> and Co(acacen)O<sub>2</sub>L (L = H<sub>2</sub>O, CN<sup>-</sup>, CO)<sup>15</sup> have shown that the bent structure is more stable than the perpendicular one.
- (13) A. Dedieu, M. M. Rohmer, M. Benard, and A. Veillard, J. Am. Chem. Soc., 98, 3717 (1976).
- (14) A. Dedieu, M. M. Rohmer, and A. Veillard, in "Metal-Ligands Interaction n Organic and Biochemistry", Part 2, B. Pullman and N. Goldblum, Ed., D. Reidel, Dordrecht, 1977, p 101.
- (15) A. Dedieu, M. M. Rohmer, and A. Veillard, J. Am. Chem. Soc., 98, 5789 (1976).
- (16) J. H. Fuhrop, Struct. Bonding, Berlin, 18, 1 (1974).
  (17) S. K. Cheung, C. J. Grimes, J. Wong, and C. A. Reed, J. Am. Chem. Soc., 98, 5028 (1976).
- (18) J. P. Collman, J. I. Brauman, T. R. Halbert, and K. S. Suslick, Proc. Natl. Acad. Sci. U.S.A., 73, 3333 (1976).
- (19) F. Basolo, B. M. Hoffman, and J. A. Ibers, Acc. Chem. Res., 8, 384 (1975).
- (20) A. Dedieu, unpublished results.
- (21) A controversy still exists over the oxidation state of iron in dioxygen adducts of iron porphyrin, either Fe<sup>III</sup> or Fe<sup>III</sup>.<sup>18,22</sup>
  (22) A. Dedieu, M. M. Rohmer, H. Veillard, and A. Veillard, *Bull. Soc. Chim. Belg.*,
- 85, 953 (1976).
- (23) The LCAO-MO-SCF calculations were carried out with the Asterix system of programs<sup>24</sup> using basis set (10,6,4) for Mn, (7,3) for first-row atoms, and (3) for hydrogen contracted to a minimal basis set (except for the 3d functions which are split). The following geometries were used: for the porphyrin ligand the same geometry that was used for Fe(P)O<sub>2</sub> calculations,  $^{13,14}$  i.e., the experimental geometry of the porphyrin ligand in Fe(TPP).<sup>25</sup> Following Day et al.,<sup>26</sup> the Mn atom was chosen out of plane, lying at 0.25 Å above the plane of the porphyrin (see for instance the out-of-plane distances quoted for Mn(TPP).<sup>3</sup> Mn(TPP)(1-Melm)<sup>3</sup> and Mn(TPP)- $(N_3)$ ,  $^{26}$  the latter structure having a strong similarity with a possible bent structure of Mn(TPP)Q<sub>2</sub>). The Mn–O and O–O bond lengths were, respectively, 1.75 and 1.25 Å both in the bent and perpendicular geometries. The porphyrin plane is in the xOy plane with the nitrogen atoms along the x and y axis, the Mn-O<sub>2</sub> plane is in the xOz plane as it is the case in Ti(OEP)-
- (24) M. Benard, A. Dedieu, J. Demuynck, M. M. Rohmer, A. Strich, and A. Veillard, Asterix, a system of programs for the Univac 1110, unpublished work
- J. P. Collman, J. L. Hoard, N. Kim, G. Lang, and C. A. Reed, J. Am. Chem. (25)Soc., 97, 2676 (1975).
- (26) V. W. Day, B. R. Stults, E. L. Tasset, R. S. Marianelli, and L. J. Boucher, Inorg. Nucl. Chem. Lett., 11, 505 (1975).
- (27) The same result was obtained for a calculation in which the O–O bond length was 1.46 Å, i.e., typical of a peroxo ligand.
- (28) If so, the  $\pi_{g}$  antibonding orbitals of O<sub>2</sub> would be stabilized by the interaction with the 3d orbitals of Mn and therefore both doubly occupied. One should then obtain a stationary value of the energy for the configurations corresponding to the  $Mn^{IV}(O_2^{2^-})$  formalism.
- (29) We thank Dr. G. H. Loew for suggesting that we consider the configurations X–XII and XIV
- (30) We also thank her for her helpful comments regarding the formalism of the Mn- $O_2$  molety. (31) S. Lahiry and V. K. Anand, *Chem. Commun.*, 1111 (1971)
- (32) J. F. Kirner, W. Dow, and W. R. Scheidt, Inorg. Chem., 15, 1685 (1976), and references therein.
- (33) Since this paper was first submitted for publication, the authors of ref 2 have now reinvestigated the ESR spectra. Although some disagreement remains, a configuration which was not previously considered and which may correspond to a  $Mn^{II}(Q_2^{-})$  formalism has been taken into account and cannot totally be discounted.<sup>34</sup>
- (34) B. M. Hoffman and F. Basolo, submitted for publication.

### A. Dedieu,\* M. M. Rohmer

E.R. No. 139 du CNRS, Université L. Pasteur 67000 Strasbourg, France Received January 17, 1977

# Preparation of a Tetraphosphine Macrocyclic Ligand

Sir:

The preparation of synthetic macrocycles and their subsequent ligation to various transition metals has been a fruitful area of research.<sup>1</sup> Macrocycles having oxygen, nitrogen, and/or sulfur donor atoms have been prepared and the chemical reactivity of their metal complexes studied.<sup>2</sup> To date, however, there is only one example of a macrocyclic ligand that contains a phosphorus atom<sup>3</sup> and the chemical reactivity of its metal complex is very much controlled by the nitrogen atoms



that are part of the ligating array. The dearth of information about phosphorus-containing macrocycles is surprising since it is well known that phosphines are excellent ligands which impart many interesting properties to their metal complexes.<sup>4</sup> In view of this, we wish to describe the first synthesis of a tetraphosphine macrocycle, 1, and in addition, we present data relating to the strong complexing ability of this macrocycle toward nickel(II).

The synthesis of the macrocycle is outlined in Scheme I. It began with the reaction of  $2^5$  with "Vitride" <sup>6</sup> and then with 3<sup>7</sup> in THF to produce 4<sup>8</sup> in 70% yield (mp 179-81 °C): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.25 (12 H, dd), 1.8 (12 H, m), 2.05 (2 H, m), 4.1 (2 H, m), 7.25 (20 H, m); IR (KBr) τ 3160, 3000, 2950 (C-H), 2000–1800 (arom), 1450 (Ph-P), 1420 (CH<sub>2</sub>-P), 1390 (i-Pr), 1230 (P=O), 1000 (P-O). Reduction of 4 with LiAlH<sub>4</sub> in THF produced an oxygen-sensitive oil displaying characteristics indicative of the tetraphosphine 5: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.7 (14 H, m), 2.4 (1 H, m, <sup>9</sup> 5.8 (1 H, m), <sup>9</sup> 7.4 (20 H, m); IR (CHCl<sub>3</sub>) 7 3065, 3000 (C-H), 2340 (P-H), 1450 (Ph-P). The tetraphosphine 5 was immediately treated with NiCl<sub>2</sub>·6H<sub>2</sub>O in ethanol to produce the stable, crystalline complex  $6^8$  in 85% yield: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.7 (14 H, m), 7.3 (20 H, m);<sup>10</sup> IR (CHCl<sub>3</sub>) τ 3060, 3000 (C-H), 2360 (P-H), 1450 (Ph-P); UV-vis (CHCl<sub>3</sub>)  $\tau_{max}$  425 nm ( $\epsilon$  1000), 375 (1500), 305 (sh, 4000), 280 (19,000). Reaction of 6 first with anhydrous K<sub>2</sub>CO<sub>3</sub> in absolute ethanol and then  $\alpha, \alpha'$ -dibromo-o-xylene produced the desired macrocyclic complex  $7^8$  in 35–40% yield. The spectral properties of 7 are very similar to those displayed by 6 except that the <sup>1</sup>H NMR exhibits new multiplets at 1.8 and 7.5 ppm of four protons each while the IR attests to the disappearance of the P-H stretch and the appearance of a new  $CH_2$ -P stretch at 1415 cm<sup>-1</sup>. In addition, a molecular weight determination by vapor phase osmometry gave an apparent

molecular weight of 762 in CHCl<sub>3</sub> (theoretical, 764) substantiating the conclusion that the template reaction leading to the macrocycle-nickel(II) complex, 7, was indeed successful. This reaction thus represents the first example of a template reaction involving a phosphine ligand at the reaction center.11

Final proof that the desired macrocycle was formed was provided by the reaction of the complex 7 with aqueous NaCN. After workup an air-sensitive oil was obtained in  $\sim$ 50% yield that showed the necessary physical and spectral characteristics of 1:<sup>8</sup> <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.65 (14 H, m), 3.45 (4 H, m), 7.45 (24 H, m); IR (film) 7 3150, 3000, 2940 (C-H), 1450 (Ph-P); mol wt (CHCl<sub>3</sub>) 634.6 (theoretical), 634 (obsd). The production of this material not only proves the structure of the macrocyclic complex 7, but it also provides a means by which the uncomplexed macrocycle is now available for further complexation to other important transition metals.

Once the structure of the macrocycle 1 was secure, we felt justified in examining the physical characteristics of the macrocyclic nickel(II) complexes that were available to better understand the interaction of this unique ligand with the metal. The complexes,  $Ni(mac)X_2$ , were prepared by metathetical reactions of the synthesized chloride and the data proves that all of the complexes studied ( $X^- = BF_4^-$ ,  $Cl^-$ ,  $SCN^-$ ) are 2:1 electrolytes with the nickel ion being in a square-planar array.<sup>12</sup> The macrocycle 1 is thus a strong field ligand, but a comparison of its ligand field strength in relation to other macrocycles is difficult to assess at present since we have been unable to induce strong monodentate ligands to form tetragonal complexes. This implies that the axial site of the complex is relatively inaccessible to nucleophilic reagents either for electronic or steric reasons. We tend to favor an electronic argument founded on the well-known back-bonding capability of a phosphine moiety and coupled with a compression by the constrained macrocycle on the d orbitals of the metal rather than a steric argument involving the axially located phenyl groups. This is partially substantiated by the reaction of cyanide ion with the metal complex; however, the magnitude, validity, and viability of these arguments will have to await future studies.<sup>13</sup>

Acknowledgment. One of us (T.A.D.) thanks the American Hoechst Corporation for financial support.

## **References and Notes**

- (1) D. H. Busch, Helv. Chim. Acta, Fasciculus Extraordinarius Alfred Werner, 174 (1967).
- L. F. Lindoy and D. H. Busch, "Preparative Inorganic Reactions", Vol. 6, W. L. Jolly, Ed., Interscience, New York, N.Y., 1971, p 1. J. Riker-Napier and D. W. Meek, *J. Chem. Soc., Chem. Commun.*, 442 (2)
- (3)(1974)
- L. Vaska, Acc. Chem. Res., 1, 335 (1968); G. Booth in "Organic Phosphorus (4) Compounds", Vol. 1, G. Kosolapoff, and L. Maier, Ed., Wiley-Interscience, New York, N.Y., 1972, Chapter 3A.
- P. Masterlerz, *Rocz. Chem.*, **38**, 61 (1964). R. B. Wetzel and G. L. Kenyon, *J. Org. Chem.*, **39**, 1531 (1974).
- (7) R. B. King, J. C. Cloyd, and P. N. Kapoor, J. Chem. Soc., Perkin Trans. 1, 2226 (1973).
- New compounds gave correct elemental analyses.
- These signals disappear on addition of D<sub>2</sub>O and are assigned to the P-H (9)resonance with  $J_{P-H} = 210$  Hz.
- The P-H resonances were not observable in the NMR spectrum. (10)
- (11) C. S. Kraihanzel, J. Organomet. Chem., 73 (2), 137 (1974).
  (12) All of the complexes, Ni(mac)X<sub>2</sub> where X = BF<sub>4</sub><sup>-</sup>, Cl<sup>-</sup>, SCN<sup>-</sup>, thus far prepared are diamagnetic and 2:1 electrolytes in water. The electronic spectra of all are nearly identical: a d-d transition at ~340 nm ( $\epsilon \approx 1000$ ) and charge-transfer bands at 280 nm ( $\epsilon \approx 2000$ ). The d-d transition is very difficult to observe because of the close proximity of the chargetransfer band
- (13) Attempts to prepare another similar macrocycle by using 1,3-dibromo-propane as alkylating reagent has thus far been unsuccessful. This despite the fact that 6 reacts smoothly with benzyl bromide and methyl iodide to produce the appropriate nickel(II) complexes.

### T. A. DelDonno, William Rosen\*

Department of Chemistry, University of Rhode Island Kingston, Rhode Island 02881 Received August 1, 1977