

Selective *ortho*-Substitution of Some Phenolic Compounds

By T. M. Cresp and M. V. Sargent*


(Department of Organic Chemistry, University of Western Australia, Nedlands, Western Australia)

and J. A. Elix


(Chemistry Department, Australian National University, Canberra, Australia)

Summary Formylation and bromination of titanium tetrachloride complexes of some *o*-hydroxycarbonyl compounds occurs, to a large extent, at the 3-position.

We have found that titanium tetrachloride exerts a profound effect on the direction of electrophilic substitution in *o*-hydroxycarbonyl compounds. Bromination of the com-



(1)



(2)

a; R¹ = OMe, R² = CO₂Me

b; R¹ = OH, R² = CO₂Me

c; R¹ = OMe, R² = CO Me


d; R¹ = Me, R² = CO₂Me

TABLE

Substrate	TiCl ₄ bromination ^{a, b}	Formylation ^b
(1a)	48% 3-Br 41% 5-Br	48% 3-formyl 8% 5-formyl
(1b)	36% 3-Br 30% 5-Br 12% 3,5-Br ₂ ^c	38% 3-formyl
(1c)	65% 3-Br 15% 5-Br	32% 3-formyl 2% 5-formyl
(1d)	32% 3-Br 48% 5-Br 4% 3,5-Br ₂ ^c	32% 3-formyl 5% 5-formyl
(2)	40% 3-Br 40% 5-Br	49% 3-formyl 5% 5-formyl

^a Bromination without TiCl₄ gives only the 5-bromo-compound. Brominations with TiCl₄ were carried out by addition of TiCl₄ (12.5 mmol) to a stirred solution of the substrate (5 mmol) in CH₂Cl₂ (10 ml), followed by Br₂ (5 mmol). The solution was stirred for 5 min, then poured on to ice and water and stirred rapidly and then worked-up in the usual way. ^b % yields given. ^c Starting material was also isolated.

pounds in the Table gives exclusively the 5-bromo-compound. Under identical conditions except for the presence of an excess of titanium tetrachloride appreciable amounts of the 3-bromo-compounds result. Since < 1 mol. equiv.



of titanium tetrachloride reduces the amount of 3-bromination, a 1:1 complex of the *o*-hydroxycarbonyl compounds, such as (3), is clearly implicated.

On formylation of these compounds using 1,1-dichlorodimethyl ether and an excess of titanium tetrachloride¹ the 3-formyl compounds are the major products. However formylation of compounds (4)—(6) occurs solely at the 5-position thus providing further evidence for the intermediacy of a complex such as (3).

Collman² has observed that transition-metal complexes of β -diketones undergo electrophilic substitution in a manner analogous to aromatic compounds. The occurrence of substitution at the 3-position in titanium tetrachloride complexes of *o*-hydroxycarbonyl compounds points to a degree of bond fixation in the benzene ring of these complexes, and by analogy with Collman's work, they may be regarded as having naphthalene-like character. Thus the metal chelate ring system behaves as a fused aromatic nucleus such that the 3-position constitutes the most reactive α -position of this naphthalenoid system.

This synthetic method is of obvious utility in the preparation of otherwise inaccessible compounds.

We thank Dr. J. R. Cannon for discussions.

(Received, 22nd November 1971; Com. 2005.)

¹ H. Gross, A. Rieche, and G. Matthey, *Chem. Ber.*, 1963, **96**, 308.

² J. P. Collman, *Angew. Chem.*, 1965, **77**, 154.